Saskatchewan
Agriculture, Food
and Rural
Revitalization

Introduction

Weed Control

2005 GUIDE TO
Crop Protection
SASKATCHEWAN
weeds, plant diseases, insects

Foliar Fungicides

Seed Treatments

This publication is only a guide.
Always refer to the product label for
application details and precautions.

Insect Control

Saskatchewan Agriculture, Food and Rural Revitalization

Agriculture Knowledge Centre 1-866-457-2377

The Agriculture Knowledge Centre is your place to start for the latest answers on topics ranging from crops and livestock to nutrient management and the economic implications of management decisions. The toll-free line puts you in touch with resource agents and specialists who draw on expertise from around the province and beyond. The friendly staff will help you find the solutions you need for your operation.

Crop Protection Laboratory (306) 787-8130
(plant disease diagnosis, weed and insect identification, herbicide resistance screening)

The **Crop Protection Laboratory** offers a range of diagnostic services to assist in the identification of problems associated with all agricultural crops and forage. The major areas of expertise include weed, insect and plant disease identification and herbicide resistance screening.

Better background information on a diagnostic sample results in more accurate diagnosis. Therefore, a Crop Protection Laboratory submission form should be completed fully and included with the sample. A printable version of the Crop Protection Laboratory submission form and instructions on sample submission can be found on the SAFRR internet site at www.agr.gov.sk.ca. From the main page follow the links *Crops-Integrated Pest Management-General-Crop Protection Laboratory Services* to locate the form and instructions. Otherwise contact the Crop Protection Lab directly at the number below or the Agriculture Knowledge Centre at 1-866-457-2377 (1-866-45-SAFRR).

Fees:

Insect or weed identification	$10
Visual plant disease identification	$20
Cultured plant disease identification	$40
Herbicide resistance screening	$35 - $100

Phone: (306) 787-8130
E-mail: croplab@agr.gov.sk.ca

Cover photos

Top: Foxtail barley nearing maturity

Middle: Canola stem shattering as a result of sclerotinia infection

Bottom: Bertha armyworm adult moth on canola leaf

Agriculture and Agri-Food Canada

Saskatoon Research Centre	(306) 956-7200
Scott Research Farm	(306) 247-2011
Melfort Research Farm	(306) 752-2776

University of Saskatchewan

Plant Sciences (formerly Crop Science & Plant Ecology)	(306) 966-5006

Shelterbelts

PFRA Shelterbelt Centre Indian Head	(306) 695-2284

Pesticide Analysis
(herbicide, fungicide or insecticide)

Enviro-Test Labs	1-800-668-9878

Additional information

For more information on herbicides, pesticides, pesticide safety and weed, insect or plant disease control, consult the publications available from the Agriculture Knowledge Centre or visit the Saskatchewan Agriculture, Food and Rural Revitalization website: **www.agr.gov.sk.ca.** An update to this publication is available in May.

Major Poison Control Centres

Saskatoon	Regina
Royal University Hospital	Regina General Hospital
(306) 655-1010	**(306) 766-4545**
1-800-363-7474	**1-800-667-4545**

Toxicology Centre

University of Saskatchewan
(306) 966-7441

Spill Report

Saskatchewan Environment
1-800-667-7525

This publication was prepared by Saskatchewan Agriculture, Food and Rural Revitalization in co-operation with Manitoba Agriculture, Food and Rural Initiatives, and with the assistance of the Weed Sub-Council of the Saskatchewan Advisory Council on Soils and Agronomy.

TABLE OF CONTENTS

1. Introduction **9**
How to Use This Book 9
Product Labels and PCP Numbers 9
Safe Use of Herbicides, Fungicides and Insecticides 10
Protecting Yourself from Exposure to Herbicides, Fungicides, and Insecticides 10
Protective Clothing 11
Protecting Your Eyes, Face and Lungs 12
Avoiding Spray Drift 12
Herbicide Efficacy With Low-drift Nozzles 13
Handling a Drift Complaint 13
Mixing Pesticides 13
Container Disposal 14
Sprayer Tank Cleaning 14
How to Identify Crop and Weed Leaf Stages 16

2. Weed Control **19**
Integrated Weed Management 19
Making Spray Decisions 19
Weed Resistance to Herbicides 27
Surfactants and Herbicide Registrations 31
Oil-based Adjuvants and Herbicide Registrations 31
Adjuvants and Your Herbicide 32
Crop and Herbicide Recommendation Tables 32
Weed Control Tables 33
Herbicide Formulations and Companies 44
Products Available as Prepackaged Tank Mixes 46
Herbicide Directory 47
Special Weed Problems 214
Soil Residual Herbicides 219
Effect of Rainfall on Herbicide Efficacy 221

3. Plant Disease Control **222**
Integrated Plant Disease Management 222
Resistance Management 224
Foliar Fungicide Directory 226
Seed Treatment Directory 254

4. Insect Control

4. Insect Control 287
Insect Control Decisions 287
Field Scouting 287
Field Scouting and Insect Management Charts (for individual crops) 289
Insecticides and Bees 287
Insecticide Poisoning in Humans 322
Resistance of Insects to Insecticides 322
Insecticide Directory 324
Insect Control in Stored Grain 351

Reference Material

Pesticide Index 5
Crop Protection Companies .... Inside Cover
Emergency Numbers .... Inside Cover
Agricultural Offices .... Inside Cover
Metric Conversion Factors 7
Tank Mix Chart - Recommended Herbicides .... Insert, Inside Back Cover
Forage Crops Chart - Recommended Herbicides ..Insert, Inside Back Cover

Pesticide Index

Herbicide Directory

a – for use in annual crops
f – for use in forages

2,4-D (a,f) 47
2,4-DB (a,f) 49
Absolute (a) 51
Accent (a) 52
Achieve Liquid (a,f) 54
Achieve Liquid Gold (a) 56
Adrenalin/Adrenalin SC (a) 57
Advance (see 'Trifluralin') 197
Afolan F (see 'Linuron') 146
Ally Toss-N-Go (a,f) 59
Amitrol 240 (a,f) 61
Approve (a) 62
Assert 300 SC (a,f) 63
Assure II (a,f) 65
Atrazine (a) 67
Attain (a) 69
Avadex BW/Avadex MicroActiv (a,f) 70
Avenge 200 C (a,f) 74
Badge (see 'Bromoxynil + MCPA ester') 81
Banvel II (see 'dicamba') (a,f) 88
Basagran/Basagran Forté (a,f) 76
Bonanza (see 'Trifluralin') 197
Bromoxynil (a,f) 79
Bromoxynil + MCPA ester (a,f) 81
Buctril M (see 'Bromoxynil + MCPA ester') 81
Caliber 400 (see '2,4-DB') 49
Casoron 83
Centurion (see 'Clethodim') 84
Clethodim (a,f) 84
Clovitox Plus (see 'MCPB + MCPA') 154
Cobutox 600 (see '2,4-DB') 49
Compitox (see 'Mecoprop') 155
Credit (see 'Glyphosate') 119
Curtail M (a,f) 86
Dicamba (a,f) 88
Dicamba + mecoprop + MCPA (a,f) 90
Dichlorprop + 2,4-D (a) 92
Dichlorprop-D (see 'Dichlorprop + 2,4-D') 92
Dual II Magnum (a) 94
DyVel (a) 95
DyVel DSp (a,f) 96
Eclipse (a) 99
Edge (a,f) 100
Elite (a) 101
Embutox 625 (see '2,4-DB) 49
Eptam 8-E (a) 102
Eradicane 8-E (a) 104
Escort (f) 105
Estaprop (see 'Dichlorprop + 2,4-D') 92
Everest (a) 107
Express Pack/Express TNG (a) 109
Factor (see' Glyphosate') 119
FlaxMax (a) 111
Frontier (a) 112
Frontline (a) 115
Frontline 2,4-D (a) 114
Fusion (a) 117
Glyfos (see 'Glyphosate') 119
Glyphosate (a,f) 119
Gramoxone (a) 129
Grazon (f) 130
Harmony Total (a) 131
Hoe-Grass 284 (a,f) 132
Hoe-Grass II (a,f) 134
Horizon (a) 135
Horizon BTM (a) 137
K2 (a) 138
Kerb 50-WSP (f) 139
Koril 235 (see 'Bromoxynil') 79
Laddok (a) 141
Liberty (a) 142
Liberty 200 SN (a) 145
Linuron (a) 146
Lontrel (a,f) 149
Lorox Toss-N-Go (see 'Linuron') 146
Maverick (see 'Glyphosate') 119
MCPA (a,f) 151
MCPB + MCPA (a,f) 154
Mecoprop (a) 155
Mextrol 400 M (see 'Bromoxynil + MCPA ester') 81
Muster Gold II (a) 156
Muster Toss-N-Go (a) 157
Odyssey (a,f) 159
Oracle (see 'Dicamba') 88
Pardner (see 'Bromoxynil') 79
Pea Pack (a) 160
Pinnacle (a) 161
Poast Ultra (a,f) 162
PrePass (a) 165
Prestige (a,f) 166
Prevail Liquid (a) 168
Primextra II Magnum (a) 169
Princep Nine-T (see 'Simazine') 189
Prism (a) 170
Puma^{120} Super (a) 171
Pursuit (a,f) 173
Pursuit Ultra (a) 175
Refine Extra Toss-N-Go (a,f) 176
Reflex (a) 178
Reglone Desiccant (a) 179
Remedy EC (f) 181
Renegade (see 'Glyphosate') 119
Reward (a) 182
Rival (see 'Trifluralin') 197
Roundup brand products (Dry, Original, Transorb, Ultra, WeatherMax) (see 'Glyphosate') 119
Rustler (a) 183
Select (see 'Clethodim') 84

Sencor (a) ... 184
Shotgun (a) ... 188
Simazine 480 (see 'Simazine') ... 189
Simazine (a,f) ... 189
Spectrum (a) ... 190
Sundance (a) ... 192
Sword (see 'Dicamba + Mecoprop + MCPA') ... 90
Target (see 'Dicamba + Mecoprop + MCPA') ... 90
Thumper (a) ... 194
Topside (See 'MCPB + MCPA') ... 154
Tordon 22K (f) ... 195
Touchdown iQ (see 'Glyphosate') ... 119
Tropotox Plus (see 'MCPB + MCPA') ... 154
Treflan (see 'Trifluralin') ... 197
Trifluralin (Fallow before wheat) (a) ... 197
Trifluralin (Foxtail control in cereals) (a) ... 199
Trifluralin (Broadleaf and grassy weed control) (a,f) ... 201
Triumph Plus (a,f) ... 204
Trophy (a) ... 206
Turboprop 600 (see 'Dichlorprop + 2,4-D') ... 92
Ultim (a) ... 207
Unity (a) ... 209
Vantage/Vantage Plus (see 'Glyphosate') ... 119
Velpar DF (a) ... 211
Venture L (a,f) ... 212

Foliar Fungicide Directory

Acrobat MZ ... 228
Allegro 500F ... 229
Bravo 500 ... 230
Copper 53W (see 'Copper-based Fungicides') ... 231
Copper Spray (see 'Copper-based Fungicides') ... 231
Copper-based Fungicides ... 231
Curzate 60 DF ... 233
Dithane DG Rainshield NT
(see 'Mancozeb-based Fungicides') ... 241
Gavel 75 DF ... 234
Headline ... 235
Kocide DF (see 'Copper-based Fungicides') ... 231
Kocide 101 (see 'Copper-based Fungicides') ... 231
Kumulus DF ... 238
Lance ... 239
Mancozeb-based Fungicides ... 241
Manzate DF (see 'Mancozeb-based Fungicides') ... 241
Parasol Flowable (see 'Copper-based Fungicides') ... 231
Parasol WP (see 'Copper-based Fungicides') ... 231
Penncozeb 75 DF (see 'Mancozeb-based Fungicides') ... 241
Polyram DF ... 242
Quadris ... 243
Reason 500SC ... 244
Ridomil Gold / Bravo ... 245
Ronilan EG ... 246
Rovral Flo ... 247
Senator 70WP ... 248
Stratego 250 EC ... 249
Tanos 50 DF ... 250
Tattoo C ... 251
Tilt 250E ... 252

Seed Treatment Directory

Agrox B-2 ... 255
Agrox CD (see 'Agrox B-2') ... 255
Admire SPT (see 'Imidacloprid-based Insecticides') ... 272
Allegiance FL (see 'Apron FL') ... 256
Apron FL ... 256
Apron Maxx RTA ... 258
Baytan 30 ... 259
Captan Flowable ... 260
Charter ... 261
Crown ... 262
Cruiser 5FS ... 263
DB-Red L ... 264
DCT ... 265
Dithane F-45 ... 266
Dividend XL RTA ... 266
Foundation Lite ... 268
Gaucho CS FL ... 269
Gaucho 480 FL (see 'Gaucho CS FL') ... 269
Genesis 240 (see 'Imidacloprid-based Insecticides') ... 272
Helix ... 271
Helix XTra (see 'Helix') ... 271
Imidacloprid-based Insecticides ... 272
Manco Plus (see 'Mancozeb-based Fungicides') ... 273
Mancozeb-based Fungicides ... 273
Maxim 480 FS ... 274
Maxim PSP ... 275
Mertect SC ... 276
Polyram 16D ... 277
Potato ST 16 (see 'Mancozeb-based Fungicides') ... 273
Poncho 600 FS (see 'Prosper FL') ... 278
Prosper FL ... 278
Raxil 250 FL (see 'Raxil FL') ... 279
Raxil FL ... 279
Senator PSPT ... 281
Thiram 75WP ... 282
Tuberseal (see 'Mancozeb-based Fungicides') ... 273
Vitaflo 280 ... 283
Vitavax rs Fungicide ... 285
Vitavax Single Solution ... 286

Insecticide Directory

Admire	324
Chlorpyrifos (see 'Lorsban')	337
Cygon	325
Decis	326
Diazinon	328
Dibrom	329
Dipel 2X DF	330
Eco Bran	331
Fulfill 50WG	332
Furadan	333
Guthion	334
Imidan	335
Insecto (see 'Protect-it')	352
Lagon (see 'Cygon')	325
Lannate	336
Lorsban	337
Malathion	339
Malathion Grain Protectant Dust	353
Matador	340
Methoxychlor	342
Monitor	343
Nufos (see 'Lorsban')	337
Orthene	344
Phostoxin	352
Pounce	345
Protect-It	352
Pyrinex (see 'Lorsban')	337
Ripcord	346
Sevin XLR / Sevin XLR Plus	347
Sniper (see 'Guthion')	334
Success 480 SC	348
Tempo 20 WP	353
Thimet	349
Thiodan	350
Thionex (See 'Thiodan')	350

Table 1: Metric Conversion Factors* (Approximate)

Metric Unit	*Metric to Imperial Multiply By*	*Imperial Unit*	*Imperial to Metric Multiply By*	*Metric Unit*
LINEAR				**LINEAR**
centimetre (cm)	x 0.39	inch	x 2.54	centimetre (cm)
AREA				**AREA**
square metre (m^2)	x 1.2	square yard	x 0.84	square metre (m^2)
hectare (ha)	x 2.5	acres	x 0.4	hectare (ha)
VOLUME				**VOLUME**
litre (L)	x 0.22	gallon	x 4.55	litre
PRESSURE				**PRESSURE**
kilopascals (kPa)	x 0.14	psi	x 6.9	kilopascals (kPa)
WEIGHT				**WEIGHT**
gram (g)	x 0.04	oz	x 28.35	gram (g)
kilogram (kg)	x 2.2	lb	x 0.454	kilogram (kg)
AGRICULTURAL				**AGRICULTURAL**
litres per hectare (L/ha)	x 0.089	gallons per acre	x 11.23	litres per hectare (L/ha)
litres per hectare (L/ha)	x 0.357	quarts per acre	x 2.81	litres per hectare (L/ha)
litres per hectare (L/ha)	x 0.71	pints per acre	x 1.41	litres per hectare (L/ha)
millilitres per hectare (mL/ha)	x 0.014	fl.oz per acre	x 70.22	millilitres per hectare (mL/ha)
kilograms per hectare (kg/ha)	x 0.89	lb per acre	x 1.12	kilograms per hectare (kg/ha)
grams per hectare (g/ha)	x 0.014	oz per acre	x 70	grams per hectare (g/ha)

*EXAMPLE: To convert centimetres to inches, multiply by 0.39; conversely, to convert inches to centimetres, multiply by 2.54. CAUTION: Herbicide labels are in metric units only. Conversion between the Metric and Imperial system may result in confusion. It is recommended to use metric units only.

Guide to Crop Protection 2005

For Reference Until December 31, 2005
This publication is updated annually and replaces
the 2004 and previous issues.

Introduction

How to Use This Book

This publication is only a guide. Always refer to the product label for application details and precautions.

The *Guide to Crop Protection* is divided into three sections: (1) Weed Control; (2) Plant Disease Control; and (3) Insect Control.

To use the information in each of these sections, use the following process:

1. Turn to the charts at the beginning of each section. There is a set of charts for weeds, plant diseases and insect control. Select the chart for the crop you want or plan to grow. Use the chart to match your weed, disease or insect problems with the products available for that crop.
2. Once you have narrowed your product choices down to a few candidates, go to the recommendation section for that product. Products are listed alphabetically. Read the recommendations thoroughly for each product you are considering.
3. Read the product label attached to the container for detailed instructions on application.

This publication is intended to be used as a guide only. Information contained herein is that available at time of printing. While every effort has been made to ensure accuracy, the provincial government does not accept responsibility for label changes. When more than one trade name is listed, not all weeds or tank mixes may appear on all labels. Consult product labels attached to pesticide containers for final detailed instructions.

Certain recommendations in this publication are given in quantity of commercial product per acre (mL, L, g or kg/acre). Product labels are given in quantity of product per hectare (mL, L, g or kg/ha). To avoid application errors be sure to read and understand label recommendations.

The *Guide to Crop Protection* includes the most recent recommendations for weed, plant disease and insect control in field and forage crops. These recommendations are based on the uses registered under the Pest Management Regulatory Agency's *Pest Control Products Act*. It is an offence under *The Pest Control Products Act* to apply any chemical in a manner not consistent with the product label. If you have any doubts regarding the instructions in this publication, or on the product label, contact the company representative, your local agricultural office or the Pest Management Regulatory Agency for further advice.

Product Labels and PCP Numbers

On each Product Page you will see a PCP number, so named because it is mandated by the ***P**est **C**ontrol **P**roducts Act*. Under the Act, every pesticide requires a unique identifier – the product's PCP number. That number must also appear on the product's label.

The pesticide label packaged with the product is the authoritative source of information on use of the product and will contain more detailed information than is included in this Guide. Some products have a number of trade names for the same active ingredient. However, each product will have its own PCP number and these appear next to the registrants' names. Users who are seeking more detailed information than is provided in this guide, prior to purchase, can use the PCP number to access a sample product label online through the Pest Management Regulatory Agency's (PMRA) website or they can contact the PMRA Hotline by phone at 1-800-267-6315.

Use this link – http://www.eddenet.pmra-arla.gc.ca/4.0/4.01.asp – to access the Electronic Label: Search and Evaluation (ELSE) site and instructions on how to find labels. The PMRA – ELSE label database can be searched by a product's trade name, active ingredient, company name or registration (PCP) number. Since several products can contain the same active ingredient and there are often several versions of the same label on this database, using the PCP number is the most direct route to finding the label that links to the product page in this Guide. ***There may be some differences between a label found on the package and the sample labels found on the PMRA-ELSE web site so always refer to the packaged product label when applying the product.***

Once the product is located, you may click on its number to view an Adobe Acrobat (PDF) document containing the label and any supplemental registrations. Some of these documents run to many pages but you can use the 'Find' capabilities of the Acrobat Reader plug-in for your browser to jump to specific areas of interest. If you do not have Adobe Acrobat Reader installed on your computer you can download a free version from www.adobe.com

Safe Use of Herbicides, Fungicides and Insecticides

Herbicides, fungicides and insecticides are classified according to the use hazard and risk involved. The categories of hazard are:

- toxicity
- flammability
- explosive potential
- corrosivity

The degree of risk is represented by symbols taken from common traffic sign shapes represented by the stop, caution and yield signs. The signal word for each of the signs is danger (high risk), warning (moderate risk) and caution (low risk). Where the risk is minimal, no designation is required. The label on the container will carry the appropriate signs for the protection of the user. Degree of risk symbols for herbicides, insecticides and fungicides used in field and forage crops are included in the product directory. The symbols are illustrated in Figure 1.

Figure 1. Degree of Risk and Hazard Symbols



LD_{50} values are used to rate the toxicity of pesticides. The LD_{50} refers to the dose of pesticide (in mg per kg of the test animal's body weight) that is lethal to 50 percent of the group of test animals. For example, if a pesticide has an LD_{50} value of 10 mg/kg, and the test animals each weigh 1 kg, then 50 percent of the animals would die if they each ate 10 mg of the pesticide.

Table 2. Oral LD_{50} Values as they relate to the Risk/Hazard Symbols



DANGER POISON
LD_{50} less than 500 mg/kg indicates high toxicity



WARNING POISON
LD_{50} 500-1,000 mg/kg indicates moderate toxicity



CAUTION POISON
LD_{50} 1,000-2,500 mg/kg indicates low toxicity

SYMBOL ABSENT
LD_{50} greater than 2,500 mg/kg indicates very low toxicity

Different types of protective equipment are required for pesticides that differ in toxicity. Special equipment requirements are described on the product label, but in general the following precautions must be taken when using pesticides of different hazard ratings.

Danger Poison - requires goggles, respirator, gloves and skin protection, avoid fumes and spray mist.

Warning Poison - requires goggles, gloves and skin protection, avoid fumes and spray mist.

Caution Poison - requires gloves and skin protection, avoid fumes and spray mist.

The absence of a hazard symbol on a pesticide label indicates low toxicity to mammals. Nevertheless, protective clothing should be worn when using pesticides that do not have a hazard symbol.

Protecting Yourself from Exposure to Herbicides, Fungicides and Insecticides

The use of protective equipment and sound safety procedures will help minimize your exposure to herbicides, fungicides and insecticides. Follow the 10 rules for safe application listed below, and wear the safety equipment recommended.

10 Rules for Safe Application

1. Never smoke or eat while applying pesticides.
2. Avoid inhaling sprays or dusts. Wear protective clothing and a respirator.
3. Sprayer lines carrying chemicals should not enter the operator's cab.
4. Have soap, water and a towel available. Should concentrated product spill on skin, hands, face or eyes, wash immediately.
5. Wash hands and face when leaving the treated area, before break periods, lunch or urination.

6. Bathe or shower and change into clean clothing after working with pesticides. Wash clothing each day before re-use.
7. Call a physician or get the patient to a hospital immediately if symptoms of illness occur during or shortly after pesticide application. Be sure to take along the product label or container.
8. Store pesticides out of reach of children and where there is no chance of contact with human food or livestock feeds. Do not store herbicides with insecticides and avoid cross-contamination. Storage areas should be locked.
9. Keep chemicals in their original containers, never in unmarked containers or bottles used for food or drink.
10. Follow proper container disposal methods. All containers should be triple rinsed or pressure rinsed, punctured to render the container non-reusable, and delivered to designated disposal sites.

Protective Clothing

Wear protective equipment as described in the chart to reduce exposure.

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Coveralls	There are two types of coveralls: disposable and reusable. Disposable coveralls are lightweight and comfortable on warm days. They can be worn for mixing and applying pesticides, then discarded at the day's end. If they become contaminated, they should be discarded at once. The second type of coverall is made of washable fabric and may be reused many times. These fabric coveralls are adequate for use with all but the most highly toxic and concentrated pesticides.	Button (or zip) right up to the neck. Loose coveralls around the neck will suck and blow pesticide in and out of the interior of the coveralls as you bend and move. Wear coveralls over a long-sleeved shirt and pants.
Aprons	When pouring or otherwise handling concentrated pesticides, it makes good sense to wear protection in the form of an apron. The apron protects the front of your body from spills or splashes of the concentrate. The apron should be made of rubber or synthetic liquid-proof material that will resist the solvents used in formulating the pesticide.	Make sure the apron covers your body from your chest to your boots.
Gloves	Protect your hands by wearing chemical-resistant gloves. Neoprene gloves provide the best protection. Natural rubber gloves may be used when handling organo-phosphorus or carbamate pesticides. Be sure that they are designed for use with solvents and pesticides. Never use lined gloves, gloves with wristbands or leather gloves.	Put gloves on and roll up the first inch or two of the cuff. That way when you lift your hands, any liquid on the gloves won't drip down your arms.
Hats	Use a chemical-resistant hat, preferably made of washable plastic. The hat may be a hard hat or made of flexible plastic. In either case, it should have a plastic sweatband. Wash and dry entire hat after each use and before storing. Ordinary baseball caps with cloth sweatbands are dangerous as they absorb the pesticide and recontaminate the forehead each time you wear them. Even small amounts of moderately or slightly toxic pesticides may cause severe skin irritation or other illness if exposure continues for several days.	
Boots	Wear chemical-resistant, unlined boots. These boots are available in a variety of styles and materials. Neoprene boots are the best. Knee-length boots offer greater protection because they extend above the lower end of the apron. Avoid leather or fabric boots and shoes because these will absorb pesticides and cannot be cleaned effectively.	Wear your pant legs outside the top of your boots. This will prevent spills and splashes from running into the boot and onto your leg.

Protecting Your Eyes, Face and Lungs

Wear the following equipment to protect your facial area from exposure:

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Goggles	Chemical-resistant goggles keep your eyes safe from both splashing and, if using dry formulations, dusts or granules. Don't use goggles with cloth or elastic headbands as these will absorb pesticides.	Wear goggles snugly on your face so that the sides of your head are protected from splashes. If you wear glasses, make sure you purchase goggles that fit snugly over them. Never wear contact lenses when working around pesticides.
Respirators	Only NIOSH-approved respirators should be used. Do not exchange parts of different respirators. (For example, do not use a cartridge produced by Company "A" with a respirator produced by Company "B" as the combination may not provide adequate protection to the user). Dust masks are ineffective in protecting against herbicide vapours. Similarly, the filters on tractor cabs are intended to remove dust and are not designed to protect against herbicide vapours or mists. Chemical cartridge respirators are recommended for outdoor use when mixing and applying herbicides.	When carrying out operations, change filters each day. The cartridge should be replaced when chemical odour becomes apparent or when breathing becomes difficult. New cartridges should always be installed at the beginning of the spray season. Prior to commencing work, check the face seal while the respirator is on the wearer's face. Regardless of design, respirators cannot be worn securely by people wearing beards, moustaches or sideburns.
Face Shields	Goggles offer some protection, but frequently full-face protection is advised or required according to the pesticide label. It is especially important to protect your eyes and face when pouring or mixing liquid concentrates. Effective face shields are made of clear plastic.	Since the shield attaches to the hard hat, you can raise or lower it as needed.

Avoiding Spray Drift

To minimize the risk of drift, follow these guidelines:

1. Do not spray in winds above 15 km/h (9 mph).
2. Do not spray under dead calm conditions in early morning, night, or late evening. These are often associated with temperature inversions and the combination of these factors can result in long-distance spray drift (2 km or more). Fog or dust that seems to hang in the air is a good indicator of an inversion.
3. Avoid nozzle pressures above 45 psi (310 kPa) for conventional flat fan tips.
4. Use a minimum of 45 L/acre water for all pesticides unless otherwise specified for the product.
5. Take note of buffer zones identified in the "Restrictions" section of this guide. Do not spray when the wind is blowing towards a nearby sensitive crop, shelterbelt, garden, or water body.
6. Use amine formulations of 2,4-D or MCPA where possible. Use special care when applying volatile herbicides (most herbicides in Group 3 and Group 4, particularly ester formulations). Avoid spraying these products on or immediately before hot days.
7. Ensure that air flow from air assisted sprayers is properly set to minimize airblast rebound and drift for different crop canopies.
8. Operate nozzles at their minimum recommended height. For 80° tips, this is 18" (45 cm), and for 110° tips, this is 12" (35 cm). Orienting nozzles forward allows further height reductions.
9. Special nozzles are now available that create coarse, low-drift sprays. Pre-orifice, Turbo-TeeJet, or venturi-type nozzles are available from a number of manufacturers, and these reduce drift by 50 to 95 percent. (Refer to the section entitled **Herbicide Efficacy with Low-Drift Nozzles**)
10. Consider equipping your sprayer with protective shrouds. A number of different designs are available that can reduce drift between 35 and 75 percent.

For more information on reducing drift refer the Factsheet entitled "Spray Drift – Causes and Solutions" available from your local extension office or at the SAF Website: www.agr.gov.sk.ca

Herbicide Efficacy with Low-drift Nozzles

A number of low-drift nozzles are now available from different suppliers. Well established nozzles, such as the Turbo TeeJet, reduce drift by about 50 percent and provide equivalent efficacy to a standard flat fan nozzle. Newer nozzles ("venturi" types) are best known for their dramatic ability to reduce drift (50 to 95 percent), but information on pesticide efficacy is still scarce. Initial data suggest that these nozzles perform well at conventional carrier volumes, travel speeds, and product rates. Some aspects require special attention:

Pressure: Most venturi-type nozzles require higher pressures to operate properly. Below 40 psi (275 kPa), patterns may deteriorate rapidly resulting in poor overlaps and erratic control. Higher pressures are recommended (60 to 80 psi, or 415 to 550 kPa). Drift potential remains low, even at these high pressures. When using automatic rate controllers, make sure your pressures remain high enough for good nozzle performance.

Water Volume: Droplet size become more important at lower water volumes. Little is known about low-drift nozzle performance at or below 5 gal/acre (23 L/acre). Since low-drift nozzles generate fewer droplets than conventional nozzles, ensure that water volumes are high enough for coverage when using coarse sprays.

Weed Type: Difficult-to-wet weeds, such as wild oats, green foxtail, lamb's-quarters, and cleavers, typically require finer sprays for effective coverage. When using venturi nozzles on these weeds, make sure your pressure is high enough to achieve good coverage. Larger weeds and reduced product rates typically make chemical control more difficult, and these conditions may also reveal some performance differences between nozzles.

Herbicide Type: According to preliminary results, herbicides that belong to herbicide Groups 2, 4, and 9 perform well with venturi nozzles, even at normal pressures (40 psi). Application of herbicides in Groups 1, 6, 8, and 10 may require higher pressures with venturi nozzles to maintain good performance, especially under challenging conditions. Wild oat control may be reduced with the coarsest sprays, even when applied at high pressure.

Check with your chemical representative to see if the manufacturer supports the use of low-drift nozzles with their products.

More information is available in the factsheet "Making Sense of New Nozzle Choices," available from your local extension office or at the SAF Website: www.agr.gov.sk.ca

Handling a Drift Complaint

When spray drift occurs, it is important to take the right steps to resolve the complaint. If you suspect that your crop or property has been damaged because of spray drift, use the following guidelines for resolving the situation.

1. Are you sure that the symptoms or damage you see has been caused by spray drift? Contact your local agricultural office to help determine if the damage is the result of spray drift.
2. Contact the suspected applicator as soon as possible. View the damage with the suspected applicator and determine if that person did, in fact, cause the damage.
3. If the damage was caused by the applicator, determine the extent of the damage and the level of compensation (if any) with the applicator.
4. If the situation cannot be resolved quickly because of disagreements on the extent of damage or level of compensation, contact your local agricultural office to discuss options on how to proceed. Documentation will be required, particularly if insurance companies are involved.
5. The involvement of a private consultant is recommended if documentation is required. Required documentation often includes samples of the damaged plants, photographs, and yield comparisons to determine losses. Your agricultural office can provide you with a list of private consultants in your area.
6. The best approach is to start an open and honest line of communication with the suspected applicator. The majority of drift complaints are resolved quickly and efficiently by communicating with the applicator, without the involvement of outside parties.

Mixing Pesticides

The ability to control a broad range of weeds or other pests in one pass is the advantage that a mix of two or more products allows. If tank mixing is not done in the correct order, the result could be a tank-load of material that may not control the target pests, cause injury to the crop, plug nozzles, or leave an undesirable residue in the tank that will require extensive cleaning. Mistakes like these are costly, could put the user at unnecessary risk of exposure to the products, or create an environmental disposal problem.

To avoid mixing that may result in incompatibilities, **always consult the label of the products that are being used** to learn the correct order. **Remember to add all like components at the same stage of mixing.** The list below is a general rule-of-thumb for mixing pesticides:

1. Fill the spray tank with 1/4 to 3/4 the amount of water required for the application and turn on the sprayer agitation. Check the products that are being used for the correct amount to add. Once agitation has begun, maintain until the tank is emptied.
2. Add any fertilizer or pH adjuster additives to the tank.
3. Add any wettable powders, water dispersible granules (DF, DG, or WDG), or flowable liquid suspensions to the tank. Add dry products slowly to prevent clogged return lines. Allow sprayer to agitate for a few minutes, allowing the product to become completely suspended in the tank, before adding the next component.

4. Shake any containers of liquid pesticide thoroughly **before adding** to ensure they are well mixed.
5. Add any pesticides that are solutions (SN) (i.e. amines and salts)
6. Add emulsifiable concentrates (EC, SC) (i.e. esters)
7. Add any surfactants or other adjuvants.

Remember to always consult the label for compatible mixes and recommended mixing order.

Many pesticides will break down if left in the tank for an extended period. Try not to mix any more than you can spray at one time. If you need to stop spraying for a short time, leave the sprayer agitation running to keep products from settling or separating in the tank.

Container Disposal

Proper disposal of used containers and unused pesticide is important to protecting the environment and preventing the potential contamination of soil and water resources. Rinse all containers prior to disposal to reduce environmental contamination caused by open dumping of unwanted containers. Only mix as much pesticide solution as is needed to treat the desired area.

Triple Rinsing

Triple rinsing renders used pesticide containers (metal, plastic, glass) more than 99.9 percent free of residues, in most cases.

Here are the steps that should be followed:

1. Empty contents of the container into the spray tank and drain in a vertical position for 30 seconds.
2. Add a measured amount of rinse water or other diluent until container is about one-fifth full.
3. Rinse the container thoroughly and pour the rinsate into the spray tank.
4. Repeat the procedure twice (it should take only about 5 minutes in total).
5. Puncture or break triple rinsed containers to render them non-reusable. Paper bags should be rinsed once prior to disposal.

Pressure Rinsing

Also available are pressure rinsers that can rinse all sizes of empty pesticide containers that can be lifted into position over the spray tank. A 30 second rinse with a pressure rinser is convenient and just as effective as triple rinsing. Pressure rinsers are constructed to be thrust into the bottom of a metal can or plastic jug. Holes, situated laterally in the rinser tip, direct water from a pressurized source against the inner sides of the container and effectively wash the residual pesticide into the spray tank. Some farmers have found it convenient to attach a rinser to the pump on their large water storage tank to minimize container handling. Pressure rinsers have the added advantage of rendering containers useless by automatically puncturing them.

Disposal of Containers

Properly rinsed containers should be delivered to a designated pesticide container disposal site. Contact your ag rep, municipal office or weed supervisor for the locations of pesticide container disposal sites in your municipality.

Sprayer Tank Cleaning

When pesticide application is completed each day it is important to empty and clean the sprayer thoroughly to prevent the breakdown of certain pesticides, prevent adhesion of the pesticide to the sprayer, and to maintain the sprayer parts in good condition. Certain pesticides break down very quickly when left in solution, and several pesticide solutions can be corrosive to sprayer parts. Tank cleaning is especially important when changing from one crop to another or from one pesticide to another. Each year several reports are logged of herbicide damage cause by carryover of product residue in the tank. To avoid the risk of contamination, sprayers should be cleaned as soon as possible after application is completed.

Do not clean sprayers where rinsate can run off into ditches or other water bodies, near sensitive plants or shelterbelts, or where other people or animals are likely to walk, to avoid unnecessary exposure to people, animals and the environment.

There are three basic types of rinse solution for cleaning sprayer tanks. Their recipes and basic procedures are outlined below:

The Ammonia Rinse – Fill spray tank and add 1 L of household ammonia (3%) for every 100 L of clean water needed for the rinse and begin agitation. Allow solution to flush through the booms until the boom is completely filled with ammonia solution and top up the tank with water. Circulate the ammonia solution through the tank and pump system for 15 minutes. Flush hoses and booms with ammonia rinse solution again (minimum 5 minutes) before emptying. Remove nozzles and screens and scrub with 0.1 L household ammonia per 10 L clean water and an old toothbrush. Perform clean water rinse to remove ammonia solution prior to next spray load. Some herbicides recommend leaving the ammonia rinse in the tank over night to improve cleaning potential.

The Fresh Water Rinse – The spray tank cleaning should begin and end with a fresh water rinse to remove the majority of potential contaminants prior to the cleansing process or prior to the next round of spraying. Drain the tank of its previous contents and fill the tank with clean water. Open nozzle valves and pump clean water through the booms and hoses. Top up the tank with more clean water and circulate/agitate for at least 10 minutes and empty the tank of waste water. If this is the first rinse after spraying, a high pressure hose could be used to clean residue from all surfaces in the tank. Do not enter the tank during the cleaning process.

The Detergent Rinse – After rinsing with clean water, fill spray tank and add a heavy-duty detergent at 0.25 L per 100 L of water (some suggest a non-ionic surfactant such as Agral 90 or Agsurf at 0.6 L per 100 L of water). Circulate the mixture for a minimum of 5 minutes and spray out through sprayer nozzles. Nozzles and screens are removed and cleaned individually with the same detergent solution in a small container. Soaking in this solution for several hours also helps to loosen any deposits.

The above solutions are just components of the overall sprayer cleaning process. Typical rinse instructions will repeat a combination of one or two or all of these basic rinses. Below we will give some generic rinse instructions utilizing the basic rinses as components of the larger cleaning procedure. Never enter the tank during the cleaning process as some cleansers may release dangerous gases.

Method A –

Drain contents of tank – 1 to 2 X Water Rinse – 2X Ammonia Rinse – 2X Water Rinse (one just prior to the next spraying event)

Products: 2,4-D, Accent, Accord, Ally, Atrazine, Avadex, Banvel II, Dichlorprop+2,4-D (Estaprop, Turboprop, Dichlorprop-D), DyVel, Escort, Everest, Express Pack, FlaxMax, Freedom Gold, Fusion, Harmony Total, MCPA, Muster, Muster Gold, Pinnacle, Poast Ultra, Prism, Pursuit Ultra, Refine Extra, Rustler, Sundance, Tordon 22K, Triumph Plus, Unity.

Method B –

Drain Contents of tank – 2x Water Rinse – 2X Detergent Rinse – 2X Water Rinse

Products: Achieve Liquid*, Achieve Liquid Gold*, Basagran, Buctril M, Centurion, Gramoxone*, HoeGrass, Liberty, Pea Pack, Prevail*, Puma^{120} Super, Reglone Desiccant*, Reward*, Select, Sencor, trifluralin products, Venture*

*Manufacturers of these products recommend adding a non-ionic surfactant such as Agral 90 or Agsurf at 0.6 L per 100 L of water.

Method C -

Drain Contents of tank – Several repetitions of the Water Rinse with nozzles and screens removed and checked for debris.

Products: Amitrol 240, Assure II, Attain, Bladex, Dual II Magnum, Eclipse, glyphosate products, Horizon, Horizon BTM, Prestige.

The above directions are general processes based on the similarities of tank cleaning recommendations between products in each of the herbicide groupings. Always follow the specific instructions on the product label.

Several products in the guide do not have label instructions regarding tank cleaning. In the case of products that have no cleaning recommendations on the label, there are some basic principals that can be applied. Products that are water based formulations can usually be cleaned from spray tanks using Method C above. Products that are formulated as an EC, SC or F (flowable) or use a petroleum based adjuvant should at least use Method B. The detergent breaks down the oil that may be sticking to the side of the tank. Products in Group 2 (most will already have a recommendation) will require the use of Method A. The ammonia in Method A either increases the solubility of the product allowing it to be easily removed from the tank surfaces or neutralizes these products. If the product that is to be cleaned out of the tank is a combination of these elements, use a combination of Methods to clean the tank. In these cases, use a good commercial tank-cleaning product from a recognized source, with both ammonia and detergent as components.

Group 2 compounds can occasionally be trapped on the tank walls by petroleum based formulations or adjuvants when tank mixed with other products, resulting in tank residues. A way to reduce the chance of this occurring is to add detergent at 0.25 L per 100 L to the Ammonia Rinse portion to assist with the breakdown of the petroleum coating so that the ammonia may rid the tank of Group 2 product.

It is very important to clean sprayers immediately after every use. With a more diverse rotation, the likelihood of damage from lack of care increases dramatically.

How to Identify Crop and Weed Leaf Stages

Recognition of plant growth stages is essential for effective weed and disease control. Many herbicides and fungicides are safe on a crop only when applied at a specific growth stage. Similarly, weeds are controlled only when they are at certain growth stages.

For most post-emergent products, growth stages are described by the number of leaves. The following is a description of how to count leaves for staging.

Cereals and Annual Grass Weeds

Manufacturers generally use two different systems of staging for grasses. The minimum stages of application are similar, while the later stages may differ.

Some manufacturers use "Total Leaf Count" stages based on the number of leaves on the entire plant, including tillers or secondary shoots. Most recommendations however, are based on the number of main stem leaves and tillers. Tillers or stools are the secondary shoots or stems of a grass plant. Similar to the branches of a broadleaf plant, tillers will emerge from the axils between the leaf and main shoot. Tillers usually begin to appear at the 3 or 4 leaf stage. When staging a plant in this manner, be sure to identify the tillers first, then count only leaves that originate from the main shoot.


leaf
tiller
main stem
prophyll - a sheath found only on tillers

Figure 2. Leaf Stages of Cereals and Annual Grass Weeds


1
1 leaf
1
2
2 leaves
3
2
1
3 leaves
3
4
2
1
4 leaves


5
4
3
2
T#2
T#1
1
5 main stem leaves, 2 tillers OR 9 total leaves

Figure 3. Leaf Stages of Broadleaf Weeds and Crops


Alternate Leaves
Cotyledon Stage
1 Leaf Stage
2 Leaf Stage
4 Leaf Stage
Opposite Leaves
Cotyledon Stage
2 Leaf Stage
4 Leaf Stage
6 Leaf Stage
Whorled Leaves
Cotyledon Stage
1 Whorl Stage
2 Whorl Stage
3 Whorl Stage

Broadleaf Weeds

Cotyledons - These are the seed leaves that usually emerge above ground. On some plants, such as fababeans, lentils and peas, they stay below the soil surface. Cotyledons are not true leaves and are not counted when determining leaf number. They are a different shape than the true leaves and may dry up and disappear at an early stage.

Alternate leaves - Some plants have one leaf at each node on the stem. The next leaf emerges at the next higher node and extends away from the stem in the opposite direction. These plants (lamb's quarters and wild mustard are good examples) are said to have alternate leaves. To determine the leaf stage, simply count the number of leaves present (Figure 3).

Opposite leaves - Plants with two leaves at each node, one on each side of the stem, are said to have opposite leaves. The next pair of leaves on the next node are rotated about 45° so that they are not directly over the previous pair. Plants with opposite leaves have even-leaf numbers only. When counting, the leaf number progresses from cotyledons to 2 leaf, 4 leaf, etc. These plants generally appear shorter than plants with alternate leaves at a similar leaf stage. **Be sure to count each pair as two leaves**. Hemp nettle is a weed that has opposite leaves (Figure 3).

Whorled leaves - More complex plants like cleavers may have whorled leaves. These plants have three or more leaves at each node on the stem. The leaf number in each whorl may vary, so be sure to count each individual leaf unless the Guide or label recommendation refers to the number of leaf whorls (Figure 3).

Figure 4. Leaf Stages of Certain Special Crops and Forages

Lentil Seedling


5th node
4th node
3rd node
2nd node
Stipule (1st node) may be below soil surface

Field Pea Seedling


4th true leaf
Node 6
3rd true leaf may have several leaflets and tendrils
Node 5
2nd true leaf
Node 4
1st true leaf with stipules
Node 3
Soil surface may be anywhere in this range depending on how deep peas are planted.
Node 2 (2nd scale leaf)
Node 1 (1st scale leaf)
Seed
NB: Scale leaves are usually found below ground but may occur above ground. Do not count as true leaves.

Dry Bean Seedling


2nd trifoliate leaf
1st trifoliate leaf
unifoliate leaf
cotyledon

Forage Legume Seedling


cotyledon
Unifoliate
1
1 Trifoliate
2
1
2 Trifoliate
2
3
1
3 Trifoliate

Stages of Alfalfa, Red Clover and Alsike Clover Leaf Development

Weed Control

The use of in crop herbicides to control weeds is often very important in determining the success or failure of a crop. However, many other practices that can be implemented before and after a herbicide application can help to reduce weed competition. The use of these practices is termed Integrated Weed Management.

Integrated Weed Management

A farming system that utilises an array of inter-dependent cultural, biological and herbicidal weed control practices is implementing Integrated Weed Management (IWM). It is essential that IWM involves an array of tools including the rotation of available herbicide groups, ensuring that weeds are exposed to a diverse range of control mechanisms. The principal aim of IWM is to reduce selection for resistance to any single control agent and to delay or prevent the development of herbicide resistant weeds.

Practising IWM does not mean abandoning chemical weed control, just relying on it less. For example:

- You may decide to choose a normal-height wheat or a tall, viny pea variety for a certain field. These crop selections will compete strongly with weeds, possibly allowing you to skip a spray operation.
- You could insert a short-term forage crop into your crop rotation. Studies show that short-term alfalfa stands can reduce wild oat and green foxtail populations by up to 80 percent the year after breaking.
- Early sown barley may give you enough of a "jump" on the weeds that you can avoid herbicide applications.
- Use of vigorous, high-quality seed, sown shallow, can give you better crop competition than poor-quality or deeply sown crop seed.
- Banding nitrogen near the seed can give your crop an advantage over weeds.

For more information, refer to "Integrated Weed Management: Making it Work on Your Farm" factsheet, available from both Manitoba Agriculture and Food and Saskatchewan Agriculture and Food.

Making Spray Decisions

Field Scouting

Field scouting is an important tool for making informed spray decisions. Accurately assessing the type and number of weeds in the field will help you determine if a spray operation is necessary. The scouting pattern diagram on this page provides a guideline for scouting a field. The entire field should be walked to get a feel for the distribution of the weeds present. A minimum of 20 weed counts should be taken across the field. A smaller number may be used, but be aware that accuracy decreases as the number of counts gets smaller. Count the number of weeds in a 1 m^2 or a 0.25 m^2 area and divide the total number of weeds by the number of counts taken to obtain an average for the field. If using 0.25 m^2 samples, make sure to multiply by four so your average is for a 1 m^2 area.

Some weeds are not distributed uniformly and may be found in patches (for example, Canada thistle) or in low spots. As well, the type and number of weeds found along the field edges may be very different from those found inside the field. These areas should be considered separate from the rest of the field. If possible, patches, low spots, and field borders should be treated separately, as field wide spraying may not be required. Look out for new invading weeds and patches of herbicide-resistant weeds. Herbicide-resistant weeds and new invaders should be controlled, regardless of their number, to prevent them from spreading and becoming a serious control problem. Mapping your field's weed problems will allow you to monitor the spread of weed patches over time and help you assess the effectiveness of your control program.



Yield Losses Caused by Weeds

Knowing the amount of crop yield loss caused by a given weed density will help you decide if a spray operation is required. The tables on the following pages give an indication of the yield loss caused by some of the important grassy weeds.

THESE TABLES SHOULD BE USED ONLY AS A GUIDE. The figures are based on Western Canadian research trials and will not be accurate all of the time. The yield loss values apply only to healthy, well fertilized crops with good stand establishment. Crops that are diseased or emerged unevenly will not compete well with weeds and will suffer larger yield losses than indicated in these tables. The yield loss figures are based on competition from single weed species only. Other weeds, such as wild mustard or Canada thistle, must be controlled if the figures are to be accurate. As well, the tables are based on competition from normal height crops. Semi-dwarf or hybrid varieties may not compete as well with weeds and the figures may not be accurate in these cases.

Table 1. Yield Losses (Percent) in Wheat Caused by Wild Oats.

	Wild Oat Density – Number Per Square Metre																
	1	**2**	**4**	**6**	**8**	**10**	**12**	**14**	**16**	**18**	**20**	**25**	**30**	**35**	**40**	**45**	**50**
Wild Oats are 1 Leaf Stage Ahead of the Crop	1	2	4	6	8	10	12	14	15	17	19	22	26	29	32	34	37
Wild Oats are the Same Leaf Stage as the Crop	1	1	2	4	5	6	7	8	9	10	11	14	16	18	20	22	24
Wild Oats are 1 Leaf Stage Behind the Crop	0	1	1	2	3	3	4	5	5	6	7	8	10	11	13	14	15

Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Figure 1. Spray Decision Guideline for Wild Oats in Wheat.


Is Your Crop Healthy and Growing Well With Good Stand Establishment?
Yes
No
Do Not Use This Guideline. Yield Losses May Be Greater and It May Be Necessary To Spray At Lower Densities Than Indicated Here.
Determine the Leaf Staging of Your Crop and the Wild Oats. Choose One of the Following Three Options
The Wild Oats Are 1 Leaf Stage Ahead of Your Crop
The Wild Oats Are The Same Leaf Stage as Your Crop
The Wild Oats Are 1 Leaf Stage Behind Your Crop
Spray Only if You Have 5 or More Wild Oats Per Square Metre
Spray Only if You Have 8 or More Wild Oats Per Square Metre
Spray Only if You Have 15 or More Wild Oats Per Square Metre

Table 2. Yield Losses (Percent) in Wheat Caused by Green Foxtail (Wild Millet).


Is the Green Foxtail Density Less Than 100 Plants Per Square Metre?
No
Yes
Has the Majority of the Green Foxtail Emerged Within One Week of the Crop?
No
Yield Loss Insignificant
Yes
Are Daytime Temperatures Near the Time of Spraying 20°C or Higher?
No
Yield Loss Insignificant
Yes

Green Foxtail Density (no. per square metre)	50	75	100	125	150	175	200	250	300	350	400	450	500
Yield Loss (percent)	2	3	5	6	7	8	9	11	14	16	18	20	23

Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Table 3. Yield Losses (Percent) in Barley Caused by Wild Oats.

Crop Density (plants/m²)	Relative Emergence	Wild Oat Density (plants/m²)											
		1	5	10	15	20	25	30	40	50	70	100	150
300	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.3	1.4	2.8	4.1	5.4	6.7	8.0	10.3	12.6	16.8	22.4	30.2
	Wild Oats are the Same Leaf Stage as the Crop	0.3	1.3	2.5	3.7	4.8	6.0	7.1	9.2	11.3	15.1	20.3	27.6
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	0.9	1.7	2.6	3.4	4.2	5.0	6.6	8.1	11.0	15.0	20.9
225	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.4	1.9	3.6	5.4	7.0	8.6	10.2	13.1	15.9	20.9	27.4	36.2
	Wild Oats are the Same Leaf Stage as the Crop	0.3	1.6	3.1	4.6	6.1	7.5	8.8	11.4	13.9	18.4	24.4	32.6
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	1.0	2.0	3.0	4.0	4.9	5.8	7.6	9.3	12.6	17.1	23.6
175	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.5	2.3	4.6	6.7	8.7	10.7	12.5	16.1	19.3	25.1	32.3	41.8
	Wild Oats are the Same Leaf Stage as the Crop	0.4	1.9	3.8	5.6	7.3	8.9	10.5	13.6	16.4	21.6	28.2	37.1
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	1.1	2.3	3.4	4.4	5.5	6.5	8.5	10.4	14.0	18.9	25.9

Table 4. Yield Losses (Percent) in Barley Caused by Green Foxtail (Wild Millet).


Is the Green Foxtail Density Less Than 200 Plants Per Square Metre?
No
Yes
Has the Majority of the Green Foxtail Emerged Within One Week of the Crop?
No
Yield Loss Insignificant
Yes
Are Daytime Temperatures Near the Time of Spraying 20°C or Higher?
No
Yield Loss Insignificant
Yes

Green Foxtail Density (no. per square metre)	50	75	100	125	150	175	200	250	300	350	400	450	500
Yield Loss (percent)	1	2	3	3	4	5	6	7	8	10	11	13	14

Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Table 5. Yield Losses (Percent) in Canola Caused by Wild Oats and Volunteer Cereals.

	Weed Density – Number Per Square Metre												
	1	**2**	**4**	**6**	**8**	**10**	**12**	**14**	**16**	**18**	**20**	**25**	**30**
Wild Oats	3	5	6	8	9	10	11	12	13	14	15	16	18
Volunteer Wheat	1	3	6	8	10	11	12	14	15	16	17	19	21
Volunteer Barley	3	5	8	10	12	14	15	17	18	19	20	23	25

Sources: Dew and Keys, Agriculture Canada (Lacombe, Alberta)
O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Figure 2. Spray Decision Guideline for Wild Oats and Volunteer Cereals in Canola.


Is Your Crop Healthy and Growing Well With Good Stand Establishment?
Yes
No
Do Not Use This Guideline. Yield Losses May Be Greater and It May Be Necessary To Spray At Lower Densities Than Indicated Here.
Choose One of the Following Four Options
Wild Oats Only. Spray Only if You Have 1 or More Weeds Per Square Metre
Volunteer Wheat Only. Spray Only if You Have 2 or More Weeds Per Square Metre
Volunteer Barley Only. Spray Only if You Have 1 or More Weeds Per Square Metre
Mixed Population of Grassy Weeds. Spray Only if You Have 1 or More Weeds Per Square Metre

Table 6. Yield Losses (Percent) in Canola Caused by Green Foxtail (Wild Millet)


Is the Green Foxtail Density Less Than 100 Plants Per Square Metre?
No
Yes
Has the Majority of the Green Foxtail Emerged Within One Week of the Crop?
No
Yield Loss Insignificant
Yes
Are Daytime Temperatures Near the Time of Spraying 20°C or Higher?
No
Yield Loss Insignificant
Yes

Green Foxtail Density (no. per square metre)	50	75	100	125	150	175	200	250	300	350	400	450	500
Yield Loss (percent)	3	4	5	6	8	9	10	13	15	18	20	23	25

Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Table 7. **Yield Losses (Percent) in Flax Caused by Wild Oats and Volunteer Cereals.**

	Weed Density – Number Per Square Metre									
	1	**2**	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**10**
Wild Oat	6	8	10	12	13	15	16	17	18	19
Volunteer Wheat	6	11	15	18	22	24	27	29	31	33
Volunteer Barley	6	12	16	21	24	28	31	34	36	39

Sources: Dew and Keys, Agriculture Canada (Lacombe, Alberta)
Friesen et al., University of Manitoba (Winnipeg, Manitoba)

Figure 3. Spray Decision Guideline for Wild Oats and Volunteer Cereals in Flax.


Flax is a Very
Non-Competitive Crop
Consider Spraying for Wild Oats
and Volunteer Cereals When You Have
1
or More Weeds For Every
4
Square Metres in Your Flax Crop

Deciding to Spray – Economic Thresholds and Herbicide Resistance

An **economic threshold** is the level of infestation at which lost yield exceeds the cost of the chemical and its application. Determining the economic threshold will help you decide if a spray operation is necessary.

The following example outlines how to determine an economic threshold:

You have a wild oat problem in your wheat. After a thorough field scouting you have determined that your field has an average density of 35 wild oats per square metre. You know that the crop and weeds are at the same leaf stage. Using Table 1, choose the "Same Leaf Stage" row and read across to **35** wild oats per square metre. You will find that your yield loss will be about **18** percent.

You think it could be a 40 bushel per acre wheat crop, and expect to get $3 per bushel for it. Therefore:

40 bushels x 0.18 (percent of expected yield loss) =
7.2 bushels per acre of lost yield

7.2 bushels x $3 per bushel =
$21.60 per acre of lost income

Now find out the price of your herbicide. Most wild oat herbicides for wheat cost about $15 per acre. In this case, lost income exceeds the cost of the herbicide and application, so spraying would be justified.

Alternatively, you may want to use the figures provided with some of the yield loss tables. These figures provide flowcharts to assist you in making spray decisions. In some cases the flowcharts may indicate to spray when you do not have an economic threshold weed density, but most times they will prevent you from spraying unnecessarily.

Another factor to consider when deciding whether to spray is your herbicide rotation. A one in three rotation of herbicide groups is currently recommended to delay the development of herbicide resistance for weeds such as wild oats and green foxtail. Skipping a spray operation will give you an extra year of flexibility in your herbicide rotation. This means that you have one extra herbicide group to choose from the year after you skipped the spray operation. When making spray decisions, the ability to rotate herbicides should be considered in addition to the economics of spraying.

Making the Spray Decision

Remember that economic thresholds should be used only as guides when making a spray decision. Lost income caused by dockage or downgrading must also be considered. **FIELDS THAT ARE NOT SPRAYED THIS YEAR HAVE A HIGHER POTENTIAL FOR PROBLEMS THE FOLLOWING YEAR BECAUSE OF WEED SEED RETURN.** A farmer's experience and common sense play an important role when deciding to spray. Used properly, however, the economic threshold can be an important tool in making spray decisions.

Weed Resistance to Herbicides

In recent years, the number of herbicide-resistant weeds and the areas they infest have increased.

Most herbicide-resistant weed infestations have developed following repeated use of the same herbicide (or herbicide group) for a number of years on the same field. Growers who have developed weed resistance on their farms will typically see a weed, which is normally controlled by a herbicide, escape uncontrolled after a number of years of use of the same product or product group. Individual plants may be resistant to 1.5 up to 10 or more times the normal field rate.

Herbicide Groups

To help you plan your herbicide program, the following table lists "herbicide groups." To slow down the process of developing weed resistance, use products from different groups from year to year on your fields.

Table 1: Herbicide Groups Based on Mode of Action

Group 1.(contain ACCase grasskillers) Achieve, Assure II, clethodim, Elite*, FlaxMax*, Fusion, Harmony Total*, Hoe-Grass 284/II*, Horizon, Horizon BTM*, Muster Gold*, Poast Ultra, Prevail*, Puma(2) Super, Pursuit Ultra*, Triumph Plus*, Venture
Group 2 (contain ALS/AHAS inhibitors) Absolute*, Accent, Adrenalin*, Ally, Approve, Assert, Champion(2) Extra*, Escort, Everest, Express Pack*, Express TNG, Frontline*, Frontline 2,4-D*, Harmony Total*, K2, Muster, Muster Gold*, Odyssey, Pinnacle, Prism, PrePass*, Pursuit, Pursuit Ultra*, Refine Extra, Spectrum*, Sundance, Triumph Plus*, Trophy*, Ultim, Unity*
Group 3 (contain mitotic inhibitors) Advance, Bonanza, Edge, Rival, Treflan
Group 4 (contain growth regulator herbicides) 2,4-D, 2,4-DB, Absolute*, Adrenalin*, Attain, bromoxynil+MCPA ester*, Curtail M, dicamba, dicamba+mecoprop+MCPA, dichlorporp+2,4-D, DyVel, DyVel DSp, Eclipse*, Elite*, Express Pack*, FlaxMax*, Frontline*, Frontline 2,4-D*, Grazon, Lontrel, MCPA, MCPB+MCPA, mecoprop, Pea Pack*, Prestige, Prevail*, Remedy, Rustler*, Spectrum*, Thumper*, Tordon 22K, Trophy*, Triumph Plus*
Group 5 (contain photosynthetic inhibitors – triazines) Atrazine, Laddok*, Pea Pack*, Primextra II Magnum*, Sencor, simazine, Velpar

Group 6 (contain photosynthetic inhibitors – nitriles/benzothiadiazoles) Approve, Basagran, bromoxynil, bromoxynil+MCPA ester *, Elite*, Hoe-Grass II*, Horizon BTM*, Laddok*, Thumper*, Unity*
Group 7 (contain photosynthetic inhibitors – ureas/amides) linuron
Group 8 (unknown mode of action) Avadex, Avenge, Eptam 8E, Eradicane 8E
Group 9 (contain inhibitors of EPSP synthase) Credit, Eclipse*, Factor, Glyfos, Maverick, PrePass, Renegade, Roundup (Dry, Original, Transorb, Ultra, WeatherMax), Rustler*, Touchdown, Vantage, Vantage Plus
Group 10 (contain inhibitors of glutamine synthetase) Liberty
Group 11 (inhibit carotenoid synthesis - triazoles) Amitrol 240
Group 14 Reflex
Group 15 (inhibit cell division - benzamides, chloroacetamides) Dual II Magnum, Frontier, Kerb, Primextra II Magnum*
Group 22 (membrane rupture, photosynthetic inhibitors) Gramoxone, Reglone, Reward

*Products contain more than one active ingredient and appear in more than one group. In some instances, both active ingredients act to kill the same weed using different modes of action. In these instances, use of tank mixes may slow down the process of developing weed resistance.

New herbicides do not necessarily have a unique mode of action and may fall within the groups listed in the charts. Herbicides that have the same mode of action may not control the same weed spectrum or have the same crop safety. For example, Assert and Ally have the same mode of action; however, Assert controls wild oats while Ally does not.

How Herbicides Work

After applying a herbicide, fields can be scouted to determine the effectiveness of the treatment. The symptoms of different herbicide groups, and the approximate time it takes to develop these symptoms, is listed in the table below. Weed patches that are not affected should be noted and checked, as they may be herbicide resistant. Note that symptoms may take longer to develop when conditions are not conducive to rapid plant growth.

Table 2: The Mode of Action, Site of Uptake and Symptoms of Different Herbicide Groups.

Herbicide Group	Mode of Action	Site of Uptake	Weed symptoms/timing	
			Grass weeds	**Broadleaf weeds**
1	Systemic	Foliar	Reduced growth, yellowing of growing point in 1-3 weeks. Newest leaf of affected plant pulls out easily in 3 to 5 days.	Tolerant
2	Systemic	Foliar/Soil	Newest leaves yellowed in 3-10 days, dead in 1-3 weeks.	Discolored (red/yellow/purple) at the growing point and spreading to the whole plant in 1-3 weeks.
3	Systemic	Soil	Reduced emergence, poor root development of emerged plants. Roots often swollen/stunted and root tips darkened.	Reduced emergence, poor root development of emerged plants.
4	Systemic	Foliar	Yellowed / purpled leaves in 1-2 weeks, plant death in 2-4 weeks.	Abnormal growth (twisted stems, cupped leaves) in 2-10 days.
5	Systemic	Soil	Wilted and yellowed oldest leaves, death in 7-10 days.	Wilted and yellowed oldest leaves, death in 7-10 days.
	Contact	Foliar	Yellowed oldest leaves, death within days."	Bleached oldest leaves, death within days.
6	Contact	Foliar	Some leaf burn possible.	Yellowed leaves in 2-4 days, death in 1-2 weeks.
7	Systemic	Soil	Yellowed and stunted plants, death in 10-14 days.	Yellowed and stunted plants, death in 10-14 days.
	Contact	Foliar	Interveinal yellowing of oldest leaves, death within days.	Interveinal yellowing of oldest leaves, death within days.
8	Contact	Foliar	Yellowed leaves in 3-7 days, stunted plants.	Tolerant
	Systemic	Soil	Reduced emergence, emerged leaves dark green/blue.	Reduced emergence, emerged leaves dark green/blue.
9	Systemic	Foliar	Wilted, yellowed leaves in 7-10 days.	Wilting, yellowing of plant in 7-10 days.
10	Contact	Foliar	Wilted, bleached leaves in 3-5 days, death in 1-2 weeks.	Wilted, bleached leaves in 3-5 days, death in 1-2 weeks.
11	Systemic	Foliar	Plants wilted in 2-3 days, bleached and purpling leaves in 1-2 weeks.	Plants wilted in 2-3 days, bleached leaves in 1-2 weeks. Perennial plants die slowly.
14	Contact	Foliar	Some leaf burn.	Leaves yellowed and dessicated in 1-3 days.
15	Sytemic	Soil	Reduced emergence, emerged plants stunted.	Reduced emergence, emerged plants stunted.
22	Contact	Foliar	Leaves wilted within hours, dessicated in 1-3 days.	Leaves wilted in 1-3 days, dessicated and dead in 3-7 days.

How to Identify Weed Resistance

It is important to avoid confusing herbicide failure caused by resistance with herbicide failure caused by various other factors (such as weather or application errors). When a herbicide fails to control weeds because of weather or application factors, that herbicide may work in the field the next season. But when herbicides fail because of the development of resistance, they will fail in subsequent years, regardless of weather or application procedures.

Herbicide resistance should be suspected under the following conditions:

- A weed species that the herbicide controlled in previous seasons now escapes the treatment, while other weeds that appear on the label continue to be controlled in the field.
- The escapes cannot be attributed to adverse weather or emergence after application (if a post-emergence product is in question).
- Irregular-shaped patches of a weed develop where the herbicide gives little or no control.
- Records of the past history of the field show repeated use of the same herbicide, or combinations of herbicides, that kill the weed in question in the same way.

Table 3: Herbicide-Resistant Weeds in Western Canada

WEED	HERBICIDE GROUP	LOCATIONS CONFIRMED
Canada Fleabane	Group 9	Occurs in several US states
Cleavers	Group 2	AB
	Group 4	AB
	Multiple combinations of:Groups 2 & 4	AB
Chickweed	Group 2	AB
Green Foxtail	Group 1	AB, MB, SK
	Group 2	Occurs in MB & Ontario
	Group 3	AB, MB, SK
	Multiple combinations of: Groups 1 & 3	MB, SK
Hemp-nettle	Group 2	MB
	Group 4	AB
Kochia	Group 2	AB, MB, SK
	Group 4	Occurs in North Dakota and Montana
Lamb's-quarters	Group 2	Occurs in Ontario
	Group 5	Occurs in Ontario
Mustard, Ball	Group 2	AB
Mustard, Wild	Group 2	AB, MB
	Group 4	MB
	Group 5	MB
Persian Darnel	Group 1	Occurs in Montana
Redroot pigweed	Group 2	Occurs in Ontario and North Dakota
	Group 5	Occurs in Ontario
Russian thistle	Group 2	Occurs in SK, North Dakota and Montana
Spiny Annual Sow-thistle	Group 2	AB
Stinkweed	Group 2	AB
Wild oat	Group 1	AB, MB, SK,
	Group 2	AB, MB, SK
	Group 8	AB, MB, SK
	Multiple combinations of: Groups 1 & 2, 1 & 8, 2 & 8, 1, 2 & 8	AB, MB, SK

See Table 1 on page 27 for a complete list of products in each Herbicide Resistance Group.

If Weed Resistance Develops on Your Farm

It is important to identify weed resistance before it spreads across your farm. Plan on conducting a "patch watch" scouting program this summer to identify suspicious patches before they become difficult to manage. Resistant weed patches have been identified on fields where producers were unaware of their existence.

Your patch watch program should begin shortly after spraying and continue through July after the crop has headed out and most weeds are visible from a distance. If you find suspicious looking patches, contact your local agricultural office or crop protection company representative to assist you in confirming weed resistance. If resistance is suspected:

1. Map the location of the patches and mark them with stakes so you will remember their location.
2. Mow, cultivate or spot spray the patches. Resistant patches should not be allowed to produce seed.
3. Patchy areas should NOT be harvested with the rest of the field. Harvest these areas separately, and make sure to clean all harvesting equipment before leaving the area to prevent the spread of seeds across the field or to a neighbouring field.
4. Check patches each year to monitor their spread. Keeping your resistant weeds isolated to a manageable patch is easier than dealing with an entire field of resistant weeds.

Surfactants and Herbicide Registrations:

Registrations listed are strictly those listed on the adjuvant label. For adjuvants recommended for use with a specific product, see the product page.

SURFACTANT	% Active ingredient	TYPE	REGISTERED HERBICDES
Agral 90, Agsurf	90	Non-ionic	Accent, Ally, Escort, Everest, K2, Glyphosate, Gramoxone, Muster, Pinnacle, Pursuit, Prism, Refine Extra, Reflex, Reglone, Reward, Ultim, Unity
Citowett Plus	50	Non-ionic	Ally, atrazine, Basagran (peas), Escort, Muster, Pinnacle, Prism, Refine Extra, Unity
Companion	70	Non-ionic	Ally, Glyphosate, Muster, Refine Extra, Unity
Permax	76	Non-ionic	Ally, Glyphosate, Muster, Refine Extra, Reglone
Super Spreader	50	Non-ionic	Accent, Ally, atrazine, Basagran, Escort, Everest, Muster, Pinnacle, Prism, Refine Extra, Reflex, Ultim, Unity
LI700	80	Non-ionic	Everest, Glyphosate

* Note – some products are specific about the concentration of active ingredient in the surfactant for product performance. Check with the herbicide page in this guide or the product label.

Oil-based Adjuvants and Herbicide Registrations:

TRADE NAME	COMPOSITION	REGISTERED HERBICIDES
Amigo	30% phosphate ester surfactant	Centurion, Select*
Assist	83% paraffin based mineral oil 17% surfactant blend	Basagran (all crops), Laddok
Crop oil or Corn oil	99% paraffin based mineral oil 1% surfactant blend	Atrazine, linuron
Merge	50% solvent 50% surfactant blend	Absolute, Elite*, FlaxMax, Muster Gold II*, Odyssey, Poast Ultra, Sundance*
Score	83% paraffin based mineral oil 17% surfactant blend	Horizon*, Horizon BMT*
Sure-Mix	60% paraffin based mineral oil 35.6% surfactant blend	Assure II*, Muster Gold II*, Refine Extra
Turbocharge	50% mineral oil 39.5% surfactant blend	Achieve Liquid*, Prevail Liquid*
Adapt (XA oil Concentrate)	83% paraffin based mineral oil 17% surfactant blend	Atrazine, Basagran

* The adjuvant is packaged with the product.

Adjuvants and Your Herbicide

Adjuvants are important ingredients in chemical weed control. Many herbicides must be applied with an adjuvant. If it is forgotten, the level of weed control can vary widely, and re-spraying may be necessary.

Most products have adjuvants built into the formulation (e.g. Puma). Other require adjuvant addition (e.g. Refine Extra). Some adjuvants were developed specifically for one herbicide, and these are either pre-packaged with the herbicide, or are identified by name on the label (e.g. Turbocharge for Achieve, Amigo for Select).

With some products, adjuvants need to be added only under certain conditions. For example, glyphosate products have built-in adjuvants, but require additional adjuvant when low rates (pre-seeding or chem-fallow), high water volumes, or certain tank mixes are used.

Adjuvants should be added only when required. If one is not required, addition can reduce weed control or injure crops. Product labels will describe when an adjuvant is required, and what type should be used.

There are two main classes of adjuvants: "activators or spray modifiers" (these include surfactants and crop oils), and "utility modifiers" (these include pH adjusters, water conditioners, low-drift adjuvants, and anti-foaming agents). The most important class of adjuvants is the activators. Surfactants, the main group within the activators, are "*surface active agents.*" These chemicals produce effects at points where two substances touch, such as between two liquids (herbicide and water) or between a solid and a liquid (herbicide and leaf surface). Some surfactants act as dispersing agents, helping to keep a pesticide suspended in water. Others work on the plant, improving the wetting, sticking and penetrating characteristics of the herbicide droplets. Oil-based adjuvants contain petroleum or vegetable oil plus an emulsifier that suspends the oil in tiny droplets within the spray solution. Oil-based adjuvants typically assist in herbicide penetration into the leaf.

There are two basic type of surfactants (ionic and non-ionic), of which the non-ionic are most common. The following table lists the surfactants registered for use with herbicides in Western Canada.

Crop and Herbicide Recommendation Tables

The ratings in the following charts give general comparisons based on rates, timing and other application instructions and precautions as outlined in this Guide.

The ratings for crop tolerance and weed control are explained below. Where ratings are not given, a dot (•) will indicate if the weed is listed on a product label. Where rate ranges are listed for controlling a given weed, ratings are based on results achieved with the higher rate.

RATING	CROP TOLERANCE	WEED CONTROL
E - Excellent	Consistently exhibits a high degree of tolerance over a wide range of growing conditions.	Consistently provides a high degree of control over a wide range of growing conditions.
G - Good	Exhibits good crop tolerance under most growing conditions.	Provides good control under most growing conditions.
F - Fair	Tolerance is acceptable depending on growing conditions.	Control is acceptable depending on growing conditions.
P - Poor		Provides suppression, but control is often unacceptable.

All ratings are primarily based on the use of the herbicides on their own. Tank mixing herbicides may reduce the control of one or both tank mix partners, or result in a reduced level of crop safety.

Weed Control Tables

Table 1. Weed Control in Barley

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D	47	G								P		G	P	E		G	E		E		G	E	F	E	P	P		E	P
Achieve Liquid	54	G	F	G			G																						
Achieve Liquid Gold	56	G	F	G			G	E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Avadex BW/MicroActiv	70	E					G																						
Ally Toss-N-Go + 2,4-D	59	G						F		G				E	G	G	E		E	E	G	E	G	E	F	F		G	
Assert	63	E					G	P											G					G					
Attain	69	G						G		F	E	G	G[5]	E	F	E	E	E	E	E	G	E	G	E	P	P	E	E	•
Avenge	74	G					G																						
Bromoxynil	79	E						G				F				F	G		F	F	G		G	F					
Bromoxynil + 2,4-D*	62, 194	E						E	F			G		E		G	E		E	G	G	G	G	G				G	G
Bromoxynil + MCPA	81	E						E	F			G		G[3]		G	E		E	G	G	G	G	G	P	P		G	G
Curtail M	86	G						G					G[5]	E		P	E		E	F		E	F	E	G	G		F	G
Dicamba +MCPA/2,4-D	88	F						G			G	G	P	E	F[2]	G	E		E	G	G	E	G	E	F	F		G	F
Dicamba + mecoprop + MCPA	90	F						G	G		G			E	F	G	E		E	G	G	E	G	E	F	F		G	G
Dichlorprop + 2,4-D	92	G						G	G			G		E		G	E	F	E	G	G	E	G	E	F	F		G	G
DyVel	95	F						G			F	G		E	F	G	E		E	G	G	E	G	E	F	F		F	F
DyVel DSp	96	F						G			F	G		E		G	E	F	E	G	G	E	G	E	F	F		G	F
Express Pack	109	G						F						E		G	E		E	G	G	E		E		F			G
Frontline	115	E						E		E	E		F[10]		G		E		E	E		E	E	E	F	F		E	
Glyphosate (preharvest)	124				G							G[6]													F	G			
Hoe-Grass 284	132	G	G	G		F	G																						
Hoe-Grass II	134	G	G	G		F	G	G	P							F	G		F	F	G		G	F					
Linuron + MCPA amine	146	F		P				G		G		G		E	F	G	E		E	G		E	F	E					
Lontrel	149	E						G																	G	G			
MCPA	151	E										G	P	E	F	F	E		E	F	F	E	P	E	P	P			P
MCPAK	151	E										G		E	F		E		E			E		E	P	P			
Mecoprop	155	G								E	G						E		E							P			
Prestige	166	E						E		G	E		G[5]	E	G	E	E	E	E	G		E	G	E	E	E	E	E	E
Prevail Liquid	168	G		G			G	F					G[5]	E		P	E		G	F		E	F	E	G	G		F	
Puma® Super	171	G	G	E			G																						
Refine Extra Toss-N-Go	176	E						G		E	F			E	E	G	E	F	E	E	E	G	E	E	F	F		G[4]	G
Sencor	184	F							G	G					F		G		G	F	G		G	G				G	
Spectrum	190	E						E		E	E		F[10]	E[5]	E		E		E	F		E	E	E	G	G		E	
Trifluralin (foxtail control)	199	E		G																									
Trifluralin (grassy and broadleaf)	201	F	G	G			F	F		G							G			G	G								
Trophy	206	E						F			G	G		E	F	E	E		E	E	G	G	F	G	P	P		E	
Unity[1]	209	E						G		G	G			G	E	G	G		E	G		G	G	E				G	

[1] Not for use in Manitoba. [2] 2,4-D mixes only. [3] MCPA K mixes only. [4] Will not control CLEARFIELD canola varieties. [5] Spring seedlings only. [6] Not registered for control with all glyphosate products. See glyphosate sections for details. [10] Seedlings and overwintered rosettes. • Ratings not yet developed. *Includes Thumper and Approve herbicides.

Table 2. Weed Control in Oats

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
Bromoxynil	79	E					G				F				F	G		F	F	G		G	F					
Bromoxynil + MCPA	81	E					E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Curtail M	86	G					G					G[8]	E		P	E		E	F		E	F	E	G	G		F	G
Dicamba + MCPA	88	F					G			G	G	P	E	F[2]	G	E		E	G	G	E	G	E	F	F		G	F
Dicamba + mecoprop + MCPA	90	F					G	G		G			E	F	G	E		E	G	G	E	G	E	F	F		G	G
DyVel	95	F					G			F	G		E	F	G	E		E	G	G	E	G	E	F	F		F	F
Frontline	115	E					E		E	E		P[8]		G		E		E	E		E	E	E	F	F		E	
Glyphosate (pre-harvest)[5]	124				G							G[6]												F	G			
Linuron + MCPA amine	146	F		P			G		G		G		E	F	G	E		E	G		E	F	E					
Lontrel	149	G					F																	F	G			
MCPA	151	E									G	P	E			E		E			E		E	P	P			P
MCPAK	151	E									G		E	F		E		E			E		E	P	P			
Mecoprop	155	G							E	G						E		E							P			
Refine Extra Toss-N-Go	176	G					G		E	F			F	E	G	E	F	E	E	E	G	E	E	F	F		G[3]	G
Spectrum	190	E					E		E	E		P[8]	E[4]	E		E		E	E		E	E	E	G	G		E	

[2] MCPAK mixes only. [3] Will not control CLEARFIELD canola varieties. [4] Spring seedlings only. [5] Not all products registered for use in oat. See glyphosate sections for details. [6] Not registered for control with all products. See glyphosate sections for details. [8] Seedlings and overwintered rosettes. • Registered for control, ratings not yet developed.

Table 3. Weed Control in Rye or Triticale

HERBICIDE	PAGE	Crop Tolerance - Rye	Crop Tolerance - Triticale	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D[3,4]	47	G											G	P	E			E		E		G	E		E	P	P			
Achieve Liquid[3]	54	G	G	F	G			G																						
Achieve Liquid Gold[3,4]	56	G	G	F	G			G	E	F			G		G		G	E		E	G	G	G	G	G	P	P		F	G
Avenge[1]	74	G	G					G																						
Bromoxynil[1]	79	E	E						G				F				F	G		F	F	G		G	F					
Bromoxynil + MCPA[1,4]	81	E							E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Dicamba + 2,4-D[2,4]	88	F							G			G	G	P	E		G	E		E	G	G	E	G	E	F	F		G	G
Hoe-Grass 284[3]	132	G	G	G	G		F	G																						
Hoe-Grass II[2]	134	G	G	G	G		F	G	G	P							F	G		F	F	G		G	F					
MCPA[3,4]	151	E											G	P	E			E		E			E		E	P	P			P

[1] Fall rye only. [2] Spring rye only. [3] Fall and Spring rye. [4] Rye only; not registered for triticale.

Table 4. Weed Control in Winter Wheat

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-Purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D	47	G										G	P	E			E		E		G	E		E	P	P			
Achieve Liquid	54	G	F	G			G																						
Achieve Liquid Gold	56	G	F	G			G	E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Avenge[3]	74	G					G																						
Bromoxynil	79	E						G				F				F	G		F	F	G		G	F					
Bromoxynil + MCPA	81	E						E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Dicamba + MCPA/2,4-D	88	F						G			G	G	P	E	F	G	E		E	G	G	E	G	E	F	F		G	F
Dicamba + mecoprop + MCPA	90	G						G	G		G			E	F	G	E		E	G	G	E	G	E	F	F		G	G
Dichlorprop + 2,4-D	92	G						G	G			G		E		G	E	F	E	G	G	E	G	E	F	F	P	G	G
DyVel	95	F						G			F	G		E	F	G	E		E	G	G	E	G	E	F	F		F	F
DyVel DSp	96	F						G			F	G		E		G	E	F	E	G	G	E	G	E	F	F		G	F
Hoe-Grass 284	132	G	G	G		F	G																						
MCPA	151	E										G	P	E			E		E			E		E	P	P			P
Refine Extra Toss-N-Go	176	G						G		E	F				E	G	E	F	E	E	E	G	E	E	F	F		G[2]	G
Sencor	184	F												F								F		G					

[1] MCPAK mixes only. [2] Will not control CLEARFIELD canola varieties. [3] Norstar only.

Table 5. Weed Control in Spring Wheat

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D	47	G								P		G	P	E		G	E		E	G	G	E	F	E	P	P		E	
Achieve Liquid	54	G	F	G			G																						
Achieve Liquid Gold	56	G	F	G			G	E	F			G		G		G	E		E	G	G	G	G	G	P	P		G	G
Adrenalin[11]	57			•			•	S		•	S	•	•	•		•	•		•	•	•	•	•	•				•	
Ally Toss-N-Go + 2,4-D	59	G						F		G				E	G	G	E		E	E	G	E	G	E	F	F		G	
Assert	63	G					G	P											G					G					
Attain	69	G						G		F	E	G	G[2]	E	F	E	E	E	E	E	G	E	G	E	P	P	E	E	
Avadex BW/MicroActiv	70	G					G																						
Avenge[7,8]	74	G					G																						
Basagran + 2,4-D	76	E										G					G		E	F	G	E		E				E	
Bromoxynil	79	E						G				F				F	G		F	F	G		G	F					
Bromoxynil + 2,4-D*	62, 194	E						E	F			G		E		G	E		E	G	G	G	G	G				G	G
Bromoxynil + MCPA	81	E						E	F			G		G		G	E		F	G	G	G	G	G	P	P		G	G
Curtail M	86	G						G					G[2]	E		P	E		E	F		E	F	E	G	G		F	G
Dicamba + MCPA/2,4-D	88	G						G			G	G	P	E	F[6]	G	E		E	G	G	E	G	E	F	F		G	F
Dicamba + mecoprop + MCPA	90	G						G	G		G			E	F	G	E		E	G	G	E	G	E	F	F		G	G
Dichlorprop + 2,4-D	92	G						G	G			G		E		G	E	F	F	G	G	E	G	E	F	F	P	G	G
DyVel	95	G						G			F	G		E	F	G	E		E	G	G	E	G	E	F	F		F	F
DyVel DSp	96	G						G			F	G		E		G	E	F	E	G	G	E	G	E	F	F		G	F
Everest	107	F		E			G																						
Express Pack	109	G						F						E		G	F		E	G	G	E		E		F			G
Frontline	115	E						E		E	E		F		G		E		E	E		E	E	E	F	F		E	
Frontline + 2,4-D	114							•		•	•	•	•[2]	•	S	•	•		•	•	•	•	•	•	S	S			
Glyphosate (preharvest)	124				G								G[4]												F	G			
Harmony Total	131	G	G	G			E	G		E	F			E	E	E	E	F	E	E	E	G	E	E	F	F		G[5]	
Hoe-Grass 284	132	G	G	G		F	G																						
Hoe-Grass II	134	G	G	G		F	G	G	P							F	G		F	F	G		G	F					
Horizon	135	G	E	G			E																						•
Horizon BTM	137	G	E	G			E	E	F			G		G		G	E		F	G	G	G	G	G	P	P		G	G
K2	138	F		E			G	G		E		F		E	E	E	E	F	E	E	E	E	G	E	F	F		G[5]	G
Linuron + MCPA amine	146	F		P				G		G		G		E	F	G	E		E	G		E	F	E					
Lontrel[7]	149	E						G																	F	G			
MCPA	151	E										G	P	E	F[6]		E		E			E		E	P	P			P
Mecoprop	155	G								E	G				F	F	E		E	F	F			P		P			
Prestige	166	E						E		G	E		G[2]	E	G	E	E	E	E	G		E	G	E	E	E	E	E	G
Prevail Liquid	168	G		G			G	F					G[2]	E		P	E		E	F		E	F	E	F	G		F	
Puma[10] Super	171	G	G	E			G																						
Refine Extra Toss-N-Go	176	E						G		E	F			E	E	E	E	F	E	E	E	G	E	E	F	F		G[5]	G
Sencor	184	F							G	G					F		G		G	F	G		G	G				G	
Spectrum	190	E						E		E	E		F	E[6]	E		E		E	E		E	E	E	G	G		E	
Sundance	192	G		F	F		G			E	E								E	E				E				G[5]	
Trifluralin (foxtail control)	199	E		G																									
Triumph Plus[7,8]	204	G		E			G	G		G		G		E	E	F	E		E	E	G	E	E	E		P		G[5]	E
Trophy	206	E									G	G		E	F	E	E		E	E	G	G	F	G	P	P	E	E	
Unity[9]	209	E						G		G	G			G	E	G	G		E	G		G	G	E				G	

[1] Green foxtail only. [2] Spring seedlings and overwintered rosettes. [3] MCPA K mixes only. [4] Not registered for control with all products. See glyphosate sections for details. [5] Will not control CLEARFIELD canola varieties. [6] Spring seedlings only. [7] Not registered for durum wheat. [8] Not recommended for all spring wheat varieties. Check product listing for details. [9] Not for use in Manitoba. [10] Tank mix with 2,4-D ester when applying to durum wheat. [11] For use on CLEARFIELD wheat varieties only.

• Registered for control, ratings not yet developed. * Includes Thumper & Approve herbicides.

Table 6. Weed Control in Corn

HERBICIDE	PAGE	Barnyard Grass	Volunteer Cereals	Foxtail, Green	Foxtail, Yellow	Wild Oats	Quackgrass	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed Annual Species, Lady's-thumb	Sow Thistle (Perennial)	Stinkweed	Sunflower, Volunteer	Thistle, Canada	Volunteer Canola
2,4-D	47									P		G	P	E	G	E	E	G	G	E	F	P	E	P	P	E
Accent	52	•		•	S	•	•																			
Atrazine	67					•		•								•	•	•			•					
Basagran/Basagran Forté	76									•		•				•	•	S	•	•	•		•		S	
Bromoxynil	79							G				F			F	G	F	F	G		G		F			
Bromoxynil + MCPA	81							•	•			•		•	•	•	•	•	•	•	•		•	•		•
Dicamba	88							•			•				•	•	•	•			•	S			S	
Dicamba + 2,4-D amine	88							•			•	•	S	•	•	•	•	•	•	•	•	S	•	•	S	•
Dual II Magnum	94	•		•	•													S								
DyVel DSp	96							G			F	G		E	G	E	E	G	G	E	G	F	E		F	G
Eradicane	104	•	•	•	•	•	S			•						•		•								
Frontier	112			•																						
Glyphosate[1,4]	125	E	E	E	E	E	F	G	G	E	E		F	E	F	E	E	E	E	E	E	F	E	E	F	
Laddok	141									•		•				•	•	•	•		•					
Liberty 200 SN[3]	145	•		•	•	•	S	•		•		•				•	•	•		•	•	•	•		S	
MCPB + MCPA	154															•	•	•		•		S	•		S	•
Primextra II Magnum	169	•		•	•			•								•	•	•			•					
Shotgun	188							•								•	•	•			•					
Ultim[2]	207	•	•	•	•	•	•											•								

• Controlled S – Suppression

[1] For use on Roundup Ready varieties only. [2] See product page for registered corn varieties. [3] For use on Liberty 200 SN tolerant corn varieties.
[4] Not all glyphosate products are registered for use on Roundup Ready Corn. [5] Registered for use only in Manitoba.

Table 7. Weed Control in Peas

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp Nettle	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada
Assure II	65	E	G	E	F	E	E	G																		
Avadex BW	70	E						G																		
Basagran/Basagran Forté	76	F									F	F	F					F	F	P	F	F	F	F		P
Clethodim	84	E	E	E	F	F	E	G																		
Edge	100	G	E	E		P	F	G	G		G	P				P	G	E		G	P		P			
Fusion	117	E	G	E[1]		E	E	G																		
Glyphosate (preharvest)	124				G									G[2]											F	G
Hoe-Grass 284	132	G	G	G				G																		
MCPA Sodium Salt/Amine	151	F													P			P	G			P		P		
MCPB + MCPA	154	F														F		G	G	G		G		G	P	P
Odyssey	159	G	G	E[1]		G	G	G	F		E	E			E	F	P	P	E	E	E	E	G	E		
Pea Pack	160	F																G	E	G				G		
Poast Ultra	162	E	E	E	F	E	E	G																		
Pursuit	173	G		F				F	G		G	G				G		P	E	G		G	G	E		
Pursuit Ultra	175	G	E	E	F	E	E	G	F		G	G				F			E	G			F	G		
Sencor	184	F									G					F		G	G				G	G		
Trifluralin (Broadleaf & grassy weeds)	201	G	G	G				F	F		G							G		G	G					
Venture L	212	E	E	F	F	E	E	G																		

[1] Green foxtail only. • Registered for control, ratings not yet developed.
[2] Not registered for control with all products. See glyphosate sections for details.

Table 8. Weed Control in Other Pulses

HERBICIDE	PAGE	CROP						ANNUAL WEEDS																				PERENNIALS			
		Bean, Dry	Fababean	Lentil	Chickpea	Soybean	Sweet White Lupin	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Volunteer Barley	Volunteer Wheat	Wild Oat	Buckwheat, wild	Chickweed	Cleavers	Cocklebur	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Nightshade, Hairy	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Canada Thistle	Dandelion	Perennial Sow-thistle	Quackgrass
Assure II	65			X		X		G	E		E	E	G																		F
Basagran/Basagran Forté[1]	76	X	X			X									F	F	F			F	F	F	P	F	F	F	F	P			
Clethodim	84	X		X	X	X		E	E	E	E	E	G																		
Dual II Magnum	94	X				X	X	G	G	G													P								
Edge	100	X	X	X		X		E	E	E	P	F	G	G	G	P		P	G	E			G	P		P					
Eptam	102	X						G	G	G	F	F	G		G					G			G								
Frontier[1]	112	X							•																						
Fusion	117			X				G	E		E	E	G																		
Glyphosate (preharvest)	124	X		X		X																						G	G[4]	F	G
Hoe-Grass 284	132	X	X	X		X		G	G	G			G																		
Liberty (desiccant only)	142			X																											
Linuron	146					X[2]	X[2]	S	•	•				•	•								•		•	•	•				
Pinnacle	161					X														•	•		•			•					
Poast Ultra	162	X	X	X	X	X		E	E	E	F	F	G																		F
Pursuit[3]	173	X[3]				X																G									
Reflex + Basagran	178	X[5]													F					•	•		•		•	•	•				
Sencor (post-emergence)	184			X	X										F			F		F	F					F	F				
Sencor + Treflan (PPI)	184		X			X		G	G	G			F	F	G			F		G	G		G	G	G	G	G				
Sencor + Edge (PPI)	184		X					G	E	E	F	F	G	G	G	F		F	G	E	F		G	F		G	F				
Trifluralin (Broadleaf and grassy weeds)	201	X	X	X		X		G	G	G			F	F	G					G			G	G							
Venture L	212			X		X[4]		E	F	F	E	E	G																		F

[1] White and Kidney beans only. [2] Pre-emergent surface treatments only. [3] Pinto, pink and red beans only. (Refer to Pursuit section for full list of weeds controlled).
[4] Not registered for control with all products. See product section for details. [5] For use on navy beans in the Red River Valley of Manitoba. Does not include Basagran Forté.
• Registered for control, ratings not yet developed.
S - Registered for suppression.

Table 9. Weed Control in Flax and Solin (Low Linolenic Acid Varieties). Not all products are registered for use on Solin.

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Quackgrass	Volunteer Cereals	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp Nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada
Assure II[1]	65	E	G	E		F	E	G																			
Avadex BW	70	E						G																			
Basagran[1]/Basagran Forté	76	F									F	F	F					F		F	P	F	F	F	F		F
Bromoxynil + MCPA[1]	81	F							E	F			G		G		G	E		E		G	G	F	G	P	P
Clethodim[1]	84	E	E	E	E	F	E	G																			
Curtail M	86								F					G[2]	E			E		E	P		E	P	E	G	G
Elite	101		•	•	•	•	•	•	•	•			•		•		•	•		•		•	•	•	•	•	•
Eptam[3]	102	F	G	G	G		F	G			G							G			G						
FlaxMax	111	F	E	E	E		G	G	F					G[2]	E			E		E	P		E	P	E	G	G
Fusion[1]	117	E	G	E			E	G																			
Glyphosate (preharvest)[4]	124					G								G[5]												F	G
Hoe-Grass 284	132	G	G	G	G			G																			
Hoe-Grass II	134	F	G	G	G			G	G	P							F	G		F	P	G		G	F		
Lontrel	149	E							G																	F	G
MCPA/MCPAK	151	F											G	P	E	F		E		E			E		E	P	P
Pardner[1] (Bromoxynil)	79	F							G				F				F	G		F	F	G		G	F		
Poast Ultra[1]	162	E	E	F	F	F	E	G																			
Trifluralin (Broadleaf and grassy weeds)	201	F	G	G	G			F	F		G							G			G	G					
Venture L	212	E	E	F	F	F	E	G																			

[1] Registered for use on Solin (low linolenic acid varieties). [2] Spring seedlings only. [3] Not recommended for use on flax in Saskatchewan. [4] Not all products registered for use in flax. See glyphosate sections for details. [5] Not registered for control with all products. See glyphosate sections for details.

Table 10. Weed Control in Canola

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada
Absolute[2]	51	G	G	E			G	G	G	G		E	G			E	E	P	P		E	E	E	E	G	E	G	G
Assure II	65	E	G	E		F	E	E	G																			
Avadex BW	70	E							G																			
Clethodim	84	E	F	E	E	F	E	E	G																			
Eclipse[2]	99	G		E		F	E	E	E	G	G	E	E		F		E	E	E		E	E	E	E		E	G	G
Edge	100	G	E	E	E		P	F	G	G		G	P				P	G	E			E	P		P			
Fusion	117	E	G	E			E	E	G																			
Glyphosate[4,5]	125	E	E	E	E	F	E	E	E	F	G	E	E		F	E	E	F	E	G	E	E	E	E	E	E	F	F
Glyphosate (preharvest)	124					G									G												F	G
Hoe-Grass 284	132	G	G	G	G				G																			
Liberty[1]	142	G	G	G		F	F	G	G	G		G	G		F	G	G	G	G	G	G	G	G	G	G	G	G	F
Lontrel	149	G								G																	F	G
Muster Gold II	156	G	G	E		F	E	E	G							G	G				G	P			G	F		
Muster Toss-N-Go	157	G														G	G				G	P			G	F		
Odyssey[2]	*159	G	G	E			G	G	G	F		E	E			E	G	P	P		E	E	E	E	G	F		
Poast Ultra	162	E	E	E	E	F	E	E	G																			
Pursuit[2]	173	E		G			P	P	F	G		G	G						P		E	G			G	E		
Pursuit Ultra[2]	175	E	E	E	E	F	E	E	G	F		G	G				F				E	G			F	G		
Trifluralin	201	E	G	G	G				F	F		G							G			G	G					
Venture L	212	E	E	F	F	F	E	E	G																			

[1] For use **only** on Liberty Link varieties. [2] For use **only** on CLEARFIELD canola varieties. [3] Ratings based on 1.35 L/acre rate of Liberty. Control may be reduced at lower rates. [4] For use **only** on Roundup Ready canola varieties. [5] Not registered for control with all products. See glyphosate sections for details. [6] Season long control.
• Registered for control, ratings not yet developed. S – Suppression.

Table 11. Weed Control in Potatoes

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green and Yellow	Volunteer Tolerant Canola	Volunteer Corn	Volunteer Barley	Volunteer Wheat	Wild Oats	Quackgrass	Chickweed	Dandelion	Hemp Nettle	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Pigweed, Prostrate	Purslane	Smartweed (Annual)	Shepherd's-purse	Stinkweed	Volunteer Canola
Clethodim	84	•	•		•	•	•	•													
Dual II Magnum	94	•	•												S						
Eptam 8-E	102	•	•			•	•	•	S	•			•		•	•	•				
Glyphosate (preplant use only)	120		•	•	•	•	•	•	•		•	•	•	•	•			•	•	•	
Hoe-Grass 284	132	•	•		•			•													
Liberty (dessicant only)	142																				
Linuron (pre-emergent use only)	146	S	•							•			•		•		•	•	•	•	
Poast Ultra	162	•	•		•	•	•	•	S												
Prism	170	•	•						•				S		•						
Reglone Desiccant	179																				
Sencor	184									•		•	•	•	•			•	•	•	•
Venture L	212	•	•		•	•	•	•	S												

• Controlled S – Suppression [1] Will control non-Roundup Ready varieties only.
Note: Before using any pesticides on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

Table 12. Weed Control in Sunflowers

HERBICIDE	PAGE	Crop Tolerance	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada
Assert	63	F																		G				G		
Clethodim	84	E	E	E	F	E	E	G																		
Edge	100	G	E	E		P	F	G	G		G	P				P	G	E			E	P	P			
Eptam 8E	102	G	G	G		F	F	G			G							G			G					
Hoe-Grass 284	132	G	G	G				G																		
Poast Ultra	162	E	E	F				G																		
Trifluralin	201	E	G	G				F	F		G							G			G	G				
Venture L	212	E	E	F	F	E	E	G																		

Table 13. Weed Control in Special Crops

HERBICIDE		CROP						ANNUAL WEEDS																								PERENNIALS			
	PAGE	Canaryseed	Safflower	Caraway	Coriander	Buckwheat	Mustard	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Volunteer Wheat	Wild Oat	Buckwheat, wild	Catchfly, night-flowering	Chickweed	Cleavers	Cocklebur	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Volunteer Barley	Volunteer Flax	Volunteer Mustard, Canola	Canada Thistle	Dandelion	Perennial Sow-thistle	Quackgrass
Avadex BW	70	✓[1]					✓					G																						
Avenge	74	✓										G																						
Bromoxynil	79	✓											G				F			F	G	F	F	G		G	F							
Bromoxynil + MCPA	81	✓											E	F			G	G		G	E	E	G	G	G	G	G			G	P		P	
Clethodim	84						✓	E	E	E	E	G																E						F
Curtail M	86	✓											F					E		P	E	E	P		E	P	E			F	G	G[4]	G	
Dicamba + MCPA	88	✓											G			G	G	E	F		E	E	G	G	E	G	E		P	G	F	P	F	
Dicamba + Mecoprop + MCPA	90	✓											G	G		G		E	F	G	E	E	G	G	E	G	E				F		F	
Edge	100		✓	✓	✓		✓[3]	E	E	E	F	G	G		G	P			P	G	E		G	P		P		P						
Fusion	117						✓	G	E			G																						
Hoe-Grass 284	132					✓	✓	G	G	G		G																						
Muster Toss-n-Go	157						✓[2]											G	G			G				G	F							
Poast Ultra	162		✓	✓	✓	✓	✓	E	E	E	E	G																E						F
Prestige	166	✓											E		G	E		E	G	E	E	E	G		E	G	E		E	E	E	G[4]	E	
Trifluralin	201		✓				✓	G	G	G		F	F		G						G		G	G										
Venture L	212						✓	E	F	F	E	G																E						F

[1] Granular formulation only. [2] Yellow mustard only. [3] Brown and oriental only. [4] Spring seedlings only.

Table 14a. Herbicides for Use Before Seeding or After Seeding but Prior to Crop Emergence

HERBICIDE	RATE L/acre	pre seeding	pre emergent	barley	canaryseed	canola	chickpea	corn, field	corn, sweet	dry beans	field pea	flax	forage grasses	lentil	oats	rye	soybean	wheat
Amitrol	1.7	✓		✓		✓		✓		✓	✓						✓	✓
Glyphosate	glyphosate[1]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + Express TNG	glyphosate[1] + 4g	✓		✓											✓			✓
Glyphosate + Pardner	glyphosate[1] + 0.51	✓	✓	✓											✓			✓
Prepass	40 acres/case	✓		✓											✓			✓
Glyphosate+ 2,4-D	glyphosate[1] + 0.23-0.58	✓	✓	✓												✓		✓
Glyphosate + Buctril M	glyphosate[1] + 0.2-0.4	✓		✓	✓			✓	✓			✓	✓		✓	✓		✓
Glyphosate + MCPA	glyphosate[1] + 0.2-0.28	✓		✓			✓	✓	✓		✓	✓		✓	✓	✓		✓
Glyphosate + MCPA	glyphosate[1] + 0.2-0.4	✓		✓				✓	✓			✓			✓	✓		✓
Rustler	1.0-1.3	✓		✓				✓							✓	✓		✓

[1] Rate of application of glyphosate vary among brands. Consult Table 10 on page 127 for specific application rates of glyphosate.

Table 14b. Weed Control Before Seeding or After Seeding but Prior to Crop Emergence

HERBICIDE	RATE L/acre	PAGE	Brome, Downy	Buckwheat, Wild	Dandelion	Flixweed	Foxtail Barley	Foxtail, Green	Hemp-nettle	Kochia	Lady's-thumb	Lamb's-quarters	Mustard, Wild	Narrow-leaved Hawk's Beard	Pigweed, Redroot	Quackgrass	Russian Thistle	Shepherd's Purse	Stinkweed	Volunteer Cereals	Volunteer Canola (including Roundup Ready varieties)	Volunteer Flax	Wild Oats
Amitrol 240	1.7 L	61			•																		
Glyphosate[1] + 2,4-D	glyphosate[1] + 0.23-0.58	119	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate[1] + Buctril M	glyphosate[1] + 0.2-0.4	119	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Express TNG	glyphosate[1] + 4g	119		•	•	•	S	•	•	•		•	•	S	•		•	•	•	•			•
Glyphosate[1] + MCPA	glyphosate[1] + 0.2-0.4	119	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Pardner	glyphosate[1] + 0.51	119		•				•		•	•	•	•				•		•	•	•		•
Glyphosate	glyphosate[1]	119	•	•	S	•	S	•	•	•	•	•	•	•	•	•	•	•	•	•		•	•
Prepass	40 acres/case	165	•	•	S	•		•	•	S	•	•	•	•	•		•	•	•	•	•		•
Rustler	1.0 to 1.3	183	•	•		•		•		•	•	•	•		•		•		•	•			•

• Controlled S – Suppression [1] Rate of application of glyphosate vary among brands. Consult Table 10 on page 127 for specific application rates of glyphosate.

Table 15. Weed Control in Summerfallow (Chem Fallow)

HERBICIDE	RATE L/acre	PAGE	Brome, Downy	Buckwheat, Wild	Dandelion	Flixweed	Foxtail Barley	Foxtail, Green	Kochia	Lady's-thumb	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Quackgrass	Russian Thistle	Stinkweed	Thistle, Canada	Thistle, Sow (Perennial)	Volunteer Cereals	Volunteer Canola	Wild Oats
Dicamba + 2,4-D	0.12 + 0.45	88		•		•			•	•	•	•	•		•	•	S	S			
Dicamba + mecoprop + MCPA	0.81	90		•		•			•	•	•	•	•		•	•	S	S		•	
DyVel DSp	0.71	96		•		•			•	•	•	•	•		•	•	S			•	
Express Pack	40 acres/pack	109				•										•					
Glyphosate	0.5 to 1.0	119	•	•		•	S	•	•	•	•	•	•		•	•			•	•[2]	•
Glyphosate + 2,4-D amine (500 g/L)	glyphosate[2] + 0.49	119				•		•	•	•	S	•	•		S	•			•	•	•
Glyphosate + dicamba	glyphosate[3] + 0.12	119		S		•		•	•	•	•	•	S		•	•			•	•[2]	•
Glyphosate + Bromoxynil	glyphosate[3] + 0.51	119		•				•	S	•		•	S			•			•	•	S
Lontrel	0.34	149		•													•	S			
Rustler	1.0 to 1.3	183	•	•		•	S	•	•	•	•	•	•		•	•			•	•[2]	•

• Controlled S – Suppression [1] Not for use in Manitoba. [2] Not including Roundup Ready canola. [3] Rate of application of glyphosate vary among brands. Consult Table 10 on page 127 for specific application rates of glyphosate.

Table 16. Fall Weed Control in Stubble

HERBICIDE	PAGE	Flixweed	Narrow-leaved Hawk's Beard	Shepherd's-purse	Stinkweed	Canada Thistle	Quackgrass	Dandelion
2,4-D	47	•		•	•	S		S
Amitrol 240	61					S		
Dicamba	88					S		S
Dicamba + Glyphosate	88	•		•	•	S	S	
Dicamba + mecoprop + MCPA	90	•		•	•	S		
DyVel DSp	96	•		•	•	S		
Glyphosate	119					S	•	•
MCPA	151	•		•	•	S		S

• Controls fall seedlings less than 2 inches (5 cm) across.
S – Suppression. Levels of suppression vary depending on the product and growing conditions in the fall. Regrowth and in-crop treatments can be expected.

Table 17. Weed Control in Grass Pastures and Hayfields

HERBICIDE	RATE L/acre or kg/acre	PAGE	Absinth	Bindweed, Field	Burdock	Canada Thistle	Dandelion	Dock, Curled	Daisy, English	Flixweed	Foxtail Barley	Gumweed	Hawk's Beard, Narrow-leaved	Knapweed	Leafy Spurge	Nodding Thistle	Poplar	Pussy Toes	Red Bartsia	Sage, Pasture	Sow-thistle, Perennial	Snowberry	Stinkweed	Tansy	Wild Rose	Willow	Wormwood, Biennial
2,4-D (500 g/L)	0.57 - 1.82	47	□	□	•	□	□	•		•		□			□	□	□	□	•	□	□	□	•				
2,4-DB	0.71 - 1.72	49		□		□		■					■								□		•				
Dicamba	0.85 - 1.86	88		□		□	□	□	□					□	□					□	□			□			
Dicamba + 2,4-D	0.85 + 0.89 - 1.62	88	□	□	•	□	□	□	□	•		□			□	□	□		•	□	□	•	•	□	•	□	
Escort	0.010 - 0.012	105				□	•														□	•		•	■		
Grazon	2.8	130			•	•	•	•																			
Kerb	0.36 - 0.45	139									•																
MCPA (500 g/L)	0.7 - 1.9	151		□		□	□	□		•		□			■						□		•	□			•
MCPB + MCPA	1.11 - 1.72	154		□																	□		•				
Remedy EC	1 - 8	181		■	•		•	•									•								•	•	
Tordon 22K	0.45 - 3.64	195		■		•								•	■					•							

• Controlled. ■ Controlled by the highest rate within this range. □ Top growth suppression.

Table 18. Weed Control in Shelterbelts

HERBICIDE	USE				SHELTERBELT SPECIES												WEEDS																				
	PAGE	Before Planting	After Planting (New & Established)	Established	American Elm	Birch	Caragana	Crabapple	Green Ash	Juniper	Lilac	Manitoba Maple	Poplar	Scots Pine	Siberian Elm	Willow	Barnyard Grass	Foxtail, Green and Yellow	Wild Oats	Buckwheat, Wild	Chickweed	Cleavers	Cocklebur	Dandelion	Flixweed	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Red Root	Quackgrass	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Stinkweed	Thistle, Sow (Perennial)	Thistle, Canada
Amitrol 240	61			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Casoron	83		•		•		•	•	•	•	•	•			•	•		•		•	•						•	•	•	•	•		•	•			
Glyphosate	119	•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Gramoxone	129		•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		•	•	•	•	•	•		•	•	•	•		
Linuron	146			•	•		•		•			•		•	•	•	•	•		•	•						•	•		•				•	•		
Simadex Simazine	189			•	•		•		•			•			•		•	•[1]	•	•								•						•			
Trifluralin Liquids	201	•		•	•		•		•					•	•		•	•	•	•	•							•		•		•					

[1] Yellow foxtail only.

Herbicide Formulations and Companies

Product	*Common Name*	*Formulation*	*Company*
2,4-D	2,4-D	500, 600, 700 g/L SN, EC	Various
Absolute	imazamox + imazethapyr + clopyralid	35% + 35% DG + 360 g/L	BASF Canada
Accent	nicosulfuron	75% DF	E. I. duPont Canada
Achieve Liquid	tralkoxydim	400 g/L SC	Syngenta
Achieve Liquid Gold	tralkoxydim+ bromoxynil + MCPA E	400 g/L SC + 280 g/L+280 g/L EC	Syngenta
Adrenalin	imazamox +2,4-D LV 700 ester	11.40% SN 660 g ae/L EC	BASF Canada
Adrenalin SC	imazamox + 2,4-D LV 700 ester	20 g/L + 560 g/L SC	BASF Canada
Advance	trifluralin	10% G	Dow AgroSciences
Ally Toss-N-Go	metsulfuron methyl	60% DF	E. I. duPont Canada
Amitrol 240	amitrole	240 g/L SN	Nufarm
Approve	bromoxynil + 2,4-D LV Ester	280 g/L + 660 g ae/L	Nufarm
Assert 300 SC	imazamethabenz	300 g/L SC	BASF Canada
Assure II	quizalofop-p-ethyl	96 g/L EC	E. I. duPont Canada
Atrazine	atrazine	480 g/L F; 90% DG	Syngenta, UAP; Syngenta
Attain	fluroxypyr+ 2,4-D E	180g/L + 564 g/L EC	Dow AgroSciences
Avadex BW/MicroActiv	triallate	400 g/L EC, 10% G	Gowan Company
Avenge200-C	difenzoquat	200 g/L SN	BASF Canada
Badge	bromoxynil+ MCPA E	225 g/L + 225 g/L EC	UAP
Banvel II	dicamba	480 g/L SN	BASF Canada
Basagran/Basagran Forté	bentazon	480 g/L SN	BASF Canada
Bonanza	trifluralin	400 g/L EC, 10% G	UAP
Buctril M	bromoxynil+ MCPA E	280 g/L+280 g/L EC	Bayer CropScience
Caliber	2,4-DB	400 g/L EC	UAP
Casoron	dichlobenil	4% G	UAP
Centurion	clethodim	240 g/L EC	Bayer CropScience
Clovitox Plus	MCPB+ MCPA	375 g/L+ 25 g/L SN	IPCO
Cobutox 600	2,4-DB	600 g/L EC	IPCO
Compitox	mecoprop	150 g/L SN	Nufarm
Credit	glyphosate	356 g/L SN	Nufarm
Curtail M	clopyralid+ MCPA E	50 g/L + 280 g/L EC	Dow AgroSciences
Dichlorprop-D	dichlorprop+ 2,4-D E	300 g/L + 282 g/L EC	IPCO
Dual II Magnum	metolachlor	915 g/L EC	Syngenta
DyVel	dicamba+ MCPA K	84 g/L + 336 g/L SN	BASF Canada
DyVel DSp	dicamba + 2,4-D A + mecoprop	110 g/L + 295 g/L + 80 g/L SN	BASF Canada
Eclipse	glyphosate + clopyralid	360 g/L SN + 360 g/L SN	Dow AgroSciences
Edge	ethalfluralin	5% G	Dow AgroSciences
Elite	sethoxydim + bromoxynil + MCPA E	450 g/L EC 200 g/L + 200 g/L	Nufarm
Embutox 625	2,4-DB	625 g/L EC	Nufarm
Eptam-8E	EPTC	800 g/L EC	Gowan Comapny
Eradicane-8E	EPTC + safener	800 g/L EC	Gowan Comapny
Estaprop	dichlorprop+ 2,4-D E	300 g/L + 282 g/L EC	Nufarm
Everest	flucarbazone-sodium	70% DG	Arvesta
Express Pack	tribenuron methyl+ 2,4-D E	75% DF + 700 g/L EC	E. I. duPont Canada
Factor	glyphosate	356 g/L SN	IPCO
FlaxMax	sethoxydim clopyralid + MCPA E	450 g/L EC + 50 g/L + 280 g/L EC	BASF Canada
Frontier	dimethanamid	900 g/L EC	BASF Canada
Frontline	florasulam+ MCPA E	50 g/L SC + 500 g/L EC	Dow AgroSciences
Frontline+2,4-D	florasulam + 2,4-D E	50 g/L + 564 g/L EC	Dow AgroSciences

Product	*Common Name*	*Formulation*	*Company*
Fusion	fenoxaprop-p-ethyl + fluazifop-p-butyl	80.5 g/L + 125 g/L EC	Bayer CropScience
Glyfos	glyphosate	360 g/L SN	Cheminova
Gramoxone	paraquat	200 g/L SN	Syngenta
Grazon	picloram+ 2,4-D E	65 g/L + 240 g/L EC	Dow AgroSciences
Harmony Total	clodinafop propargyl + thifensulfuron methyl + tribenuron methyl	240 g/L EC 50% + 25% DF	E. I. duPont Canada
Hoe-Grass 284	diclofop methyl	284 g/L EC	Bayer CropScience
Hoe-Grass II	diclofop methyl + bromoxynil	230 g/L + 80 g/L EC	Bayer CropScience
Horizon	clodinafop-propargyl	240 g/L EC	Syngenta
Horizon BTM	Clodinofop propargyl+ bromoxynil + MCPA E	240 g/L EC + 280 g/L + 280 g/L EC	Bayer CropScience
Kerb 50W	propyzamide	50% WP	Dow AgroSciences
Koril 235	bromoxynil	235g/L EC	Nufarm
Laddok	bentazon+ atrazine	200 g/L + 200 g/L SN	BASF Canada
Liberty	glufosinate ammonium	150 g/L SN	Bayer CropScience
Liberty 200SN	glufosinate ammonium	200 g/L SN	Bayer CropScience
Linuron 480/Linuron 400	linuron	480 g/L F; 400 g/L F	UAP
Lontrel	clopyralid	360 g/L SN	Dow AgroSciences
Lorox DF	linuron	50% DF	E. I. duPont Canada
Lorox L	linuron	480 g/L F	E. I. duPont Canada
Maverick	glyphosate	356 g/L SN	Dow AgroSciences
MCPA	MCPA	300,400,500,564 g/L SN, EC	Various
Mecoprop	mecoprop	150 g/L SN	UAP
Mextrol 400 M	bromoxynil+ MCPA E	200 g/L + 200 g/L EC	Nufarm
Muster Gold II	ethametsulfron methyl+ quizalofop-P-ethyl	75% DF + 96 g/L EC	E. I. duPont Canada
Muster Toss-N-Go	ethametsulfron methyl	75% DF	E. I. duPont Canada
Odyssey	imazamox + imazethapyr	35% + 35% DG	BASF Canada
Oracle	dicamba	480 g/L SN	Gharda/UAP/AgWest
Pardner	bromoxynil	280 g/L EC	Bayer CropScience
Pea Pack	metribuzin + MCPA Na	75% DG + 300 g/L SN	Bayer CropScience
Pinnacle	thifensulfuron methyl	75% DF	E. I. duPont Canada
Poast Ultra	sethoxydim	450 g/L EC	BASF Canada
PrePass	glyphosate + florasulam	360 g/L SN + 50 g/L SC	Dow AgroSciences
Prestige	fluroxypyr+ clopyralid + MCPA E	180 g/L + 50 g/L + 280 g/L EC	Dow AgroSciences
Prevail Liquid	tralkoxydim + clopyralid + MCPA E	400 g/L SC + 50 g/L + 280 g/L EC	Dow AgroSciences
Primextra II Magnum	metolachlor + atrazine	330 g/L + 170 g/L F	Syngenta
Princep Nine T	simazine	90% DG	Syngenta
Prism	rimsulfuron	25% DF	E. I. duPont Canada
Puma™ Super	fenoxaprop-p-ethyl	120 g/L EC	Bayer CropScience
Pursuit	imazethapyr	240 g/L SN	BASF Canada
Pursuit Ultra	sethoxydim + imazethapyr	450 g/L EC + 240 g/L SN	BASF Canada
Refine Extra Toss-N-Go	thifensulfuron methyl+ tribenuron methyl	50% + 25% DF	E. I. duPont Canada
Reglone Desiccant	diquat	240 g/L SN	Syngenta
Remedy	triclopyr	480 g/L EC	Dow AgroSciences
Renegade	glyphosate	356 g/L SN	Monsanto
Rival	trifluralin	500 g/L EC, 10% G	Bayer CropScience
Roundup Original/Transorb	glyphosate	356-360 g/L SN	Monsanto
Roundup WeatherMax/Ultra	glyphosate	540 g/L SN	Monsanto
Rustler	glyphosate + dicamba	132 g/L + 60 g/L SN	Monsanto
Select	clethodim	240 g/L EC	Arvesta
Sencor	metribuzin	75% DF, 75% DG, 500 g/L F	Bayer CropScience
Shotgun	atrazine + 2,4-D	280 g/L + 120 g/L F	UAP
Simazine 500	simazine	500 g/L F	UAP
Spectrum	florasulam + clopyralid + MCPA E	50 g/L SC + 50 g/L + 280 g/L EC	Dow AgroSciences
Sundance	sulfosulfuron	75% DG	Monsanto
Sword	dicamba + mecoprop + MCPA A	62.5 g/L + 62.5 g/L + 275 g/L SN	UAP
Target	dicamba + mecoprop + MCPAA	62.5 g/L + 62.5 g/L + 275 g/L SN	Syngenta
Thumper	bromoxynil + 2,4-D E	280 g/L + 280 g/L EC	Bayer CropScience
Tordon 22K	picloram	240 g/L SN	Dow AgroSciences
Touchdown IQ	glyphosate	360 g/L SN	Syngenta
Treflan	trifluralin	480 g/L EC, 5% G	Dow AgroSciences
Triumph Plus	fenoxaprop-p-ethyl + MCPA E + thifensulfuron methyl	56 g/L EC + 256 g/L EC + 75% DF	Bayer CropScience
Trophy	fluroxypyr + MCPA E	180 g/L + 500 g/L EC	Nufarm
Tropotox Plus	MCPB + MCPA A	400 g/L SN	Nufarm
Turboprop	dichlorprop + 2,4-D E	300 g/L + 282 g/L EC	UAP
Ultim	nicosulfuron + rimsulfuron	37.50% + 37.50% WDG	E. I. duPont Canada
Unity	bromoxynil + triasulfuron	280 g/L EC + 75% WG	Bayer CropScience
Vantage/Vantage Plus	glyphosate	356-360 g/L SN	Dow AgroSciences
Velpar DF	hexazinone	75% DF	E. I. duPont Canada
Venture L	fluazifop-p-butyl	125 g/L EC	Syngenta

Abbreviations

WP	wettable powder	EC	emulsifiable concentrate	WSP	water soluble powder	SN	solution	A	amine
DG	water dispersible granule	G	granule	DF	dry flowable	F	flowable	E	ester
WSG	water soluble granule	SC	suspension concentrate						

Products Available as Prepackaged Tank Mixes

Product Name (Manufacturer)	Component 1 or A	Component 2 or B	Component 3 or C	Crops	Weeds Controlled	Area Treated	Cost per acre (2004)
Absolute (BASF Canada)	Odyssey	Lontrel		CLEARFIELD canola	See component products	40 acres or 16 ha	$33.32
Achieve Liquid Gold (Syngenta)	Achieve Liquid	Buctril M	Turbocharge (adjuvant)	Wheat (spring, durum, winter), barley,	See the weeds of the component products plus tame buckwheat	20 acres or 8 ha	$23.27
Approve (NuFarm)	Koril	2,4-D		Spring wheat (incluing durum), barley	See component products	20 acres or 8 ha	$6.35
Eclipse (Dow AgroSciences)	Lontrel (Eclipse A)	Vantage (Eclipse B)		Roundup Ready Canola varieties	See component products	30 acres	$13.90
Elite (Nufarm	Poast Ultra	Mextrol	Merge (adjuvant)	Flax	See component products	20 acres or 8 ha	$21.65
FlaxMax (BASF Canada)	Curtail M	Poast Ultra		Flax	See component products	18-23 acres or 7.2-9.3 ha	$22.63 to $28.92
Harmony Total (E. I. duPont Canada)	Refine Extra	Horizon	Score (adjuvant)	Spring wheat (including durum)	Weeds controlled by Refine Extra plus wild oat, green foxtail	40 acres or 16 ha	$23.11
Horizon BTM (Syngenta)	Buctril M	Horizon	Score (adjuvant)	Spring wheat (including durum)	Weeds controlled by Buctril M plus wild oat.	20 acres or 8 ha	$23.64
K2 (E. I. duPont Canada)	Refine Extra	Everest	Add 2,4-D amine and a non-ionic surfactant	Spring wheat (NOT durum)	See component products	40 acres or 16 ha	$21.46 plus 2,4-D, surfactant
Muster Gold II (E. I. duPont Canada)	Muster	Assure II	Sure-Mix or Merge (adjuvant)	Canola, including herbicide tolerant canola	See component products	40 acres or 16 ha	$21.25
Pea Pack (Bayer CropScience)	Sencor	MCPA sodium salt		Pea	Lamb's-quarters, stinkweed, wild mustard, redroot pigweed, volunteer canola	40 acres or 16 ha	$5.95
Prevail Liquid (Dow AgroSciences)	Achieve Liquid	Curtail M	Turbocharge (adjuvant)	Spring wheat (including durum), barley	Wild oat, green foxtail, persian darnel; weeds controlled by Curtail M, plus Russian thistle	20 acres or 8 ha	$26.25
Pursuit Ultra (BASF Canada)	Poast Ultra	Pursuit		Field pea, CLEARFIELD canola varieties	See Poast Ultra plus Pursuit tank mix	40 acres or 16 ha	$26.34

**Not recommended for use on flax by itself.*

Note: *See the component products listed for information concerning staging, application information, safety precautions, the effect of weather and grazing, recropping, harvest interval and storage precautions. The more stringent recommendation of the two products should be followed. Mix products in the order listed.*

Herbicide Directory

2,4-D

Herbicide Group – 4
(Refer to page 27)

Company and Formulation:

	PCP #			
	Amine*		Ester**	
	500	600	600	700
IPCO	–	17511	–	20310
NuFarm	14725	–	14739	23508
United Agri-Products	9547	5931	9561	23563

* formulated as a solution.
**formulated as an emulsifiable concentrate.

Crops, Rates and Staging:

Application rates for individual products may vary from those listed. Refer to the label for product specific use rates. Rates greater than those listed may cause crop injury.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	MAXIMUM SAFE RATE (L/ACRE)	STAGE
Wheat, barley, spring rye	0.45 (500 g/L) 0.38 (600 g/L) 0.32 (700 g/L)	4 leaf to early flag leaf
Fall rye, winter wheat	0.45 (500 g/L) 0.38 (600 g/L) 0.32 (700 g/L)	In spring, apply after winter cereals begin grow but before emergence of the flag leaf. Winter wheat only may be treated in the fall for winter annual weed control if wheat is in the tillering phase of growth *(Caution: this use not registered and is at the risk of the user).*
Corn	0.42 (500 g/L) 0.35 (600 g/L)	Apply as an overall spray before corn is 6 inches (15 cm) tall and before the 6-leaf stage. After 6 inches (15 cm) use a directed spray.
Seedling and established grasses for forage and seed production*	0.45 (500 g/L) 0.38 (600 g/L) 0.32 (700 g/L)	Apply from the 3 leaf stage to emergence of the flag leaf of seedling grasses. For established grasses for seed production, apply in spring up to emergence of the flag leaf.
Established forage grass *(not for seed production)**	0.93 (500 g/L) 0.78 (600 g/L) 0.67 (700 g/L)	Apply in spring up to emergence of the flag leaf of established grasses, or in the fall after harvest.
Established grass pastures	1.81 (500 g/L) 1.51 (600 g/L) 1.30 (700 g/L)	No restrictions, apply when weeds are actively growing. For control of brush species, apply at time of rapid growth (usually May to mid-June, and September prior to colour change).
Turf	0.91 (500 g/L) 0.76 (600 g/L)	Apply in spring and early September on established turf. Do not use on bent grasses.
Fall stubble – Canada thistle suppression	0.91 (500 g/L) 0.76 (600 g/L) 0.65 (700 g/L)	Canada thistle (suppression only) - Apply when thistle plants are actively growing and have 6 to 8 inches (15 to 20 cm) of new growth. Regrowth will be present the following spring and in-crop treatments will be required.

* Note: Registered for use only with certain brands of 2,4-D; use of non-registered products is at the risk of the user.

Weeds, Rates and Staging:

Apply at lower rates when weeds are small (2 to 4 leaf stage) and actively growing. Higher rates are needed when weeds are larger, in heavy populations, or growing under stressful conditions (excessively cold, hot, dry or wet). Lower rates may be applied in late fall to control winter annual weeds.

Note: The rates listed differ slightly from product to product. Check individual product labels for exact use rates.

Susceptible Weeds:
0.27 to 0.45 L/acre (500 g/L)
0.22 to 0.38 L/acre (600 g/L)
0.19 to 0.32 L/acre (700 g/L)

Bluebur
Burdock
Cocklebur
False Flax
Flixweed (Late fall application or spring seedlings)
Goat's-beard
Kochia
Lamb's-quarters
Mustards (except dog and tansy mustard)
Narrow-leaved hawk's-beard (Late fall application or spring seedlings)
Plantain
Prickly lettuce
Ragweed
Russian pigweed
Russian thistle
Shepherd's-purse**
Stinging nettle
Stinkweed**
Sweet Clover
Thyme-leaved spurge
Volunteer canola (including all herbicide tolerant varieties)
Wild Radish
Wild Sunflower

Harder to kill weeds:
0.48 to 0.74 L/acre (500 g/L)
0.40 to 0.61 L/acre (600 g/L)
0.34 to 0.53 L/acre (700 g/L)

Blue lettuce*
Burdock (top growth only of bolting plants)
Canada thistle***
Common chickweed
Common groundsel**
Common peppergrass
Dandelion*
Flixweed (spring prior to bolting)
Knotweed
Lady's-thumb
Leafy spurge*
Mustard (Dog & Tansy)
Narrow-leaved Hawk's-beard (spring prior to bolting)
Oak-leaved goosefoot
Pineappleweed
Prostrate pigweed
Redroot pigweed
Russian thistle
Sheep sorrel
Tumble pigweed

Top Growth Control only (at rates for harder to kill weeds):

Biennial wormwood
Bull thistle
Buttercup
Curled dock
Field bindweed
Gumweed
Hedge bindweed
Hoary cress
Horsetail
Mouse-eared chickweed
Perennial Sow-thistle
Russian knapweed
Tartary buckwheat

* Control of seedlings at rates given above and top growth control only of established plants.

** Spring seedlings. Winter annual weeds - apply in late fall or early spring prior to bolting.

*** Suppression only - Apply when Canada thistle plants are actively growing and have 6 to 8 inches (15 to 20 cm) of new growth. Regrowth will be present the following spring and in-crop treatments will be required.

Cost (2004 suggested retail prices):

Amine 500 g/L - $6.25 to $6.47 per L
Amine 600 g/L - $7.41to $7.80 per L
LV Ester 600 g/L - $8.05 to $8.53 per L
LV Ester 700 g/L - $9.40 to $10.45 per L

Formulation Characteristics:

Formulation	Risk of Vapour Drift	Activity on Weeds	Risk of Crop Injury
LV Ester	Medium	Fast	Medium
Amine	Very Low	Medium	Low

Application Information:

Water Volume: Minimum 20 L per acre - ground application. Water rates depend on product and use. Consult label for details.

Higher application volumes (40 L/acre or greater), reduce the risk of crop injury.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Best weed control occurs when temperatures are above 21°C (daytime) or 10°C (nighttime) and humidity is above 70 percent. Do not apply if temperature exceeds 27°C.

Tank Mixes:

Mixes listed may not occur on all products. Check individual products for registered mixes.

Note: Following list is for mixes listed on 2,4-D labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Herbicides:

Spring wheat (including durum), winter wheat and Barley:
Avenge (2,4-D ester only)
Banvel (2,4-D amine only)

Spring wheat (including durum), and Barley:
Pardner and Sencor (2,4-D amine only)
Assert, Express DF, Refine Extra* and Ally* (2,4-D ester only)

* The addition of a non-ionic surfactant is required.

Fertilizers: None registered.

Insectcides: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: 2,4-D amine - within 4 hours will reduce control. 2,4-D LV ester - within 2 hours will reduce control.

Grazing: Do not permit lactating dairy animals to graze fields within 7 days of application. Do not harvest forage or cut for hay within 30 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Recropping: No restrictions the year after treatment.

Aerial Application: Some formulations may be applied by air. Check the label for detailed instructions.

Storage: 2,4-D LV ester may be frozen. 2,4-D amine requires heated storage.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

2,4-DB

Herbicide Group – 4

(Refer to page 27)

Company:

IPCO (Cobutox 600 – PCP#22404)
Nufarm (Embutox 625 – PCP#19217)
United Agri Products (Caliber 400 – PCP#16736)

Formulations:

Cobutox 600: 600 g/L 2,4-DB formulated as an emulsifiable concentrate. Container size - 10 L.

Embutox 625: 625 g/L 2,4-DB formulated as an emulsifiable concentrate. Container size - 10 L.

Caliber 400: 400 g/L 2,4-DB formulated as an emulsifiable concentrate. Container size - 10 L.

Crops and Staging:

NOTE - When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	STAGE
Seedling alfalfa, bird's-foot trefoil*	1 to 4 trifoliate leaf stage
Clover (alsike, red, white, Dutch but NOT sweet clover)*	As soon as possible after emergence of the 1st trifoliate leaf
Wheat, barley or oats (when used as a cover crop)	5 leaf to emergence of the flag leaf

CROP	STAGE
Field corn	15 inches (38 cm) to prior to tassling using drop nozzles.
Pastures containing forage legumes	After cutting or grazing and regrowth less than 3 inches (7.5 cm)

*with or without a cereal cover crop.

Seedling Forage Grasses*:

Apply from 2 to 4 leaf stage of:

Bromegrass (smooth)
Fescue (creeping red, meadow, tall)
Orchard grass
Timothy
Wheatgrass (crested, intermediate, streambank, tall)

* Not for seed production. Not for feeding in the establishment year.

Weeds and Staging:

Weeds controlled from the 2 to 4 leaf stage at lower recommended rates include:

Ball mustard
Lamb's-quarters
Ragweed
Redroot pigweed
Shepherd's-purse
Stinkweed
Wild mustard
Wormseed mustard

Weeds controlled at higher recommended rates include:

WEED	STAGE
Bull thistle	Rosette
Canada thistle*	6 in. (15 cm) to early bud
Chicory	Rosette
Curled dock**	Young and actively growing
Dandelion*	Prior to bud
Field bindweed*	Late summer
Green smartweed**	Seedlings
Horsetail*	4 to 5 inches (10 to 13 cm)
Narrow-leaved hawk's- beard	Apply at rosette stage after alfalfa has gone dormant
Oak-leaved goosefoot	Up to 2 leaf stage
Perennial sow-thistle*	Rosette
Lady's-thumb**	Seedlings
Plantain	Prior to flowering
Wild buckwheat	Up to 2 leaf stage
Yellow rocket	Late September to mid-October

* Top growth control
** Suppression
Refer to individual product labels for details on application rates to use for different weed species.

Cost (2004 suggested retail prices):

Cobutox - $21.00 per L
Embutox,- $21.30 per L
Caliber - $14.38 per L

Rates:

When applying to seedling forage legumes and grasses:

HERBICIDE	RATE (L/ACRE)	ACRES TREATED PER CONTAINER
Cobutox 600	0.73 - 0.93	13.7 - 10.8
Embutox 625	0.71 - 0.91	14.1 - 11.0
Caliber 400	1.1 - 1.4	9.0 - 7.0

When applying to corn and to pastures containing forage legumes:

HERBICIDE	RATE (L/ACRE)	ACRES TREATED PER CONTAINER
Cobutox 600	0.73 - 1.13	13.7 - 8.8
Embutox 625	0.71 - 1.11	14.1 - 9.0
Caliber 400	1.11 - 1.72	9.0 - 5.8

Application Information:

Water Volume: 61 to 81 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Severe damage to legumes can occur if high temperatures (more than 27°C) or high humidity prevail at the time of application. Do not apply under dry soil/drought conditions.

Tank Mixes:

Herbicides:

Underseeded Legumes: Cobutox (0.53 L/acre), or Caliber 400 (0.81 L/acre), plus MCPA amine (500 g/L) at 28 mL/acre or Embutox 625 (0.51 L/acre) plus MCPA sodium salt (300 g/L) at 47 mL/acre. This tank mix may increase crop damage (stunting). Follow all precautions and restrictions on both product labels.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on 2,4-DB labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour will reduce control.

Grazing: Do not graze or cut for feed for 30 days after applying Embutox or Cobutox. Do not graze treated crop or cut for feed in year of treatment after using Caliber.

Recropping: No restrictions the year after application.

Aerial Application: No restriction on label. Reduced water volumes used with aerial applications will lead to increased injury to forage legumes.

Storage: Do not freeze Caliber 400.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger Poison (Cobutox 600)

Caution Poison (Embutox 625)

Warning Poison (Caliber 400)

Absolute

Herbicide Group – 2,4

(Refer to page 27)

This product is a prepackaged tank mix of Odyssey at 17g/acre (page 159) and Lontrel at 0.17 L/acre OR Lontrel Dry (Lontrel Dry is only available with Absolute) at 80 g/acre (page 149). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the effect of growing conditions, and restrictions for the component products see the product pages listed above.

Company:

BASF Canada

Formulation:

Odyssey (PCP#25111): 35% imazamox and 35% imazethapyr formulated as a dispersible granule.
Absolute may be prepackaged with either:

Lontrel (PCP#23545): 360 g/L clopyralid formulated as a solution.
Container size – 8 x 86.5 g water soluble bags of Odyssey plus 2 x 3.4 L jug of Lontrel (one case treates 40 acres).

OR

Lontrel Dry (PCP#27306): 75% clopyralid formulated as a dispersible granule.
Container size – 4 x 173 g Odyssey plus 800 g Lontrel Dry, (one case treats 40 acres).

Crops and Staging:

CLEARFIELD canola varieties from the 2 to 6 leaf stage. Apply **only** to CLEARFIELD canola varieties; application to any other variety of canola or any other crop will result in crop death.

Weeds and Staging:

Grasses:

Apply from 1 to 4 main stem leaves, until tillers are visible.

Barnyard grass
Green foxtail
Persian darnel
Volunteer cereals (wheat, barley, oats)
Wild oats

Broadleaves:

Unless otherwise indicated, apply to broadleaf weeds up to 4-leaf stage.

Alsike clover ***
Canada thistle (rosette to pre-bud stage)*
Chickweed
Cleavers (up to 4 whorls)
Common groundsel***
Flixweed
Hemp-nettle
Kochia**
Lamb's-quarters**
Perennial Sow-thistle*
Redroot pigweed
Russian thistle
Scentless chamomile***
Shepherd's-purse
Smartweed
Stinkweed
Stork's-bill
Tufted vetch***
Volunteer alfalfa***
Volunteer canola (non-CLEARFIELD varieties)
Volunteer tame mustard
Wild buckwheat
Wild Mustard

* top growth control for 6-8 weeks
** suppression
*** not listed on Lontrel Dry label.

Cost (2004 suggested retail price):

$33.32 per acre

Rate:

Odyssey: 17.3 g/acre
Lontrel: 0.17 L/acre or
Lontrel Dry: 80 g/acre.
Merge: 0.5 L per 100 L of spray solution

Absolute must be applied with Merge adjuvant (not supplied). Mix to a final ratio of 0.5 L of Merge in 100 L of spray solution (0.5% Merge volume to Spray solution volume). At a spray volume of 40L/acre one 8.1 L jug of Merge will treat 40 acres.

One case treats 40 acres.

Do not apply more than once per year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan nozzles calibrated for 20-40 psi. Use a 50 mesh or coarser screen and filter system.

See component products for additional information including restrictions.

Accent*

Herbicide Group – 2
(Refer to page 27)

Company:

E. I. duPont Canada (PCP#25116)

Formulation:

75% nicosulfuron formulated as a water dispersible granule.

Container size - 133.6 g (4 x 33.4 g water soluble bags).

Crops and Staging:*

Field Corn - 1 to 8 leaf stage (six visible collars), coleoptile (short, blunt leaf) is counted as the first leaf.

Sweet corn (Krispy King, Jubilee and Jubilee Supersweet varieties only) – 1 to 6 leaf stage (4 visible collars).

** NOTE - Since applications to field and sweet corn in western Canada has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. **Application to corn is at the risk of the user.***

Weeds and Staging:

WEEDS	STAGING
Barnyard grass, fall panicum, green foxtail, yellow foxtail*, old witchgrass	1-6 leaves (up to 2 tillers)
Quackgrass	3-6 leaves (with extended leaf 4-8 inches (10 to 20 cm) long)
Wild oats	3-6 leaves

* suppression only

The most benefit is gained by applying at the earlier end of the leaf stage ranges.

Cost (2004 suggested retail price):

$24.32 per acre.

Rates:

Apply Accent at 13.5 g/acre. Add non-ionic surfactant (Citowett Plus, Agsurf or Agral 90) at 0.2 L per 100 L of spray solution.

One bag of Accent will treat 2.5 acres (1 ha).

One container will treat 10 acres (4 ha).

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 40 L per acre; optimum 56 to 77 L per acre.

Pressure: 25 to 40 psi (175 to 275 kPa).

Nozzles: Flat fan nozzles tilted forward at a 45° angle. Screens of 50 mesh or larger are recommended.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Poor weed control or crop injury may result if at the time of application, plants are under stress from disease, insect or nematode injury, carryover of herbicide from a previous years application, abnormally hot or cold weather, drought, water-soaked soils, hail damage or frost. Delay application until stress passes and both corn and weeds have resumed growth. When corn is injured by frost, wait 48 to 72 hours after normal growing conditions have resumed before applying Accent. Stress conditions after application may also result in injury or poor weed control.

Tank Mixes:

Herbicides:

Field corn only:

Banvel II (0.24 L/acre) plus surfactant.
Pardner (0.4 L/acre) plus surfactant.

Fertilizers: Do not mix with fertilizers.

Insecticides: None registered. Accent should not be applied to corn that has been treated with Counter, Cygard, Thimet, or Di-Syston. Leave seven days between the application of Accent and that of a foliar organophosphorous insecticide.

Fungicides: None registered.

Note: The above mixes are those listed on the Accent label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: Within 2 to 4 hours of application may result in reduced weed control.

Grazing: Do not graze treated crops or cut for hay.

Preharvest: Leave at least 30 days in field corn and 40 days in sweet corn from application to harvest.

Recropping: Corn and spring cereals may be seeded the year following Accent application. For all other crops, a field bioassay is recommended before planting.

Aerial Application: Do not apply by air.

Storage: Store product in original containers in a secure, dry area, away from other pesticides, food or feed.

Environment: Leave a 5 m buffer between the last spray path and woodlots or shelterbelts. Leave a 22 m buffer before wetland areas or water bodies.

Tank Cleaning:

Drain tank and hose down interior surfaces. Flush tank, hoses, boom, and nozzles with clean water for a minimum of 5 minutes. Fill spray tank with a water-ammonia cleaning solution (1 litre of a minimum 3% household ammonia for every 100 L of water). Flush hoses, boom and nozzles with the cleaning solution, then add more water to completely fill the tank. Circulate for 15 minutes, then flush hoses, boom and nozzles with the cleaning solution, and drain the tank. Remove and clean the nozzles and screens separately in a bucket containing the cleaning solution as above. If the spray equipment is to be used to spray crops other than corn, repeat the above process and thoroughly wash the spray mixture from the outside of the spray tank and the boom. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom. Prior to using the sprayer again, flush the tank, boom and hoses for 5 minutes with fresh water. Do not clean equipment where cleaning solution could flow towards water bodies, ditches, cropland, shelterbelts, or areas where people are likely to frequent or walk.

For additional information, refer to page 14.

Hazard Rating:

KEEP OUT OF REACH OF CHILDREN.
Caution Eye Irritant

Avoid breathing spray mist.
Avoid contact with skin, eyes and clothing.

Achieve Liquid

Herbicide Group – 1
(Refer to page 27)

Company:

Syngenta (PCP#27011)

Formulation:

Achieve Liquid: 400 g/L tralkoxydim formulated as a suspension concentrate.
Container size – 8 L plus 8 L of Turbocharge adjuvant.

Crops and Staging:

No staging restrictions.

Spring wheat (including durum), winter wheat, barley, triticale, rye (spring and fall)

May be used on wheat and barley crops undersown to forage legumes (if not tank mixed with a broadleaf herbicide):

- Alfalfa
- Bird's-foot trefoil
- Clovers
- Sanfoin

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Wild oats - 1 to 6 leaf stage (total leaves including tillers), with a maximum of 2 tillers.

Volunteer tame oats - 1 to 6 leaf stage.

Green and yellow foxtail - 1 to 5 leaf stage (total leaves including tillers), with a maximum of 1 tiller.

Barnyard grass, Persian darnel - 1 to 4 leaf stage (total leaves including tillers).

Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

Cost (2004 suggested retail price):

$16.63 per acre.

Rates:

0.2 L per acre. One case treats 40 acres. Add Turbocharge at a rate of 0.5 L per 100 L spray solution.

If water analysis shows application water bicarbonate levels are 400 ppm or greater, add ammonium sulfate at 1.65 to 3.3 lb (0.75 to 1.5 kg) per 100 L of spray water prior to mixing.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:
Ground: 20 to 40 L per acre. Application in less than 20 L per acre water volume may result in mixing problems or unacceptable crop injury.
Aerial: 14 to 18 L/acre.

Pressure:
Ground: 40 psi (275 kPa).
Aerial: 20 to 40 psi (140 to 275 kPa)

Nozzles: 80° or 110° flat fan nozzles. All strainer and nozzle screens must be 50 mesh or coarser.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Cereal crops that have set tillers may incur injury (yellowing and/or stunting) if applications are made within 48 hours of freezing temperatures. Cereal crops that have not set tillers may be injured if exposed to temperatures of 4°C or less up to 48 hours before or after application. Tank mixing with a broadleaf weed herbicide under adverse conditions may increase severity of crop injury. Crops under stress from foliar diseases or low fertility are more susceptible to injury from application. Temporary crop injury may occur when Achieve tank mixes (particularly Estaprop, Turboprop, Dichlorprop D, and Buctril M + MCPA) are applied under extreme environmental conditions (dry or wet, cool or hot weather) resulting in crop stress. Control of grasses could be reduced when they are stressed due to drought, heat, lack of fertility, flooding or prolonged cool temperatures.

Tank Mixes:

For all tank mixes with Achieve, add Turbocharge at a rate of 0.5 L per 100 L of spray solution.

Herbicides:

Do NOT tank mix Achieve products with a broadleaf herbicide when applying to underseeded forage grasses or legumes.

In wheat and barley only:

Tank mix Partner	CROPS						
	Spring wheat	Durum	Winter wheat	Barley	Spring Rye	Fall Rye	Triticale
2,4-D ester (0.38 L/acre)Δ	✓	✓	✓	✓	✓	✓	
Attain (80 acres per case)**	✓			✓			
Buctril M (0.4 L/acre)*	✓	✓	✓	✓		✓	
Curtail M (0.81 L/acre)	✓	✓		✓			
Dichlorprop + 2,4-D (0.71 L/acre)*	✓	✓	✓	✓			
Lontrel (0.11 L/acre) + MCPA ester (0.45 L/acre)	✓	✓		✓			
MCPA ester (0.45 L/acre)	✓	✓	✓	✓	✓	✓	
Pardner (0.40 L/acre)*	✓	✓	✓	✓		✓	✓
Prestige (20 acres per case)**	✓			✓			
Thumper (0.40 L/acre)*	✓	✓		✓			

Δ 600 g/L formulations

* Tank mixes may result in some temporary initial injury under adverse environmental conditions.

** Temporary crop injury can occur if applied prior to the 4 leaf stage. A reduction in wild oat control may occur with this mix.

Do NOT tank mix Achieve with herbicides or formulations of herbicides not listed above as loss of grass control may result. When applying broadleaf herbicides not listed above, in the same field, always apply Achieve first. Apply the broadleaf product no sooner than seven days after application of Achieve.

Fertilizers: None registered.

Insecticides: Decis Flowable at 49 mL per acre. This mix may also be mixed with Buctril M or Pardner,.

Fungicides: None registered.

Note: The above mixes are those listed on the Achieve label only. To check for other possible mixes see the blue chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour will reduce control.

Grazing: Straw from treated crops may be fed to livestock.

Preharvest: Leave at least 60 days from application to harvest.

Recropping: Do NOT replant treated areas to tame oats or corn for at least 4 weeks after application.

Aerial Application: Achieve may be applied by air in 14 to 18 L/acre. Do not apply within 50 m of fish bearing waters and wildlife habitat.

Storage: Store in a dry place. Do not freeze.

Environment: Do not apply within 15 m by ground (50 m by air) of fish bearing waters, wetlands and wildlife habitat.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Caution Eye Irritant

Achieve Liquid Gold

Herbicide Group – 1,4,6
(Refer to page 27)

This product is a prepackaged tank mix of Achieve (page 54) and Buctril M (page 81). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Syngenta

Formulation:

Achieve Liquid Gold contains 3 components:

Achieve Liquid (PCP#27011): 400 g/L tralkoxydim formulated as a suspension concentrate. Container size – 4 L

Buctril M (PCP#18022): 280 g/L bromoxynil and 280 g/L MCPA ester formulated as an emulsifiable concentrate. Container size - 8 L.

Turbocharge adjuvant (PCP#23135). Container size - 4 L.

Crops and Staging:

Spring wheat (including durum), barley - 2 leaf to early flag leaf stage.

Winter wheat, rye (spring and fall) - from the time growth commences in the spring to the early flag leaf stage.

Weeds and Staging:

Grassy weeds :

Wild oats - 1 to 6 total leaves to a maximum of 2 tillers

Barnyard grass, Persian darnel - 1 to 4 total leaves including tillers

Green & yellow foxtail
- 1 to 5 total leaves, maximum of 1 tiller

Volunteer tame oats - 1 to 6 leaf stage.

Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are controlled before tillering.

Broadleaf weeds:

Up to the 4 leaf stage:

American nightshade
Ball mustard
Bluebur
Cocklebur
Cow cockle
Flixweed
Green smartweed
Kochia*
Lady's-thumb
Night-flowering catchfly
Pale smartweed
Redroot pigweed
Russian thistle*
Scentless chamomile (spring annuals only)
Shepherd's-purse
Velvetleaf**
Volunteer canola
Volunteer sunflower

Up to the 6 leaf stage:

Wild tomato

Up to the 8 leaf stage:

Common groundsel
Common ragweed
Lamb's-quarters
Stinkweed
Tartary buckwheat
Volunteer tame buckwheat
Wild buckwheat
Wild mustard
Wormseed mustard

Perennials (top growth control):

Canada thistle
Perennial sow-thistle

* Spray before plants are 2 inches (5 cm) in height.
** Spray before plants are 3 inches (8 cm) in height.

Cost (2004 suggested retail price):

$23.27 per acre.

Rates:

One case treats 20 acres. Add Turbocharge at a rate of 0.5 L per 100 L spray solution.

If water analysis shows bicarbonate levels are 400 ppm or greater, add ammonium sulfate at 1.65 to 3.3 lb (0.75 to 1.5 kg) per 100 L water.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 20 to 40 L per acre. Do not apply with air assist sprayers set to apply less than 20 L per acre water volume as mixing problems or unacceptable crop injury could occur.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° degree flat fan nozzles. All strainer and nozzles screens must be 50 mesh or coarser.

Tank Mixes:

Herbicides: MCPA ester (0.22 L/acre) (500 g/L formulations). A reduction in green foxtail control may occur with this tank mix.

Insecticides: Decis Flowable (49 mL/acre).

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Adrenalin/Adrenalin SC

Herbicide Group – 2,4
(Refer to page 27)

Company:

BASF Canada

Formulations:

Adrenalin contains two separate components:
AC 299,263 (PCP#26705): 11.4% imazamox present as an ammonium salt and formulated as a solution.
2,4-D LV700 (PCP#23508): 660 g ae/L 2,4-D LV ester formulated as an emulsifiable concentrate.
Container size – Each case includes 2 jugs that are split internally. Each jug contains 1.34 L of AC 299,263 and 6.8 L of 2,4-D LV700 Ester.

Adrenalin SC (PCP#27879): 20 g/L imazamox and 560 g/L 2,4-D ester formulated as a suspension concentrate.
Container size 2 X 8L.

Crops and Staging:

CLEARFIELD wheat varieties - 4 to 6 leaf stage. Apply **only** to CLEARFIELD wheat varieties: application to any other variety of wheat or any other crop will result in crop death.

NOTE: The labeled rate of 2,4-D ester in these products is greater than the recommended safe rate for 2,4-D on wheat and injury to the crop is possible.

Weeds and Staging:

Grasses:
Apply from 1 to 4 main stem leaves, to a maximum of two tillers.

Green foxtail
Volunteer cereals (non-Clearfield wheat, barley)
Yellow foxtail[†]
Wild oats

[†] Adrenalin SC formulation only

Broadleaves:
The following weeds will be controlled up to the 4-leaf stage under a wide variety of conditions. Application at an earlier weed stage provides the best yield response.

Annual Smartweed species (including Lady's Thumb)[†]
Annual sow-thistle
Bluebur
Cleavers[*]
Cocklebur
Daisy fleabane
False flax
Flixweed
Goat's-beard
Kochia
Lamb's-quarters
Mustards (except dog and tansy)
Plantain
Prickly lettuce
Ragweeds
Redroot pigweed
Russian pigweed
Russian thistle
Shepherd's purse
Stinging nettle
Stinkweed
Sweet clover
Volunteer canola (all varieties)
Wild buckwheat[*]
Wild mustard
Wild radish
Wild sunflower

[*] suppression
[†] Adrenalin SC formulation only

As packaged the following weeds are controlled to the two leaf stage when conditions are optimal for growth. Control may be reduced if weeds are found in high population densities, or are growing under stressful conditions (drought, cold, heat).

Biennial wormwood[Δ]
Blue lettuce[Δ]
Chickweed
Curled dock[Δ]
Dandelion[Δ]
Groundsel
Gumweed[Δ]
Hedge bindweed[Δ]
Hemp-nettle[†]
Horsetail[†]
Knotweed
Leafy spurge[Δ]
Mustards (dog and tansy)
Oak-leaved goosefoot[†]
Peppergrass[Δ]
Pineappleweed
Prostrate pigweed
Purslane
Russian knapweed[Δ]
Sheep sorrel[Δ]
Smartweed (including lady's-thumb)[**]
Wild buckwheat
Yellow rocket

[Δ] Only seedlings of biennial and perennial weeds will be controlled. Established plants emerging from roots will not be controlled.
[**] Adrenalin only, listed as full control on the Adrenalin SC label.
[†] Adrenalin SC formulation only

As packaged the 2,4-D component of Adrenalin and Adrenalin SC will give top growth control of the following weeds . Control may be reduced if weeds are found in high population densities, or are growing under stressful conditions (drought, cold, heat).

Bull thistle
Burdock
Buttercup
Canada thistle
Curled dock
Field bindweed
Hoary cress
Mouse-eared chickweed**
Perennial sow-thistle
Tartary buckwheat

** Adrenalin formulation only

Cost (2004 suggested retail price):

$23.83 per acre.

Rate:

Aderenalin:

AC 299,263: 71 mL per acre.

2,4-D LV700: 0.34 L per acre as labelled (0.36 L per acre as packaged).

(One case treats 37.7 acres).

Adenalin SC: 0.4 L per acre (One case treats 40 acres).

Agral 90 Ag-Surf or ***Surf 92*** at.0.25 L per 100L of spray solution.

Use of an anti-foam agent is suggested.

Do not apply more than once per year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: No nozzle preference is indicated. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury will occur as a result of spray overlap. AVOID SPRAYER OVERLAP.

Do not spray if freezing or near freezing (+2 Celsius or lower) temperatures are forecast. Treat CLEARFIELD wheat during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced. Application must be made before the crop canopy shields the weeds.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: DO NOT graze the treated crop within 14 days of application or cut for hay within 42 days of application.

Preharvest Interval: Do not apply within 79 days of harvest.

Recropping: The following crops may be grown safely the year following an application:
Field peas, CLEARFIELD canola varieties, spring wheat, durum, barley, oats.
Conduct a field bioassay the year before growing any other crop than those listed above.

Aerial Application: Do not apply by air.

Storage: Do not freeze. Store in a cool, dry place above 5°C.

Environment:
Adrenalin/Adrenalin SC - These products are highly toxic to non-target plants. Avoid overspraying and applying in situations where drift may occur. Leave a buffer zone of 11 metres between the last spray swath and sensitive areas. Areas to be buffered include shelter belts, hedgerows, wetlands, woodlots, vegetated ditchbanks, ponds, streams, potholes, sloughs, and other cover on the edges of fields.
Adrenalin SC – This product contains petroleum distillate which is toxic to aquatic organisms. Avoid contamination of aquatic systems during application and cleaning operations.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Keep Out of Reach of Children (AC 299,263), Adrenalin SC

Warning Poison (2,4-D LV700)

Ally Toss-N-Go

Herbicide Group – 2
(Refer to page 27)

Company:

E. I. duPont Canada Inc. (PCP#24388)

Formulation:

60% metsulfuron methyl formulated as a dry flowable. Container size - 122 g package (4 x 30.5 g water soluble bags).

Crops, Rates and Staging:

Cereals – up to 3 g/acre:
Wheat (spring and durum), barley - 2 leaf to emergence of the flag leaf.

Forage Grasses (seedling or established)* - up to 3 g/acre: Apply from the 2 leaf to flag leaf stage and before canopy is dense enough to prevent thorough leaf coverage.

For forage or seed production*:
Crested wheatgrass
Intermediate wheatgrass
Orchard grass

For seed production only:
Creeping red fescue*

Pasture and Rangeland – Up to 12 g/acre

** NOTE - Since applications to forage grasses have been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance.* ***Application to forage grasses is at the risk of the user.***

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Cereals and forage grasses:

Unless otherwise indicated, apply to weeds at the 2 to 4 leaf stage.

Alone or in tank mix with a grass control herbicide use 3 g/acre. Use 2 to 3 g/acre (One 122 g package treats 40 to 60 acres) of Ally when tank mixing with 2,4-D or MCPA Add non-ionic surfactant such as Agral 90, Agsurf, Companion, Super Spreader or Citowett Plus at 0.2 L per 100 L spray volume.

Weeds Controlled:
Ball mustard
Bluebur
Chickweed
Common groundsel
Corn spurry
Cow cockle
Flixweed
Green smartweed
Hemp-nettle
Kochia
Lady's-thumb
Prostrate pigweed
Redroot pigweed
Scentless chamomile
Shepherd's-purse
Stinkweed
Stork's-bill
Tartary buckwheat
Volunteer canola*
Wild mustard

* CLEARFIELD varieties will be controlled only with the addition of 2,4-D or MCPA.

Weeds Suppressed:
Canada thistle**
Lamb's-quarters
Russian thistle
Sow-thistle (annual and perennial)
Toadflax
Wild buckwheat***

** Apply when thistles are less than 6 inches (15 cm) tall.
*** Apply to wild buckwheat up to the 3 leaf stage.

Pastures and Rangelands:
Western snowberry - 10 g/acre
Wild rose~ 12 g/acre

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$3.94 to $5.91 per acre.

Application Information:

Water Volume:
Cereals and forage grasses: Minimum 40 L per acre.
Pastures and Rangelands: 40 to 91 L per acre.

Pressure: No application pressures are recommended by the manufacturer. Typical application pressures for standard flat fan nozzles are from 35 to 40 psi (240 to 275 kPa).

Nozzles: Flat fan nozzles. Use 50-mesh screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Ally may injure crops stressed by heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures, drought, or water-saturated soils, either before or after application. Weed control will be reduced under dry, cold conditions.

Tank Mixes:

Do not mix the soluble bags with any substance containing boron or which releases chlorine.

Herbicides:

In wheat and barley:
2,4-D Amine or Ester (0.34 to 0.45 L/acre of 500 g/L formulation), plus surfactant.
Avenge (1.72 L/acre), no surfactant required.
Avenge + MCPA Ester, no surfactant required.
MCPA Amine or Ester (0.28 to 0.45 L/acre of 500 g/L formulation), plus surfactant.

In spring wheat (including durum):
Horizon (95 or 115 mL/acre) plus Score adjuvant
Puma^{120} Super (0.15 to 0.31 L/acre), no surfactant required.

In Creeping Red Fescue:
Assure II (0.2 to 0.3 L/acre) plus Sure-Mix adjuvant.

Consult tank mix partner labels for additional crop staging and variety restrictions.

Fertilizer: None registered. Do not mix the soluble bags with fertilizers.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Ally label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rain within 4 hours of application of tank mixes with 2,4-D amine, 2 hours of application of tank mixes with 2,4-D ester, will reduce weed control.

Grazing: No restrictions.

Recropping: Refer to table on the following page. The following recropping intervals, based on soil pH, should be considered as guidelines only. Ally residues may affect crops for a longer period of time than outlined in the following table. Add 12 months to recommendations if less than 5 inches (130 mm) of rainfall in brown and dark brown soils or less than 10 inches (250 mm) rainfall in black or grey wooded soils in any year following application.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. May be frozen.

Caution:

Ally residues can persist for long periods, potentially limiting recropping options. Degradation of Ally is dependent on the pH, moisture, and temperature of the soil. Refer to the label for details on rotation and minimum recropping intervals.

MINIMUM RECROPPING INTERVAL (MONTHS)

SOIL PH	BARLEY, WHEAT	OAT*	CANOLA*	FLAX*	LENTILS	CANARY-SEED	YELLOW MUSTARD
less than 7.0	10	10	10 (22)	10 (22)	34	48	48
7.0 - 7.9	10	10 (22)	22 (34)	34	48	48	48

* Figures in brackets refer to recropping intervals in brown and dark brown soil zones. ALL OTHER DATA refer to recropping intervals in all soil zones.

On black and grey wooded soils with pH of 7.5 or less, fescue may be planted 10 months after application and alfalfa, red clover, peas and flax may be planted 22 months after application. Do not use on soils with pH greater than 7.9.

Effects of Ally residues on crops other than those listed in the table have not been fully evaluated. Because of the length of recropping restrictions and the lack of information on many rotational crops, Ally is not recommended for use on farms where special crops are grown (such as fababeans, beans, sunflowers, buckwheat, corn, sugar beets, etc.).

Tank Cleaning:

Ally can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray Ally should be flushed out immediately after Ally is used. The manufacturer recommends that sprayers used to apply this product be flushed 2 times with a water/household ammonia rinse (1 L of 3% ammonia per 100 L water). All nozzles, screens and filters should be removed and cleaned after applying this product.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

For additional information refer to page 14.

Hazard Rating:

☠ Caution Poison

Caution Eye Irritant

Amitrol 240

Herbicide Group – 11
(Refer to page 27)

Company:

Nufarm (PCP#25684)

Formulation:

231 g/L amitrole formulated as a liquid.
Container size - 10 L.

Crops, Rates and Staging:

Fall Stubble: Perennial weed control prior to spring seeding. No planting restrictions for barley, canola, field corn, field pea, soybean, wheat, or white bean, but leave 8 months between application and the seeding of any other crops.

Alfalfa stand renovation/removal: 4 to 6 inches (10 to 15 cm) high.

Pastures (spot treatment only): For non-selective patch treatment of dandelion, Canada thistle, perennial sow-thistle, hoary cress, milkweed, poison ivy and toadflax apply 0.165 L in 25 L of water to treat a 10 m x 10 m area. For treatment of leafy spurge and horsetail, apply 0.460 L in 25 L of water to treat a 10 m x 10 m area. Do not mow treated plants for 3 weeks after application.

Established shelterbelts: up to 11.3 L per acre - Keep spray away from tree foliage or trunks.

Pre-seeding:

CROP	RATE (L/ACRE)	Delay seeding after application
Barley, wheat, canola	1.7	0 to 1 days
Field pea	1.7	5 to 7 days
Soybean (low rate)	1.7	6 days
Field corn, white bean, soybean (high rate)	3.3 to 6.7	10 to 14 days

Avoid using rates higher than 6.7 L per acre for preplant applications on very light textured soils with low organic matter, as crop damage can occur.

Fallow areas: Apply according to weed stage and rates below.

Weeds, Rates and Staging:

Fall stubble: Canada thistle, perennial sow-thistle - Spray when thistle has 4 to 6 inches (10 to 15 cm) of new growth. Do not cultivate for 2 weeks after application. Do not apply after October 1. Do not replant crops in treated areas within 8 months of application except those registered for pre-seeding uses.

Pre-seeding: Dandelion and annual weeds Apply 1.7 L per acre to actively growing weeds less than 10 cm tall or across. Do not cultivate for 10 to 14 days after treatment.

Fallow, Pastures and Shelterbelts:

WEED	RATE (L/ACRE)	WEED STAGE
Canada thistle	5.0 to 6.7	Early bud to bloom stage.
Cattails	15.2 to 18.5	After seed heads have formed.
Dandelion	1.7 to 5.0	Young and actively growing plants.
Hoary cress	7.6 to 11.3	Advanced rosette and bud stage.
Horsetail	5.0 to 6.7	Actively growing plants.
Leafy spurge	15.2 to 18.5	Advanced flowering to early seed set
Milkweed	7.6 to 11.3	Early summer after majority of shoot emergence.
Perennial sow-thistle	5.0 to 6.7	Early bud to bloom stage.
Poison ivy	3.7	Fully developed green foliage.
Toadflax	7.6 to 11.3	Advanced rosette to prebud.

Cost (2004 suggested retail price):

$6.70 per L.

Application Information:

Water Volume:
Fall stubble: 20 to 81 L per acre.
Pastures, shelterbelts: 40 to 121 L per acre. For poison ivy, apply 202 to 405 L per acre.
Pre-seeding: 20 to 81 L per acre.

Pressure: 45 psi (Less than 300 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Less than acceptable results may occur in dry weather.

Tank Mixes:

Herbicides: Mix 1.68 L per acre Amitrol 240 with 0.5 L per acre Roundup for improved control of certain weeds, including dandelion. Follow directions on the Amitrol 240 and Roundup labels for timing and use precautions.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze treated crops or weeds or use for hay or feed.

Recropping: Do not plant any crop for 8 months following application except those registered for pre-seeding uses.

Aerial Application: Do not apply by air.

Storage: Do not store where temperatures may exceed 50°C or near open flames. Do not store below 4°C.

Environment: Do not contaminate any body of water. Use cautions to prevent spray, spray mist, or vapours from drifting off target. Spray drift may cause damage to crops or vegetation.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Caution eye irritant

Avoid breathing spray mist or vapours.

Do not re-enter treated areas within 12 hours.

Approve

Herbicide Group – 4, 6

(Refer to page 27)

This product is a prepackaged tank mix of Koril (Page 79) and 2,4-DLV700 Ester (Page 47). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Nufarm

Formulation:

The Approve package has 2 components:

Koril 235 (PCP#25341): 235 g/L bromoxynil formulated as an emulsifiable concentrate

2,4-D LV700 (PCP#23508): 660 g/L of 2,4-D ester formulated as an emulsifiable concentrate

Container size: Koril – 9.71 L 2,4-D LV700 Ester – 3.43 L.

One case treats 20 acres

Crops and Staging:

Spring wheat (including durum), barley - 4 leaf to flag leaf stage.

Weeds and Staging:

Weeds controlled at the 1 to 4 leaf stage:

- American nightshade
- Annual smartweeds (green, pale, lady's-thumb)
- Ball mustard
- Bluebur
- Cocklebur
- Cow cockle*
- Flixweed
- Kochia**
- Night-flowering catchfly
- Ragweed (common)
- Redroot pigweed*
- Shepherd's-purse
- Stinkweed*
- Russian thistle**
- Velvetleaf***
- Volunteer sunflower
- Wild mustard*

Weeds controlled at the 1 to 8 leaf stage:

- Common groundsel
- Lamb's-quarters
- Tame buckwheat
- Tartary buckwheat
- Stinkweed*
- Wild buckwheat
- Wild mustard*

* Controlled only when plants are actively growing. Plants under stress, or larger than the 4 leaf stage will not be controlled.

** Apply before plants are 2 inches high.

*** Apply before plants are 3 inches high.

Cost:

$6.35 per acre

Rates:

Koril 0.49 L per acre

2,4-D Ester 0.17 L per acre

One case treats 20 acres

Application Information:

Water Volume:

Ground: Minimum 40 L/acre

Aerial: Minimum 8.1 L/acre. Optimum 16.2L for heavy weed populations.

Pressure: 45 psi (275 kPa)

Nozzles: 80° or 110° flat fan nozzles. All strainer and nozzle screen must be 50 mesh or coarser.

Tank Mixes:

None Registered.

Restrictions:

Use the most limiting restrictions listed for either Koril (page 79) or 2,4-D (page 47).

Assert 300 SC

Herbicide Group – 2

(Refer to page 27)

Company:

BASF Canada (PCP#21032)

Formulation:

Assert: 300 g/L imazamethabenz formulated as a suspension concentrate. Container size - 2 x 10.8 L jugs per case.

pH adjuster: 94.5% sodium bisulfate formulated as a soluble granule. Container size - 2 x 0.5 kg bags (1 bag per 10.8 L jug of Assert).

Crops, Rates and Staging:

pH adjuster: 1 packet per jug of Assert to be used.

Assert up to 0.67 L/acre (16.1 acres per jug):
Barley, spring wheat (including durum) - 1 to 6 leaf stage.
Annual ryegrass (seed production only) – 4 to 6 leaf stage.

Assert at 0.32 L/acre (32 acres per 10.8 L jug):
Sunflower - plants (not under drought stress) that are in the 2 to 8 leaf stage. Crop must be less than 15 inches (38 cm) tall except for semi-dwarf varieties, which must be less than 12 inches (30 cm), and dwarf varieties, which must be less than 4 inches (10 cm). Stunting and head deformation can occur from applications made beyond recommended stages.

Do not apply Assert to the same field more than once in two years.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Weeds, Rates and Staging:

Assert at 0.32 L/acre rate:
Stinkweed and wild mustard - up to 6 leaves.

Assert at 0.54 L/acre (20.3 acres per 10.8 L jug – weeds above plus:
Wild and tartary buckwheat (suppression) – up to 4 leaves.
Wild oats – majority from 1 to 3 leaves.

Assert at 0.67 L/acre rate – weeds above plus:
Wild oats - 1 to 6 total leaves (including tillers) up to 2 tillers.
Wild oats when tank mixing with a broadleaf partner.

Cost (2004 suggested retail price):

Cereals and Annual Ryegrass - $13.65 to $17.21 per acre
Sunflower - $8.66 per acre (wild mustard only).

Application Information:

Water Volume: In cereals only, Assert may be applied in 20 to 40 L of water per acre when applied alone or when tank mixed with dichloroprop+2,4-D, 2,4-D ester, or MCPA ester. For all other applications, apply in 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply Assert 24 hours before or after a frost. It works best at warm temperatures. Performs relatively consistently under dry conditions. If cold, wet soil conditions persist in the days after application, retillering of wild oats may occur. Do not apply to drought stressed sunflowers.

Tank Mixes:

Herbicides:

Assert must be applied at 0.53 to 0.67 L/acre when tank mixed with other herbicides for additional broadleaf weed control, depending on soil zone.

In spring wheat (including durum) and barley:
2,4-D Ester* (up to 0.32 L/acre) (700 g/L formulations)
Curtail M (0.80 L/acre)
Dichlorprop + 2,4-D (0.7 L/acre) (Estaprop & Turboprop 600 only)*
Express Pack*** (40 acres per case)
Frontline (40 acres per case)
MCPA Ester* (up to 0.45 L/acre) (500 g/L formulations)
Puma^{120} Super** (0.115 L/acre)
Puma^{120} Super** + MCPA Ester, or Refine Extra (rates listed)
Refine Extra*** (8 g/acre)
Refine Extra*** (8 g/acre) + MCPA Ester (0.28 to 0.45 L/acre) (500 g/L formulations)
Spectrum (20 acres per case)
Unity*** (40 acres per case)

In spring wheat (NOT including durum) and barley:
Attain (40 acres per case)
Prestige (20 acres per case)

In spring wheat and durum wheat only:
Dichlorprop + 2,4-D (0.7 L/acre) (Turboprop 600 only)

* apply in 23-45 L/acre. For all other tank mixes use 45 L/acre.

** use the 0.54 L/acre rate of Assert when tank mixing with Puma^{120} Super.

*** when tank mixing dry broadleaf products, add products to the tank in the following order: dry broadleaf products, acidifier, Assert, and other liquid herbicides if required. For repeat tanks, dry broadleaf products need to be mixed with water to form a slurry prior to adding to the remaining spray solution in the tank.

Refer to Assert label for specific mixing order and application details when tank-mixing. Refer to tank mix partner for additional crop staging restrictions. Tank mixes containing Refine Extra must be used within 12 hours or product degradation may occur.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Assert label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Grazing: Do not graze treated fields or cut treated forage for silage or hay. Mature barley and wheat grain or straw from fields treated with Assert can be fed to livestock. Do not feed or graze treated annual ryegrass.

Preharvest Interval: Do not apply beyond the recommended crop stage.

Recropping:

Do not apply Assert to the same field more than once in two years.

Year After Application	Black and Grey Wooded Soils	Brown and Dark Brown Soils
Year 1	Spring wheat (including durum), barley, canola, field peas, flax, sunflowers	Spring wheat (including durum), CLEARFIELD canola, barley, sunflowers
Year 2	Spring wheat (including durum), barley, canaryseed, canola, field peas, flax, oats, sunflowers	

Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop not listed in the table. Lentils are known to be particularly sensitive to Assert residues in the soil. The additive effect of soil residues from the use of Assert and sequential applications of Ally or Unity (not for use in Manitoba) herbicides on the same land area has not been determined. Crop rotation guidelines are not known and injury to rotational crops other than wheat (excluding durum) may occur. Plant only wheat (excluding durum) on fields where these herbicides have been used until a field bioassay demonstrates other crops can be grown successfully.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environment: Leave a distance of 15 m between the outside boundary of the sprayed area and bodies of water (lakes, ponds, sloughs, streams, rivers, etc.).

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

[illegible]

Assure II

Herbicide Group – 1

(Refer to page 27)

Company:

E. I. duPont Canada (PCP#25462)

Formulation:

96 g/L quizalofop-p-ethyl formulated as an emulsifiable concentrate. Container size – 8 L.

Sure-mix adjuvant. Container size – 8 L.

Crops and Staging:

No leaf stage restrictions, but do not apply beyond preharvest intervals listed in the table:

CROP	PREHARVEST INTERVAL (DAYS)
Canola	64
Dry bean (navy and pinto)*	65
Flax (including Solin)	82
Soybeans	80
Lentils	65
Peas (field and processing)	65

Forage Crops (no preharvest interval restrictions): alfalfa (seed production), and seedling bird's-foot trefoil, sainfoin, clovers (white, red, alsike and sweet) for seed production* and creeping red fescue for seed*.

* *NOTE - Since applications to these crops have been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance.* ***Application to these crops is at the risk of the user.***

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Costs (2004 suggested retail price):

$12.22 to $25.38 per acre.

Application Information:

Water Volume: Minimum 40 L per acre. Up to 162 L per acre of water may be used under heavy populations to improve coverage.

Pressure: 30 to 40 psi (210 to 275 kPa).

Nozzles: Flat fan with 50 mesh or coarser nozzle screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop injury may occur if crops are stressed because of drought or flooding. Less than acceptable weed control may be expected if weeds are under stress because of drought, flooding or cool weather.

Weeds, Rates and Staging:

Add Sure-Mix or Merge at 0.5 to 1.0 L per 100 L. Use 1.0 L per 100 L when wild oats or quackgrass are present in the field or when growing conditions are poor.

WEED	STAGE	RATE	
		L/ACRE	ACRES PER CASE
Wild oat*	1 to 5 leaf prior to tillering	0.15	54
	up to 2 tillers	0.20	40
Green foxtail	2 leaf to early tillering	0.15	54
Volunteer wheat, barley & oat	2 leaf to early tillering	0.15	54
Volunteer corn	2 to 6 leaf stage	0.15	54
Barnyard grass, Yellow foxtail, Proso millet, Fall pannicum, Old witchgrass	2 leaf to early tillering	0.20	40
Quack grass suppression	2 to 6 leaf stage	0.20	40
Quack grass - season long control	2 to 6 leaf stage	0.30	26

* Best results are likely to occur if applications are made before tillering begins. Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Tank Mixes:

Herbicides:

In Canola: Muster (8 to 12 g/acre)

In Soybeans:

Pinnacle (2.2 to 3.3 g/acre).

Assure II (0.25 L/acre) plus Pinnacle (2.2 to 3.3 g/acre) plus Basagran Forte (0.71 to 0.91 L/acre) plus Merge or SureMix to control grasses controlled by Assure II alone at 0.20 L/acre plus the broadleaf weeds controlled by Pinnacle and Basagran Forte.

In Established creeping red fescue for seed: Assure II at 0.2 to 0.3 L/acre may be tank mixed with Ally (3 g/acre). Allow 24 hours after application before applying a broadleaf herbicide. If the broadleaf herbicide is applied first, wait 7 days before application of Assure II.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Assure II label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Grazing: Do not graze treated crops or cut for feed prior to crop maturity.

Preharvest Interval: See Crop chart above.

Recropping: No restrictions the year after treatment.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environment: Do not apply within 15 m of sensitive habitats such as shelterbelts, wetlands, sloughs, or woodlots.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger – Corrosive to eyes

Warning eye and skin irritant

Atrazine

Herbicide Group – 5

(Refer to page 27)

Company:

Syngenta
(Aatrex Liquid – PCP#18450).

United Agri Products
(Atrazine 480 – PCP#20997, Atrazine 90 WG – PCP#23583).

Formulations:

Aatrex Liquid/Atrazine 480: 480 g/L atrazine formulated as a liquid.

Atrazine 90WG: 90% atrazine formulated as a dispersible granule.

Container sizes - various.

Crops and Staging:

May be applied to corn (silage, field, sweet) using the following applications:

Pre-plant incorporated (PPI).

Pre-emergent surface (after planting but before emergence of weeds and crop) - are recommended only on irrigated fields. Inconsistent weed control will occur if 0.5 inches (1.25 cm) of water/precipitation does not occur within 7 days of application.

Post-emergence - 1 to 6 leaf stage and when corn is less than 12 inches (30 cm) tall.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

For pre-plant incorporated, pre-emergent and post-emergent (when weeds are less than 4 inches or 10 cm tall) of the following weeds::

Common purslane
Lady's-thumb
Lamb's-quarters
Ragweed
Redroot pigweed
Smartweed
Volunteer clover
Wild buckwheat
Wild mustard
Wild oats
Wormseed mustard

Rates:

	ATRAZINE 90 WG	ATRAZINE 480, AATREX LIQUID
Preplant incorporated; Pre-emergence (irrigated corn only)	0.45 to 0.69 kg/acre	0.85 to 1.25 L/acre
Post-emergence	0.45 to 0.69* kg/acre	0.85 to 1.25* L/acre

* Add 1.11 to 2.23 L/acre of oil concentrate or 6.88 L/acre crop oil with post-emergence applications of these products.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Crop injury may occur when Atrazine and oil is applied post-emergence during cold weather.

Use the low rate on crops grown on sandy soils, and where weed infestations are light.

Cost (2004 suggested retail price):

Aatrex Liquid - $7.26 per L
Atrazine 480 - $7.30 per L
Atrazine 90WG - $12.47 per kg

Application Information:

Water Volume: Minimum 61 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan with 50-mesh or coarser screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Post-emergent applications made during periods of cold weather may cause crop lightening. Hot, dry weather preceding post-emergent applications may result in reduced weed control. Atrazine will move with soil if eroded.

Tank Mixes:

Herbicides:

Pre Plant Incorporated - Eradicane (not sweet corn)
Post-Emergent* - Banvel II (not sweet corn), Pardner.

* Do not use oils or adjuvants with post-emergent tank mixes.

Fertilizers: For pre-emergence applications, nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for some formulations of atrazine. Aatrex Liquid may be impregnated onto dry granular fertilizers. Do not impregnate onto nitrate or super-phosphate.

Do not apply atrazine with nitrogen fertilizer after corn has emerged, as crop injury will occur.

Insecticides: None registered.

Note: The above mixes are those listed on the atrazine labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. See the general guidelines for mixing pesticides for more information.

Restrictions:

Rainfall: Within 2 hours of postemergence applications may result in reduced weed control.

Grazing: Do not graze or cut for feed before ear emergence.

Preharvest: Leave at least 45 days from application to harvest.

Recropping: All crops, except corn and triazine-tolerant canola, may be affected the year following the use of atrazine. Flax, peas and fababeans have some tolerance to atrazine residues and are usually not affected by rates of up to 0.9 L/acre (500 g/L formulations) or 0.5 kg/acre (90 percent atrazine formulations) applied the previous year. Other more sensitive crops may be affected 2 or more growing seasons after application.

Aerial Application: Do not apply by air.

Storage: Do not freeze Aatrex Liquid or Atrazine 480.

Environment: Do not mix or load within 30 m, or apply within 10 m, of any wells, lakes, streams, ponds, dugouts or sinkholes.

Tank Cleaning:

When finished spraying atrazine, run clean water through the tank, pump and lines. Drain and refill with 1 L of 3% household ammonia solution per 100 L water. Circulate the solution through lines and nozzles. Let solution stand for several hours. Scrub inside surfaces but do not enter tank. Flush sprayer system with water.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 14 for additional information.

Hazard Rating:

◈ Warning Poison (Atrazine 90WG)

NOTE: While Atrazine 90WG is the only product with a hazard rating, other formulations of atrazine could be expected to have similar hazard ratings if rated.

Attain

Herbicide Group – 4

(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

The Attain package has 2 components:

Attain A (PCP #24834): 180 g/L fluroxypyr

Attain B (PCP #24833): 564 g/L 2,4-D LV ester.

Both formulated as emulsifiable concentrates.

Container sizes - Attain A - 9.6 L, Attain B - 2 x 8 L.

Crops and Staging:

Spring wheat (including durum) barley and tall fescue (seedling and established for seed production only)* at the 4 leaf to flag leaf stage.

* *Since this product is registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance.* ***Users of this product on tall fescue do so at their own risk***

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 2 to 4 leaf stage, unless specified:

Bluebur
Burdock
Cleavers (1 to 4 whorls)
Clover (sweet)
Cocklebur
Field horsetail*
Flixweed
Goat's beard
Hoary cress*
Kochia
Lamb's-quarters
Mustards
  (except dog and tansy)
Plantain
Prickly lettuce
Ragweed
Round leaved mallow
  (1 to 6 leaf)
Shepherd's-purse
Stinkweed
Stork's-bill
  (1 to 8 leaf)
Sunflower (annual)
Vetch
Volunteer canola
Volunteer flax (1-12cm)
Wild radish
Wild buckwheat (1 - 4 leaf)
Wild mustard

Weeds controlled only when growing rapidly (control may be reduced when weed infestations are heavy):

Blue lettuce*
Dandelion**
Docks
Field bindweed*
Field peppergrass
Lady's-thumb
Leafy spurge*
Mustard (dog and tansy)
Oak-leaved goosefoot
Redroot pigweed
Russian thistle
Smartweed
Tartary buckwheat

Weeds suppressed:

Annual sow-thistle
Canada thistle*
Common chickweed up to 3 inches (8 cm) in height
Hemp-nettle (2 to 6 leaf stage)
Perennial sow thistle*

* Top growth only
** Spring rosettes only.

Cost (2004 suggested retail price):

$9.13 per acre.

Rate:

Apply Attain A at 0.24 L/acre and Attain B at 0.40 L/acre. One case of Attain treats 40 acres. Make only one application per year.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 45 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa)

Nozzles: Use nozzles that deliver a uniform coverage while minimizing fine droplets prone to drift.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Attain activity is influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought, heat or cold stress) or if extremely heavy infestations exist.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:
Achieve Liquid (0.2 L/acre) plus Turbocharge adjuvant.*
Assert (0.53 to 0.65 L/acre)

In spring wheat (including durum) only:
Horizon (95 mL/acre) plus Score adjuvant.*
Puma^{120} Super (0.16 to 0.31 L/acre)
Everest at 17.4 g/acre plus recommended surfactant

Fertilizer: None registered.

Insecticides: None registered.

Fungicides: None registered.

*Temporary crop injury (Achieve only) or reduced wild oat control may occur with these tank mixes.

Note: The above mixes are those listed on the Attain label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not permit lactating dairy animals to graze fields within 7 days of application. Do not harvest forage or cut hay within 30 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Preharvest Interval: Leave 60 days between application and harvest.

Recropping: Wheat, barley, oats, rye, forage grasses, flax, canola, mustard, lentils and peas may be grown the year after an Attain application. There are no recropping restrictions the second year after application.

Aerial Application: Do not apply by air.

Storage: May be frozen. If frozen, bring to room temperature and agitate before use. This product is combustible. Do not store near heat or open flame.

Environment: Avoid drift onto non-target areas. Leave a buffer of 15 meters from water bodies, wetland areas and plants that may be injured by Attain.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Danger Poison.

Warning Eye Irritant.

Caution Skin Irritant.

Avadex BW/ Avadex MicroActiv

Herbicide Group – 8
(Refer to page 27)

Company:

Gowan Company*

Formulation:

400 g/L triallate formulated as an emulsifiable concentrate. Container sizes - 22.7 L, 115 L.

10% triallate formulated as a granular. Container size - 22.7 kg, 454 kg.

***Note: The production and marketing rights to Avadex products have recently been purchased by Gowan Co. While limited quantities still remain in the supply system, Gowan expects to have new product available for the fall of 2005. Remaining material originally produced by Monsanto will remain labeled as such, but will be supported in future by Gowan Co.**

Crops, Rates and Application Timing:

Avadex Liquid Rates – Spring Treatment

CROP	APPLICATION TIMING	RATE (L/acre)		ACRES TREATED PER 22.7 L CONTAINER	
		Organic Matter		Organic Matter	
		4% or less	Greater than 4%	4% or less	Greater than 4%
Spring and durum wheat	Before Seeding* After Seeding	1.21 1.42	1.42 1.72	18.8 16.0	16.0 13.2
Barley	Before and After Seeding	1.42	1.72	16.0	13.2
Canola, flax, mustard	Before Seeding	1.72	2.23	13.2	10.2
Peas (dry)	Before Seeding	1.72	1.72	13.2	13.2

* Do not apply this product before seeding wheat in soils with 4 percent or less organic matter (brown, dark brown or grey wooded soils) where discers are to be used for seeding. If an air seeder is to be used, it must be equipped with a depth control device to ensure accurate seed placement, otherwise crop injury may occur.

Avadex Granular Rates – Fall Treatment

CROP	RATE (KG/ACRE)			ACRES TREATED PER 22.7 KG CONTAINER		
	Organic Matter			Organic Matter		
	Less than 2%*	2-4%	Greater than 4%	Less than 2%*	2-4%	Greater than 4%
Spring and durum wheat	4.45	5.67	6.88	5.1	4.0	3.3
Barley, canaryseed	4.45	5.67	6.88	5.1	4.0	3.3
Canola, flax, mustard	5.67	6.88	8.90	4.0	3.3	2.6

* Fall treatments conducted under minimum tillage are not recommended on soils with less than 2 percent organic matter.

Avadex Granular Rates – Spring Treatment

CROP	APPLICATION TIMING**	RATE (KG/ACRE)		ACRES TREATED PER 22.7 KG CONTAINER	
		Organic Matter		Organic Matter	
		4% or less*	Greater than 4%	4% or less*	Greater than 4%
Spring and durum wheat	Before seeding*** After seeding	4.45 5.67	5.67 6.88	5.1 4.0	4.0 3.3
Barley, canaryseed	Before and after seeding (barley only)	5.67	6.88	4.0	3.3
Canola, flax, mustard	Before seeding	6.88	8.90	3.3	2.6

* Minimum tillage treatments must be applied to fields with at least 2 percent organic matter.

** Minimum tillage treatments must be applied 10 to 14 days before seeding or incorporating. For minimum tillage treatments on spring and durum wheat, apply 5.67 kg/acre on soils with 4% organic matter or less and 6.88 kg/acre on soils with greater than 4 percent organic matter.

*** Do not apply this product before seeding wheat in soils with 4% or less organic matter (brown, dark brown or grey wooded soils) where discers are to be used for seeding. If an air seeder is to be used, it must be equipped with a depth control device to ensure accurate seed placement, otherwise crop injury may occur.

Seedling Forage Legumes (under-seeded only):

Apply recommended rates for the companion crop.

Alfalfa
Bird's-foot trefoil
Clover (alsike, red, sweet)

Weeds and Staging:

For pre-emergent control of Wild oats.

Cost (2004 suggested retail price):

Liquid: $9.47 to $17.45 per acre

Granular: $12.20 to $24.40 per acre.

Application Information:

Water Volume (Liquid formulations only):
45 L per acre.

Pressure: 30 psi (200 kPa), liquid formulation only.

Nozzles: Flat fan, liquid formulation only.

General Information: The liquid formulation must be incorporated into soil that is free of lumps or trash. The liquid formulation is recommended for spring use because soils are left in an erosion prone state if the liquid is fall-applied. The granular formulation may be incorporated into trashy soil and is best suited for fall use.

Fall Applications (Conventional Tillage): Apply Avadex granules to fields that are in good working condition, without excessive trash. Heavy trash or lumpy, wet fields may require tillage prior to application. Avadex must be applied after October 1 but before soil freeze-up. Application before October 1 may result in reduced weed control. Only one incorporation is required in the fall. The second incorporation may be done in the fall (before soil freeze-up) or in the spring.

Fall Application (Minimum Tillage): Applications of Avadex granules should be made to standing stubble, chemical fallow, or summerfallow fields that are not prone to erosion. Do not apply to smooth, hard packed soils that may allow granules to drift. If excessive crop residue exists at the time of application, harrowing should be conducted to ensure the granules are in good contact with the soil. Apply when the soil begins to cool (less than 4°C) and within 3 weeks of soil freeze-up. Incorporation can be performed in the spring before seeding or as part of the seeding operation.

Spring Application (Conventional Tillage): Apply Avadex (liquid or granules) to fields that are in good working condition, without excessive trash. Heavy trash or lumpy, wet fields may require tillage prior to application. Liquid formulations should be applied to fields with 30 percent or less trash cover. Avadex may be applied before or after seeding of wheat, barley, or canaryseed and before seeding of canola, flax, mustard or peas (liquid only). If wheat is being seeded into soils with an organic matter content of less than 4 percent, Avadex should be applied after seeding.

Spring Application (Minimum Tillage): Avadex granules should be applied in spring and when the soil temperature is 4°C or less. Apply granules 10 to 14 days before incorporation. Do not apply more than 4 weeks before seeding is intended.

Incorporation:

Conventional Tillage: Avadex applications require 2 incorporations, with the second incorporation at right angles to the first. Using a seeder that provides soil disturbance equivalent to a cultivator may replace 1 incorporation. The first incorporation of the granular formulation should be completed within 48 hours of application and the second incorporation should be delayed an additional 48 hours or more. The first incorporation of the liquid formulation should be completed as soon as possible after spraying, while the second incorporation may be done any time prior to crop emergence.

Incorporate to a depth of 2 inches (5 cm) by setting disc or cultivator implements to cut 3 inches (7.5 cm) into the soil. Mixing the product to greater depths will dilute the herbicide, decrease wild oat control, and may cause injury to cereals. Ensure that cereals are seeded below the treated layer (2 to 3 inches or 5 to 7.5 cm). Incorporations performed after seeding should be conducted with harrows or other suitable tillage equipment adjusted so as not to disturb the seed. Harrowing does not provide effective incorporation if compact soil prevents penetration of harrow teeth, if trash accumulates in the harrow sections, or if the harrows bounce.

Minimum Tillage: Incorporation of Avadex granules in minimum tillage systems is achieved with one high disturbance incorporation, which can be conducted prior to seeding, or as part of the seeding pass. A high disturbance system is one that disturbs the soil enough so that emerged weeds are controlled by the operation (example - air seeder with cultivator shovels). Harrowing after the incorporation operation is recommended for best results.

For optimum results in minimum tillage systems, incorporate when wild oat growth is noticeable in the field, as this will ensure that the soil is warm enough for activation of Avadex.

Under excessively warm or wet conditions between application and crop emergence, control may be reduced. For best results on heavy wild oat infestations, use the conventional tillage guidelines for incorporation.

Summer Fallow: Incorporation can be done by a disc followed by harrowing at right angles, a vibrashank cultivator followed by harrowing at right angles, or double harrowing. The second operation can be delayed until spring. If summerfallow must be ridged to prevent soil erosion the granular formulation should not be used in the fall. Note that fall minimum tillage applied granules do not require incorporation in the fall. If soils must be ridged following application of the liquid formulation, ridging depth should be kept to a minimum as deep ridging may reduce wild oat control and increase crop injury.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Reduced control may result if prolonged cool conditions or dry soil conditions prevail at the time weeds are emerging. If conditions are dry or wild oats germinate from below the treated zone, the weeds may emerge, but will usually be controlled. Thinning of wheat can occur under conditions of heavy rainfall or if cold soil conditions persist as the crop emerges.

Tank Mixes:

Herbicides: Avadex liquid may be tank mixed with liquid formulations of trifluralin for control of wild oats, green and yellow foxtail in wheat and barley. Apply after seeding but prior to crop emergence. Consult the recommendations for trifluralin for rates in different soil types.

Fertilizer: Avadex liquid alone, or tank mixed with liquid formulations of trifluralin, may be tank mixed with liquid fertilizer. Compatibility of the herbicide and liquid fertilizer should be checked. Follow the instructions on the herbicide label prior to adding the herbicide to the spray tank.

Avadex liquid may be sprayed on dry urea fertilizer. A minimum of 150 kg/ha (60 kg/acre) of dry urea fertilizer must be used. Only commercial blending is recommended.

Insecticides: None registered.

Note: The above mixes are those listed on the Avadex labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: At least 0.5 inches (1.5 cm) within 2 weeks of application is required for activation.

Grazing: Do not graze the treated crop or use as hay or feed prior to crop maturity or in year of treatment.

Recropping: Do not seed tame oats the year after treatment.

Aerial Application: May be applied by air with attachments designed for applying low volumes of granules.

Storage: Do not freeze liquid formulations. Store granular formulations in a cool, dry place.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison (liquid formulation)

Skin and Eye Irritant (granular formulation)

Avenge 200 C*

Herbicide Group – 8

(Refer to page 27)

***Note Avenge will no longer be available following the 2005 growing season, as production of this product has been discontinued.**

Company:

BASF Canada (PCP#12853)

Formulation:

200 g/L difenzoquat formulated as a solution.
Container size - 20 L.

Crops and Staging:

Apply up the 6 leaf stage of cereal and grass crops:
Barley
Canaryseed
Fall rye (Cougar, Frontier, Kodiak, Puma, Rymin only)
Triticale (Carman and Welsh)
Wheat (AC Barrie, Biggar, Bluesky, CDC Makwa, CDC Teal, Columbus, Conway, Cutler, Fielder, Genesis, Glenlea, Katepwa, Lancer, Leader, Neepawa, Oslo, Pasqua, Selkirk and Wildcat wheat varieties).
Winter wheat (Norstar only)

Forages (only when underseeded to wheat and barley varieties listed above):
Alfalfa
Bird's-foot trefoil
Bromegrass
Creeping red fescue
Crested wheatgrass
Kentucky bluegrass
Meadow fescue
Orchardgrass
Red clover
Reed canarygrass
Russian wild ryegrass
Sweet clover
Timothy

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Wild oats at the 3 to 5 leaf stage.

Cost (2004 suggested retail price):

$18.56 to $22.47 per acre.

Rate:

Apply Avenge 200 C at 1.42 to 1.72 L/acre. Use the higher rate if wild oat population exceeds 200 wild oats per square metre.

One 20 L container treats 14.1 to 11.6 acres.

Application Information:

Water Volume:
Ground application: 40 L per acre.
Aerial application: 8 to 20 L per acre

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan directed 45° forward.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop injury is worse on cold or hot, humid days than when temperatures are moderate. Best weed control is attained when temperatures are 20 to 30°C, particularly when these temperatures follow application. Do not spray when crop or weeds are wet with heavy dew or rain. Do not apply when crop is under stress from drought, excessive moisture, or heat.

Tank Mixes:

Herbicides:

In wheat and barley:
2,4-D ester (up to 0.45 L/acre of 500 g/L formulation). Do not use amine formulations.
2,4-DB (see 2,4-DB section for rates). May be used on cereals underseeded to forages.
Ally (2 to 3 g/acre)*
Ally + MCPA ester*
Buctril M (0.4 L/acre)
Curtail M (0.8 L/acre)
Dichlorprop + 2,4-D (0.71 L/acre)
MCPA ester (up to 0.45 L/acre of 500 g/L formulation). Do not use amine formulations.
Pardner (0.405 to 0.485 L/acre)
Pardner + MCPA ester
Refine Extra (8 g/acre)
Thumper (0.4 L/acre)

In canaryseed:
MCPA ester (0.38 to 0.45 L/acre of 500 g/L formulation)
Pardner (0.405 L/acre)
*No additional adjuvant required.

Insecticides: None registered

Fertilizers: None registered.

Fungicides: None registered.

Allow a 5-day interval between the application of Avenge and other pesticides.

Note: The above mixes are those listed on the Avenge label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours of application will reduce control.

Grazing: Do not graze or feed treated cereal crop for 8 weeks after treatment. Do not graze or feed treated forages in the year of treatment. Straw from treated fields can be fed to livestock 8 weeks after application.

Preharvest Interval: Do not apply beyond the recommended crop stage.

Recropping: No restrictions the year after treatment.

Aerial Application: May be applied by air.

Storage: May be frozen.

Environment: Do not contaminate water bodies (sloughs, dugouts, lakes, streams, ponds, etc.).

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

Banvel II

Herbicide Group – 4
(Refer to page 27)

See dicamba on page 88.

Basagran/Basagran Forté

Herbicide Group – 6
(Refer to page 27)

Company:

BASF Canada

Basagran (PCP#12221)
Basagran Forté (PCP#22006)

Formulation:

480 g/L bentazon formulated as a solution. Basagran Forté has a built-in adjuvant.

Container size - 2 x 9 L jugs.

Crops and Staging:

Basagran and Basagran Forté:

CROP	STAGE
Soybeans	No restrictions
Dry beans	After the first trifoliate leaf
Corn	No restrictions
Peas*	After 3 pairs of leaves
Fababeans	At least 4 inches (10 cm) tall
Flax	After 2 inches (5 cm) in height

Basagran Forté only:

CROP	STAGE
Forage millet and Forage sorghum*	3 to 6 leaf prior to canopy closure

Basagran only:

CROP	STAGE
Spring wheat (excluding durum)**	No restrictions (limited to the 4 leaf to flag leaf by 2,4-D staging)
Solin	After 2 inches (5 cm) in height
Forage grasses for seed production*: Bromegrass, creeping red fescue, crested wheatgrass, meadow foxtail, orchardgrass, timothy.	1 to 7 leaf stage
Forage legumes (seedlings) for seed production*: Alfalfa, alsike clover, red clover, sainfoin.	After the third trifoliate leaf
Alfalfa for seed production.	Established prior to flowering
Sweet clover for seed production.	3 to 10 inches (7.5 to 25 cm) high

* One application per season.
** Basagran only at 0.4 L/acre. Must be tank mixed with 2,4-D (no adjuvant required).

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Cost (2004 suggested retail price):

Basagran: $29.43 per L not including adjuvant.

Basagran Forté: $ 25.18 per L adjuvant included in formulation.

Weeds, Rates and Staging:

Add Assist or XA oil concentrate at 0.41 to 0.81 L/acre to Basagran only. Basagran Forté does not require the addition of Assist or XA oil concentrate. If hot, humid conditions prevail (above 28°C and 80% relative humidity), use only the low rate of Assist or XA oil concentrate. Citowett Plus may be used on peas at 0.25 L per 100 L spray mixture.

Apply the rate listed when weeds in the table are within the recommended height:

ANNUAL WEEDS	0.71 L/acre		0.91 L/acre	
	Inches	**Maximum Leaf Stage**	**Inches**	**Maximum Leaf Stage**
Buttercup			2 to 4	6*
Cleavers			1 to 3 whorl stage	
Cocklebur	3 to 7	6*	7 to 12	10*
Common chickweed			1 to 3 weeks after emergence	
Common groundsel			2 to 4	
Common ragweed			1 to 2	6
Corn spurry			1 to 4	
Giant ragweed			2 to 6	4
Hairy galinsoga			2 to 3	6*
Hairy nightshade			0.2 to 0.8	6
Lady's-thumb (smartweed)	1 to 3	6*	3 to 8	10
Lamb's-quarters			0.5 to 1.0	8
Purslane			1 to 2	6
Redroot pigweed (suppression only)			0.5 to 1.5	4
Russian thistle (suppression only*)			1 to 3	4*
Shepherd's-purse	Rosette to 4	6*	4 to 10	6
Stinkweed	Rosette to 2	6*	2 to 6	6
Stork's-bill			1.5 to 4	2 to 6 leaf stage
Flower of an hour	1 to 2	6*	2 to 4	10*
Volunteer canola	0.75 to 6	8	0.75 to 6	8
Wild mustard	1 to 5		5 to 10	10
Wild radish			1 to 2	6
PERENNIAL WEEDS	Repeat application 7 to 15 days after first application (if necessary)			
Canada thistle	6 to 8			
Field bindweed	1 to 2.5			
Yellow nutsedge	6 to 8			

*Applies to Basagran Forté only.

Basagran may be applied in wheat at 0.4 L/acre when tank mixed with 2,4-D amine or ester (500 g/L) at 0.3 to 0.4 L/acre to control Lady's-thumb, redroot pigweed and daisy fleabane in addition to the weeds controlled by 2,4-D. No adjuvant is required for this mix.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 to 160 L per acre. A minimum of 80 L per acre. is recommended for optimum control.*

Pressure: 40 to 60 psi (275 to 425 kPa)*

Nozzles: Flat fan nozzles capable of delivering high water volumes. Direct nozzles 45° forward.

* Higher water volumes and pressures should be used when the weeds are at the upper end of their recommended treatment stage.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Poor results will occur if temperatures are cool. Optimum results are achieved when applied at daytime temperatures between 20 and 28°C. Applications at temperatures greater than 28°C may result in crop injury.

Tank Mixes:

Herbicides:

In soybeans:

Pinnacle at 2.2 to 3.2 g/acre.

***In dry beans** (in the Red River Valley of Manitoba only):*

Basagran only at 0.71 L/acre can be tank mixed with Reflex at 0.23 L/acre plus Agral 90.

***In spring wheat** (not including durum):*

Basagran only at 0.4 L/acre can be tank mixed with 2,4-D amine or ester (500 g/L) at 0.3 to 0.4 L/acre. This tank mix DOES NOT need any adjuvant.

Fertilizers: Use of fertilizer mixes is not recommended for the western Canadian environment.

Insecticides: None registered.

Fungicides: None registered.

Allow 4 days between application of Basagran and other herbicides, fertilizers or insecticides.

Note: The above mixes are those listed on the Basagran/Basagran Forté labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 to 8 hours will reduce control.

Grazing: Allow 30 days between treatment with Basagran Forté and harvest of forage sorghum and millet for hay. Otherwise do not graze treated crops or cut for feed prior to crop maturity.

Preharvest Interval: 50 days for Basagran + 2,4-D in wheat, 84 days for Basagran + Reflex in Dry beans in Manitoba. Other uses are restricted only by appropriate staging.

Recropping: No restrictions the year after application.

Aerial Application: May be applied by air for weed control in dry beans or soybeans only. Use 23 to 45 L/acre water volume. Assist or XA oil concentrate at 0.05 to 0.1 L/acre must be added. Do not use Assist or XA oil concentrate in excess of 0.1 L/acre as substantial crop injury could occur. Do not apply fertilizer mixes in soybean or 2,4-D tank mix in wheat by air. Crop canopy should not cover the weeds.

Storage: May be frozen.

Environment: Avoid drift onto non-target areas. Do not contaminate domestic, irrigation or natural water sources.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison – Basagran

Warning Poison – Basagran Forté

Bromoxynil

Herbicide Groups – 6
(Refer to page 27)

Company:

Bayer CropScience (Pardner) - PCP#18001
Nufarm (Koril 235) - PCP#25341

Formulation:

Pardner: 280 g/L bromoxynil formulated as an emulsifiable concentrate.
Container size - two 8 L containers per case.

Koril 235: 235 g/L bromoxynil formulated as an emulsifiable concentrate.
Container size – two 9.71 L containers per case.

Crops and Staging:

Registered uses for both Pardner and Koril 235:

CROP	STAGE
Barley, oats, triticale, wheat (spring and durum)	2 leaf stage to early flag
Winter wheat	2 to 4 leaf stage (fall application) First growth to early flag leaf (spring application)
Corn (field or sweet)	4 to 8 leaf
Corn (field or sweet) with drop pipes	Beyond 8 leaf

Registered uses for Pardner only:

CROP	STAGE
Canaryseed	3 to 5 leaf
Seedling alfalfa	2 to 6 trifoliate leaf stage
Established alfalfa (seed production only)	In spring, before the crop begins to shield weeds. Apply no more than twice in one growing season.
Fall rye	In spring only, from first growth to early flag
Flax (including Solin)	2 to 4 inches (5 to 10 cm)
Forage millet and sorghum	4 leaf to 8 inches (20 cm)
Seedling grasses: Bromegrass, Fescue (creeping red, meadow), Orchardgrass, Reed canary grass, Russian wild ryegrass, Timothy, Wheatgrass (crested, intermediate, slender, tall)	2 to 4 leaf
Summerfallow	Apply according to weed stage.

Weeds and Staging:

Weeds controlled at the 1 to 4 leaf stage:

American nightshade
Bluebur
Cocklebur
Common ragweed
Cow cockle*
Green smartweed
Kochia**
Lady's-thumb
Pale smartweed
Pigweed*
Stinkweed*
Russian thistle**
Wild mustard*

* Use high rate.
** Apply before plants are 2 inches high.

Weeds controlled at the 1 to 8 leaf stage:

Common groundsel
Lamb's-quarters
Tame buckwheat
Tartary buckwheat
Wild buckwheat

Cost (2004 suggested retail price):

Pardner: $8.73 to $10.60 per acre.
Koril: $8.73 to $10.20 per acre.

Rate:

Pardner: 0.405 to 0.486 L/acre (one 8 L container treats 20 to 16.5 acres).

Canaryseed, flax, seedling alfalfa, forage millet and sorghum - 0.405 L/acre (one 8 L container treats 20 acres).

Koril 235: 0.486 to 0.57 L/acre (one 9.71 L container treats 20 to 17 acres).

Use the higher rates if weeds are at the more advanced stages of the recommended treatment stages.

Application Information:

Water Volume:
Corn, Millet & Sorghum: 80 to 120 L per acre.
Seedling grasses: 60 L per acre.
Other crops: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Avoid spraying if temperatures are greater than 25°C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (greater than 27°C) and humid weather.

Tank Mixes:

Pardner Herbicide Tank Mixes :

CROP	PARDNER TANK MIXES
Spring wheat	2,4-D, Avenge, Avenge + MCPA (ester only), MCPA, Horizon, Hoe-Grass 284
Winter wheat	2,4-D, MCPA
Barley	2,4-D, Avenge, Hoe-Grass 284, Avenge + MCPA (ester only), MCPA
Oats	MCPA
Fall rye, canaryseed, seedling forage grasses	MCPA*
Corn	Accent§ + surfactant, Atrazine**, Banvel II (field corn only)**

* The ester formulations are preferred but other formulations can be used.

** Do not add oil or surfactant to this mix. Do not use atrazine formulations that contain oil.

§ *Since the use of this tank mix on corn is registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance.* ***Users of this tank mix on corn do so at their own risk.***

Koril 235 Herbicide Tank mixes:
Wheat, Barley and Oats: MCPA
Wheat and Barley only: 2,4-D
Corn only: atrazine*

* Do not add oil or surfactant to this mix. Do not use atrazine formulations that contain oil.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the bromoxynil labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour of application will reduce control.

Grazing: Do not graze treated wheat, barley, oats or seedling alfalfa crops or cut for feed within 30 days of application.

Do not graze other treated crops or cut for hay prior to crop maturity.

Recropping: No restrictions.

Aerial Application: Registered for aerial application on wheat and barley. The use of low water volumes, 9 to 18 L per acre may result in less effective weed control than seen with ground application.

Storage: May be stored at freezing temperatures.

Will return to original state by warming to room temperature (20 to 22°C) and agitating thoroughly.

Environment: Avoid drift onto sensitive plants and contamination of water or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Warning Poison

Danger Eye and Skin Irritant (Koril 235)

Bromoxynil+MCPA ester

Herbicide Groups – 4, 6
(Refer to page 27)

Company:

Bayer CropScience (Buctril M) - PCP#18022
Nufarm (Mextrol 400 M) - PCP#25342
United Agri Products (Badge) - PCP#16164

Formulation:

Buctril M: 280 g/L bromoxynil present as heptyl and octyl esters and 280 g/L of MCPA ester formulated as an emulsifiable concentrate.
Container size - 8 L.

Mextrol 400 M: 200 g/L bromoxynil present as octyl ester and 200 g/L of MCPA ester formulated as an emulsifiable concentrate.
Container size – 11.36 L.

Badge: 225 g/L bromoxynil present as octyl ester and 225 g/L of MCPA ester formulated as an emulsifiable concentrate.
Container size – 10 L.

Crops and Staging:

Field Crops:

All Products:

Barley, oats, spring wheat (including durum) - 2 leaf to early flag.
Winter wheat - 2 to 4 leaf stage in the fall or after growth resumes in the spring, but prior to early flag leaf.
Fall rye - apply from the time growth commences in spring to early flag leaf.
Canaryseed - 3 to 5 leaf stage.
Flax, Solin - 2 to 4 inches (5 to 10 cm) tall.
Corn - 4 to 6 leaf stage.

Seedling forage grasses:
2 to 4 leaf stage.

All Products:

Bromegrass
Fescue (creeping red, meadow)
Reed canarygrass
Russian wild ryegrass
Timothy
Wheatgrass (crested, intermediate, slender, tall)

Buctril M and Badge only:

Fescue (tall)
Meadow bromegrass
Meadow foxtail
Wheatgrass (streambank)

Buctril M only:

Orchard grass

Established Forage Grasses (Buctril M and Badge only):
Timothy for seed (or hay Buctril M only) - apply prior to the emergence of the flag leaf.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds up to 4 leaf stage:

American nightshade
Bluebur
Ball mustard
Cocklebur
Cow cockle
Flixweed
Green smartweed
Kochia**
Lady's-thumb
Night-flowering catchfly
Pale smartweed
Redroot pigweed*
Russian thistle**
Scentless chamomile***
Shepherd's-purse
Volunteer canola
Volunteer sunflower

* May not be controlled in flax.
** Control before plants are 2 inches tall.
*** Spring seedlings only.

Weed up to 6 leaf stage:
Wild tomato (Buctril M & Badge only)

Weeds up to 8 leaf stage:

Common groundsel
Common ragweed
Lamb's-quarters
Stinkweed
Tame buckwheat
Tartary buckwheat
Wild buckwheat
Wild mustard
Wormseed mustard

Weeds where top growth is controlled:

Canada thistle
Perennial sow-thistle

Cost (2004 suggested retail price):

Buctril M: $6.95 per acre.
Mextrol 400M: $6.68 per acre.
Badge: $6.41 per acre.

Rate:

Buctril M: 0.4 L per acre. One 8 L jug treats 20 acres.
Mextrol 400 M: 0.57 L per acre. One 11.36 L jug treats 20 acres.
Badge: 0.5 L per acre. One 10 L jug treats 20 acres.

Application Information:

Water Volume:
Corn: 80 to 120 L per acre.
Flax, Solin: 20 to 40 L per acre.
Cereals: 20 to 40 L per acre.
Seedling forage grasses: 60 L per acre.

Established timothy: 60 L per acre.
Aerial: 8 to 20 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan nozzles.

Refer to specific labels for recommended water volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Best weed control when humidity is high at the time of spraying and for the following day or two. Prolonged cool conditions may result in reduced weed control. Spraying during early morning may increase the risk of flax injury.

Avoid spraying in temperatures greater than 25°C.

Do not apply to flax, canaryseed or corn if daytime temperatures exceed 27°C within 48 hours before or after application.

Tank Mixes:

Herbicide Tank Mix Table:

Products listed below are at label rates for each crop. See labels for details.

CROP & TANK MIXES	Badge	Buctril M	Mextrol 400M
Flax (✓✓ = also registered for use on Solin):			
Fusion		✓✓	✓
Poast Ultra + Merge adjuvant	✓✓		✓✓
Select + Amigo adjuvant	✓✓	✓✓	✓✓
Spring wheat (including durum) and Barley:			
Ally	✓	✓	✓
Avenge	✓	✓	✓
MCPA (amine, ester & K)	✓	✓	✓
Puma(2) Super		✓	
Refine Extra (2.7 g/acre)	✓	✓	
Refine Extra + Puma(2) Super (rates above)		✓	
Spring wheat (including durum) only:			
Everest	✓		
Horizon	✓	✓	✓
Winter Wheat:			
Refine Extra (2.7 g/acre)	✓	✓	

CROP & TANK MIXES	Badge	Buctril M	Mextrol 400M
Oats:			
MCPA (amine, ester & K)	✓	✓	✓
Corn:			
Atrazine	✓	✓	✓

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the bromoxynil + MCPA ester labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rainfall within 1 hour of application may reduce weed control.

Grazing: Do not graze treated grain or established timothy crops or cut for feed within 30 days of application. Do not graze meadow foxtail in the year of treatment. Do not graze other treated forage grasses within 56 days of treatment.

Preharvest Interval: Do not harvest Flax or Solin within 60 days of application.

Recropping: No recropping restrictions the year after treatment.

Aerial Application: May be applied by air to wheat, barley, and oats only. Must be applied in 8 to 18 L per acre water volume. Use higher water volume when the majority of weeds are cow cockle, smartweed, hemp-nettle, pigweed, and Canada thistle.

Storage: May be frozen.

Environment: Apply when wind is blowing away from sensitive plants and water or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Warning Poison (Badge and Buctril M)

Danger Poison (Mextrol 400M)

Warning Skin Irritant (Mextrol 400M)

Buctril M/Mextrol 400 M

See Bromoxynil + MCPA section on page 81

Casoron

Herbicide Group – 20
(Refer to page 27)

Company:

Crompton Co. (PCP#12533)
(distributed by United Agri Products)

Formulation:

4% dichlobenil formulated as a granular.
Container size - 3 kg, 15 kg.

Crops:

Poplar plantations

Shelterbelts consisting of the following species:

- Ash
- Barberry
- Birch (cutleaf-weeping)
- Boxwood
- Caragana
- Cedar (White, Eastern Red)
- Crabapple
- Elm
- Euonymus (Burning bush)
- Forsythia
- Honeysuckle
- Juniper
- Lilac
- Linden
- Locust
- Maple
- Mock orange
- Poplar
- Rose
- Spirea
- Willow

NOTE: Do not apply to shelterbelts with mugo pine, firs, hemlock, holly, spruce or other shallow rooted species or injury may result. Do not apply in or around greenhouses.

Weeds and Staging:

Apply in early spring or late fall prior to annual weed emergence, or after cultivation has removed existing weeds.

- Annual blugrass
- Artemisia (absinthe, * wormwood, sage)
- Bindweed*
- Canada thistle*
- Chickweed
- Dandelion*
- Foxtail (green and yellow)
- Groundsel
- Horsetail
- Knotweed
- Kochia
- Lamb's-quarters
- Loosestrife
- Mustard
- Nutsedge*
- Pigweed
- Plantain
- Purslane
- Quack grass*
- Sheep sorel*
- Shepherd's-purse
- Smartweed
- Sow-thistle
- Spurge
- Vetch*
- Wild buckwheat*

*Controlled with fall applications at the higher rates.

Cost (2004 suggested retail price):

$8.78 per kg.

Rates:

45 to 70 kg/acre. At the low rate, a 15 kg bag will treat a 4 yd by 407 yd (4 m by 340 m) strip of shelterbelt. At the high rate, a 15 kg bag will treat a 4 yd by 256 yd (4 m by 214 m) strip of shelterbelt. If application is followed by 0.5 to 1.0 inches (1.3 to 2.5 cm) of irrigation, the lower rates are recommended.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply during periods of high soil temperatures (more than 15°C).

Tank Mixes:

None registered.

Restrictions:

Grazing: Do not graze in treated area.

Storage: Store in a cool, dry place. May be frozen.

Equipment Cleaning:

Refer to page 14.

Hazard Rating:

No specific rating. Keep out of reach of children.

Harmful if swallowed.

Avoid skin or eye contact.

Clethodim

Herbicide Group – 1

(Refer to page 27)

Company:

Arvesta (Select) - PCP#22625

Bayer CropScience (Centurion) - PCP#27598

Formulation:

Select: 240 g/L of clethodim formulated as an emulsifiable concentrate. Container size - 3 L.

Centurion: 240 g/L of clethodim formulated as an emulsifiable concentrate. Container size - 3 L.

Crops, Rates and Staging:

Crops are tolerant at all growth stages although maximum rates and preharvest intervals must be observed to prevent excess residue in the grain.

To a maximum rate of 80 mL/acre:

Chickpeas*

Dry beans (black, great northern, navy, pink, pinto, red)

To a maximum rate of 150 mL/acre:

Alfalfa (seedling only)
Canola
Field peas
Flax (including Solin)
Lentils
Mustard (brown, oriental, yellow)
Potatoes
Soybeans
Sunflowers

* Apply up to the 9 node stage (7 inches or 18 cm maximum height).

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

WEED	RATE (mL/ACRE)	ACRES TREATED PER 3 L CONTAINER	STAGE
Barnyard grass, green[Δ] and yellow foxtail[Δ], Persian darnel, Proso millet, Volunteer cereals (barley[Δ], canary seed, corn, oat[Δ], wheat[Δ]), Wild oat[Δ] (light infestations only*)	50**	60	[Δ]2 to 4 leaf stage when treated with the 60 acre rate. For best results, apply at the 2 to 3 leaf stage.
Moderate to heavy infestations of the above grasses	80	40	2 to 6 leaf stage. For best results, apply at the 2 to 3 leaf stage.
Quackgrass (suppression only)	80	40	2 to 6 leaf stage when 3 to 6 inches (6 to 15 cm) tall. For best results, apply at the 3 to 5 leaf stage.
Quackgrass (season long control)	150***	20	

* The manufacturers do not provide guidelines for weed densities under light infestations. When in doubt as to the level of weed infestation, use the higher rate or contact the manufacturer.

** At this rate, clethodim should not be tank mixed with any other pesticide and should only be applied under the following growing conditions: good crop stand, within the recommended leaf staging (2 to 3 leaf is optimum timing) prior to tillering, light weed infestations, adequate moisture and fertility, absence of stress, and good growing conditions.

*** Apply with 10 L of Amigo per 1000 L of spray solution

Amigo: 0.5 L of Amigo per 100 L of spray solution (unless otherwise indicated on the label)

Refer to the product labels for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

Select $11.08 to $33.25 per acre.

Centurion $11.06 to $33.20 per acre.

Application Information:

Water Volume: 20 to 40 L per acre. 40 L per acre under dense weed infestations or dense crop canopies.

Pressure: 40 psi (275 kPa).

Nozzles: Stainless steel 80° flat fan nozzles tilted forward at a 45° angle.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Clethodim will be less effective when plants are stressed by lack of moisture, excessive moisture, low temperature and/or very low relative humidity. Regrowth of tillers may occur if application is made under any of the above stress conditions.

Tank Mixes:

Clethodim may be tank mixed with other pesticides at the 80 to 150 mL/acre rates. Add the recommended amount of Amigo with all tank mixes unless otherwise indicated.

Herbicides:

In flax (not including solin):

Buctril M (0.4 L/acre)
Curtail M (0.6 to 0.8 L/ac)*
MCPA ester (rates for flax).
Lontrel at 0.23 to 0.34 L/acre.*

In solin (low linolenic flax varieties):

Buctril M (0.4 L/acre)

In canola:

Lontrel at 0.17 to 0.34 L/acre*.
Muster at 8 to 12 g/acre* (redroot pigweed is controlled at the 8 g/acre rate of Muster in this tankmix).

In Clearfield canola only:

Pursuit at 40 to 85 mL/acre* At the low rate of Pursuit, the following weeds are controlled: chickweed, hemp-nettle, redroot pigweed* stinkweed, volunteer canola (NOT CLEARFIELD varieties) wild buckwheat*, wild mustard, plus weeds listed on clethodim labels.
*Light infestations only. For heavy infestations, use high rate of Pursuit.

In Liberty Link canola only:

Liberty at 1.1 to 1.35 L/acre (clethodim at 25.5 mL/acre may be tank mixed with to enhance control of wild oat and volunteer cereals). When mixing add Amigo adjuvant to the water first, then Liberty, then clethodim. Consult labels for detailed mixing instructions.

In field peas:
Pursuit at 85 mL/acre.*

* Apply with the 80 mL/acre rate of clethodim only.

Allow 4 days between application of clethodim and any other chemical not recommended as a tank mix combination on the label.

Fertilizer: None registered.

Insecticide: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the clethodim labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Grazing: Do not graze or cut treated crops for forage until 60 days after application of clethodim to annual crops, and 30 days after application to seedling alfalfa.

Preharvest Interval: Do not apply within 60 days of harvesting canola, dry beans, flax (including Solin), lentils, potatoes, chickpeas (Desi and Kabuli) or mustard (brown, yellow, oriental). Do not apply within 75 days of harvesting soybeans or field peas. Do not apply within 30 days of harvesting seedling alfalfa. Do not apply within 72 days of harvesting sunflowers.

Aerial Application: Do not apply by air.

Storage: May be stored at any temperature. Shake well before use.

Other: Do not apply more than 0.15 L/acre to the same land area per season.

Environment: Leave a buffer of 15 meters between the last past of the sprayer and open water, wellheads, wetlands and other non-target areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Skin and Eye Irritant

Curtail M

Herbicide Group – 4
(Refer to page 27)

Company:

Dow AgroSciences (PCP#22764)

Formulation:

50 g/L clopyralid and 280 g/L MCPA ester formulated as an emulsifiable concentrate.

Container size - 8 L.

Crops and Staging:

Spring wheat (including durum), barley, oat, canaryseed* and timothy (established for hay and forage production)* at the 3 leaf to just before the flag leaf stage.

Flax (NOT including low linolenic acid varieties) at 2-6 inches (5 to 15 cm) height.

Since the application of Curtail M to established timothy and canaryseed is registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance.* ***Users of Curtail M on established timothy and canaryseed do so at their own risk.*

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

The following weeds are controlled at the 1 to 4 leaf stage unless specified:

At 0.61 L/acre (8L jug treats 13 acres) for light infestations of:

Canada thistle***
Flixweed**
Lamb's-quarters
Shepherd's-purse**
Stinkweed**
Volunteer sunflower
Wild mustard

At 0.8 L/acre (8L jug treats 10 acres) for medium to heavy infestations of the above weeds and:

Annual sow-thistle
Canada Thistle***
Common groundsel
Dandelion*
Kochia (suppression only)**
Perennial sow-thistle
(top growth control only)
Redroot pigweed
Russian pigweed
Scentless chamomile**
Smartweed
Tartary buckwheat
Volunteer canola
Wild buckwheat

* Spring rosettes only
** 2 to 4 leaf stage, (spring seedlings only for winter annuals).
*** Season long control, some regrowth may occur in the fall. Apply from the 4 inch (10 cm) to prebud stage.

Cost (2004 suggested retail price):

$11.00 per acre.

Application Information:

Water Volume:

Canary seed and timothy 40 to 80 L per acre
All other crops 40 to 60 L/acre

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan nozzles tilted forward at a 45° angle.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

When weeds are stressed because of drought, flooding, hot or cool (less than 15°C) temperatures, weeds are not actively growing and control may be reduced. Do not apply to weeds stressed longer than 20 days from lack of moisture as poor control can result.

Tank Mixes:

Curtail M at 0.8 L/acre should be used in all tank mixes unless otherwise indicated. See labels for adjuvant rates.

Herbicides:

In spring wheat (including durum) and barley:
Achieve Liquid at 0.20 L/acre plus Turbocharge adjuvant
Assert at 0.52 to 0.64 L/acre plus water pH adjuster
Puma^{120} Super (0.16 to 0.31 L/acre).

In spring wheat (not including durum) and barley:
Avenge at 1.4 to 1.7 L/acre (no adjuvant required)[Δ*]
Refine Extra at 8 g/acre plus a non-ionic surfactant with Curtail M at 0.61 L/acre.

In spring wheat (including durum):
Horizon at 92 to 116 mL/acre plus Score.
Everest at 17.4 g/acre plus Ag-Surf or Agral 90 adjuvant.

[Δ] Use the 0.66 to 0.8 L/acre rate of Curtail M only. The 0.66 L/acre rate will control weeds listed for the 0.61 L/acre rate above.

[*] See Avenge label for varietal restrictions. Allow a minimum of 5 days between the application of Curtail M plus Avenge and the application of any other pesticide, as poor weed control may result.

Check product labels for additional crop staging restrictions.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Curtail M label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Grazing: Do not graze treated fields or cut for hay within 7 days of application.

Preharvest Interval: Leave 60 days between application and harvest.

Recropping: Wheat, barley, oats, rye, corn, flax, canola, forage grasses and mustard may be planted the year after application. Do not underseed crops to forage legumes the year after treatment.

Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Do not sow *any other* crops until the second year after application. Apply manure bedded with straw from treated crops only to the crops listed above.

Aerial Application: Do not apply by air.

Storage: Store in a cool (above 5°C), dry area. If product is frozen, bring to room temperature and agitate before use.

Environment: Avoid drift onto sensitive plants and contamination of water or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Caution Eye Irritant

Dicamba

Herbicide Group – 4
(Refer to page 27)

Company:

BASF Canada (Banvell II) - PCP#23597

Gharda (Oracle) - PCP#26722 (distributed by United Agri Products and AgWest Inc.)

Formulation:

480 g/L dicamba formulated as a solution.
Container sizes - 2 x 10 L jugs.

Crops, Rates and Staging:

CROP	STAGE	RATE (L/acre)
Spring wheat*	2 to 5 leaf.	0.093 to 0.117
Barley*	2 to 5 leaf.	0.093
Oats*	2 to 5 leaf.	0.093 to 0.117
Canaryseed*	3 to 5 leaf.	0.117
Winter wheat*	In spring - 6 to 10 inches (15 to 25 cm) - prior to flag leaf.	0.093
Spring rye*	2 to 3 leaf	0.093 to 0.117
Corn, field	Broadcast up to 8 inches (20 cm). When higher, use drop-nozzles. Apply no later than 2 weeks prior to tassle emergence and prior to 20 inches (50 cm).	0.243
Corn, field + 2,4-D	Up to 4 inches (10 cm)	0.117
Red fescue	*Seedling:* 2 inches (5 cm) tall. *Established:* up to the flag leaf stage.	0.243
Pastures	Established and actively growing	0.84 to 1.86
Seedling grasses: Fescue (creeping red, meadow, tall), Meadow foxtail, Orchardgrass, Smooth bromegrass, Timothy, Wheatgrass (crested, intermediate, pubescent, slender, streambank, tall)	2 to 4 leaf	0.093 to 0.117
Fall stubble	Apply according to weed stage.	1.0
Fall stubble + glyphosate	Apply according to weed stage.	0.5
Pre-seeding cereals	Apply according to weed stage.	0.127
Chemfallow + 2,4-D	Apply according to weed stage.	0.093 to 0.117
Chemfallow + glyphosate	Apply according to weed stage.	0.243

* Should be mixed with a tank mix partner for broad spectrum control

Weeds, Rates and Staging:

Apply to annual broadleaf weeds at the 2 to 3 leaf stage and to winter annual rosettes up to 2 in. (5 cm) across.

Dicamba applied alone at 0.093 to 0.117 L/acre will control:

Cleavers (high rate only)
Cow cockle
Corn spurry
Canada thistle*
Lady's thumb
Perennial sow-thistle*
Smartweed (green)
Tartary buckwheat
Wild buckwheat

Dicamba at 0.25 to 0.5 L per acre control the weeds above plus:

Canada thistle**
Canada Fleabane
Field bindweed**
Mustard (hare's-ear, Indian, tumble, wild, wormseed)
Perennial sow-thistle**
Pigweed (redroot, Russian)
Ragweed (common, false, giant)

Dicamba at 0.85 L/acre will control the weeds above plus:

Curled dock*
English daisy
Goldenrod

Dicamba at 1.86 L/acre will control the weeds above plus:

Diffuse knapweed
Goat's-beard
Ground cherry
Pasture sage
Povertyweed
Sheep sorrel
Thyme-leaved spurge

* Top growth only.

** Three consecutive years of treatment are required for complete control.

Canada thistle, Perennial sow-thistle in summerfallow: Apply prior to the bud stage. Must be applied to thistle plants with 6 to 10 inches (15 to 25 cm) of new growth.

Brush control in pastures: When brush is actively growing and is 6 feet (2 m) in height or less (in spring or early summer).

Canada thistle control in fall stubble: When thistles exhibit new growth and at least 2 weeks prior to a killing frost.

Refer to label for full lists of weeds controlled by dicamba plus tank mixes in cereals, pastures, summerfallow and other situations.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Cost (2004 suggested retail price):

Banvell II Cost per L - $37.82.

Oracle Cost per Liter - $37.75

Application Information:

Water Volume:

Preseeding burnoff: 23 to 45 L per acre.
Annual crops: at least 45 L per acre.
Pastures, summerfallow and stubble: 45 to 90 L per acre.
Corn: 90 to 140 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop damage (stunting, reduced seed set) can occur if the chemical is applied at any time other than the recommended stage. Do not apply to crop under stress from adverse environmental conditions, such as excess moisture, drought and disease. Apply when air temperature is between 10 and 25°C.

Do not apply:

- when there is a risk of severe temperature fall in the night;
- under high humidity, temperatures above 30°C, or fog conditions, to prevent drift to sensitive crops;
- when wind is blowing toward a nearby sensitive crop;
- when winds are gusty up to 5 mph (8 km/hr).

Tank Mixes:

Herbicides:

	Spring wheat	Winter wheat	Barley	Oats	Seedling Grasses
2,4-D amine (500g ai/L) (0.34 L/acre)	✓	✓	✓		✓
MCPA Amine (0.34 L/acre)	✓	✓	✓	✓	✓
MCPA K (0.44 L/acre)	✓	✓	✓	✓	✓
Sencor (0.11 to 0.17 L/acre)	✓		✓		
Ally (2 g/acre)	✓		✓		

In Canaryseed: MCPA amine (0.34 L/acre)
In Corn, Spring rye: 2,4-D amine (500 g ai/L)(0.34 L/acre)
In Corn: Accent (13.5 g/acre) plus non-ionic surfactant
In Chemical fallow, stubble: 2,4-D, glyphosate products.
In Red fescue: 2,4-D amine (0.61 L/acre)
In Preseeding burnoff: glyphosate (0.38 L/acre)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Banvel II label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing and Harvest Intervals:

Canaryseed: Use only as birdseed.

Corn: Do not graze cattle or harvest for silage until 7 days after treatment of dicamba or dicamba + 2,4-D amine.

Cereals, seedling grasses, pasture: If treated vegetation has been consumed by meat animals within 30 days of dicamba application, feed the animal with untreated diet for 30 days before slaughter. Meat animals may graze or feed on treated pasture 30 days after dicamba application without restrictions on slaughter.

Dairy cattle: no restrictions up to 0.50 L/acre.

Use the following guidelines for feeding dicamba treated forage to dairy animals:

RATE/ACRE	DELAY BETWEEN TREATMENT AND HAYMAKING/ GRAZING (DAYS)
up to 0.5 L	0
0.501 L to 0.93 L	7
0.931 L to 1.86 L	14
1.861 L to 2.87 L	30

Recropping: Grow only cereals, corn, soybeans or white beans the year after treatment with the 1.0 L/acre rate. Grow only cereals, corn, field beans, soybeans or canola the year after applications of 0.5 L/acre. If applications are made after September 1, or if dry weather persists after application, crop injury may occur the following spring.

Aerial Application: May be applied by air on cereals only. Use a minimum of 8 L/acre.

Storage: May be stored at freezing temperatures.

Environment: Do not contaminate domestic or irrigation water. Do not apply around desirable trees or plants or near the root zone of these plants.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

▽ Caution Poison

Warning Eye Irritant

Dicamba+Mecoprop+MCPA

Herbicide Group – 4
(Refer to page 27)

Company:

Syngenta (Target) – PCP#13761
United Agri Products (Sword) – PCP#25583

Formulation:

275 g/L MCPA, 62.5 g/L mecoprop and 62.5 g/L dicamba formulated as a solution.
Container size - 10 L.

Crops and Staging:

Cereals:

CROP	STAGE
Barley	2 to 4 leaf (3 leaf for best crop safety)
Canaryseed, Oats, Spring wheat (including durum)	2 to 5 leaf (3 to 4 leaf for best crop safety)
Winter wheat	spring application only; up to 12 inches (30 cm) high (top leaf extended)
Summerfallow	Fall stubble

Seedling grasses grown for forage only (NOT for seed production)*:

Apply at the 2 to 4 leaf stage.

Creeping red fescue
Crested wheatgrass
Intermediate wheatgrass
Meadow foxtail
Orchardgrass
Smooth bromegrass
Timothy

Established grasses for forage only (NOT for seed production)*:

Apply up to flag leaf stage.

Bromegrass (meadow, smooth)
Fescue (creeping red, meadow, tall)
Kentucky bluegrass
Meadow foxtail
Orchardgrass
Timothy
Wheatgrass (crested, intermediate, pubescent, slender, streambank, tall, western)

* *NOTE: Use only one application per year by ground. Since applications to forage grasses in western Canada has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance.* ***Application to forage grasses are at the risk of the user.***

Cost (2004 suggested retail price):

Target: $13.52 per L.
Sword: $13.47 per L.

Weeds and Staging:

Weeds controlled at the 2 to 3 leaf stage unless otherwise indicated:

Annual sow-thistle
Ball mustard
Canada thistle (6 to 8 inches (15 to 20 cm) and actively growing)*
Cleavers (1-2 whorls)
Common ragweed
Corn spurry
Cow cockle
Field bindweed*
Flixweed
Green smartweed
Hedge bindweed*
Hemp-nettle (less than 2 pairs of true leaves)
Knotweed
Kochia
Lady's-thumb
Lamb's-quarters
Night-flowering catchfly
Perennial sow-thistle*
Prostrate pigweed
Redroot pigweed
Russian thistle (less than 2 inches-5 cm)
Shepherd's-purse
Stinkweed
Tall mustard
Tartary buckwheat
Volunteer canola
Volunteer sunflowers
Wild buckwheat
Wild mustard
Wormseed mustard
Yellow mustard

* Top Growth Control only

Rates:

0.4 to 0.6 L per acre (10 L treats 25 to 16.7 acres).

Use the higher rate under adverse weather conditions, when weed density is high, for cleavers control, winter annual control and for suppression of Canada thistle and perennial sow-thistle.

Although diacamba + mecoprop + MCPA is registered up to the 5 leaf stage of the crop for the rates listed here, the low rate should be used when the crop is at the 5 leaf stage for optimum crop safety.

For Canada thistle, post-harvest or summerfallow application, use 0.81 L/acre (one 10 L container treats 12.4 acres).

Application Information:

Water Volume: Minimum 40 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan with 50 mesh screens and filters.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Hot and dry or cold and wet weather prior to spraying may result in reduced weed control and increased crop injury. Do not apply within 2 weeks of a killing frost.

Tank Mixes:

Herbicides:
Spring wheat (including durum):
Horizon.
Wheat and Barley:
Sencor or linuron for chickweed control.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the dicamba +mecoprop+MCPA labels only. To check for other possible mixes see the blue chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Activity may be reduced if rainfall occurs within 3 hours of application. Contact manufacturer for more information.

Grazing: Do not graze or harvest for livestock feed within 7 days of application.

Preharvest: Leave at least 80 days from application to harvest.

Recropping: No restrictions the year after application.

Aerial Application: Both Target & Sword may be applied by air with a minimum of 12 L/acre water.

Storage: Do not freeze.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Dichlorprop + 2,4-D

Herbicide Group – 4

(Refer to page 27)

Company/Products:

United Agri Products (Turboprop 600 – PCP#16532)
Nufarm (Estaprop – PCP#14803)
IPCO (Dichlorprop-D – PCP#21717)

Formulation:

300 g/L of dichlorprop and 282 g/L of 2,4-D ester formulated as an emulsifiable concentrate.

Container size – 2 x 10 L.

Crops and Staging:

Wheat (spring, durum) and barley - 4 leaf until prior to the emergence of the flag leaf.

Winter wheat - in spring after tillering but prior to the emergence of the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Treat weeds when young and actively growing and before they are shielded by the crop. Additional stage restrictions are indicated.

Annual sow-thistle
Ball mustard
Bluebur
Burdock
Canada thistle*
Cocklebur
Curled dock*
Dog mustard
Flixweed
Hare's-ear mustard
Ragweed
Redroot pigweed
Round-leaved mallow
Russian pigweed
Russian thistle$^{\wedge\wedge\wedge}$
Shepherd's-purse
Smartweed$^{\wedge\wedge}$
Stinkweed
Stork's-bill
Tartary buckwheat$^{\wedge\wedge}$
Indian mustard
Kochia$^{\wedge\wedge\wedge}$
Lady's-thumb$^{\wedge\wedge}$
Lamb's-quarters
Night-flowering catchfly$^{\wedge}$
Oak-leaved goosefoot
Perennial sow-thistle*
Tumble mustard
Volunteer canola$^{\wedge\wedge}$
Volunteer sunflower$^{\wedge\wedge}$
Wild buckwheat$^{\wedge\wedge}$
Wild mustard
Wormseed mustard
Toadflax**

* Top growth control only
** Suppression only. Treat before the majority reach 6 inches (15 cm).
$^{\wedge}$ Spring annuals only
$^{\wedge\wedge}$ Treat prior to the 4 leaf stage
$^{\wedge\wedge\wedge}$ Treat up to 2 inches (5 cm)

Cost (2004 suggested retail price):

Dichlorporp-D: $6.00 per acre
Estaprop: $5.93 per acre
Turboprop: $6.22 per acre.

Rates:

0.71 L/acre (one 10 L container treats 14 acres).

Application Information:

Water Volume:
Ground: 20 to 97 L per acre*. Use a minimum of 40 L of water per acre to reduce the risk of drift.
Aerial: Minimum 12 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

* May vary by product. Check label closely.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Applications made under dry conditions may result in reduced control. Crops under stress from adverse environmental conditions, such as excess moisture, frost or drought, may be injured. Best weed control when adequate soil moisture is present and warm temperatures prevail. Do not apply when daytime temperatures exceed 27°C.

Tank Mixes:

Herbicides:

	Spring Wheat (including durum)	**Barley**
Estaprop	Avenge	Avenge
Turboprop	Achieve, Avenge, Assert, Horizon, Puma(2) Super	Achieve, Avenge, Puma(2) Super
Dichlorprop-D	Avenge	Avenge

Note: Always refer to the label or the Guide section of the tank mix partner for additional restrictions on staging and varieties.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on dichlorprop + 2,4-D labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Estaprop - Do not graze the treated crop or harvest for hay or feed within 40 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Dichlorprop-D/Turboprop 600 - Do not graze treated crops or cut for feed in the year of treatment.

Recropping: No restrictions the year after application.

Aerial Application: May be applied by air. Refer to specific product labels for full details for application by air.

Storage: May be frozen.

Environment: Do not spray when winds are calm or greater than 8 km/hr. Apply when downwind of sensitive non-target plants, open water and wetland areas. No specific distance is indicated for an upwind buffer zone from these habitats. Contact Manufacturer for details.

Tank Cleaning:

Manufacturers of this product warn that even after thorough cleaning, the use of a sprayer that has come in contact with this product may cause damage to susceptible crops.

Do not use spray equipment to apply other pesticides to crops sensitive to these products. To clean sprayer, rinse all parts several times with water, then fill sprayer with a water/ammonia solution (1 L of a 3% household ammonia solution per 100 L of water) and let stand for 24 hours. Rinse several times after with clean water.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Hazard Rating:

Warning Poison

Dual II Magnum

Herbicide Group – 15
(Refer to page 27)

Company:

Syngenta (PCP#25729)

Formulation:

915 g/L metolachlor formulated as an emulsifiable concentrate. Container size - 12 L.

Crops and Staging:

Preplant incorporated. In areas with good rainfall or under irrigation, may be applied as a pre-emergence surface treatment. When Dual II Magnum is applied to the soil surface, after planting but prior to emergence, at least 0.5 inches of water (1.25 cm) is required within 10 days of application for proper activity.

Corn (field, sweet, silage),
Dry beans (white, kidney
  and pinto)
Potatoes
Soybeans
Sweet white lupins

Weeds and Staging:

Apply prior to weed emergence.

American nightshade
Barnyard grass
Eastern black nightshade
Fall panicum
Giant foxtail
Green foxtail
Hairy crabgrass
Old witch grass
Redroot pigweed*
Smooth crabgrass
Yellow foxtail
Yellow nutsedge**

* Suppression only.,
** Preplant incorporated treatment only.

Cost (2004 suggested retail price):

$18.60 to $26.25 per acre.

Rates:

0.5 to 0.7 L/acre (12 L treats 24 to 17 acres). Use higher rates on heavy textured soils or when high populations of weeds are expected. Make only one application per season.

Application Information:

Water Volume: 70 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Use flat fan nozzles, 50 mesh screens.

Incorporation:

Apply to a firm seed bed free of large clods or lumps. If using tandem disks, set disks to work the soil at a depth of 4 inches (10 cm) and operate at a speed of 6 km/hr (4 mph). If using an S-tine cultivator, set the implement to work the soil to a depth of 4 inches (10 cm) and operate at a speed of 10 km/hr (6 mph). Incorporation equipment should include rolling or western harrows.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

A moderate rainfall or equivalent irrigation is required within 10 days to activate pre-emergent surface treatments. If rain does not occur, a shallow cultivation or use of a rotary hoe is necessary. Drought conditions that persist after any application may reduce annual grass control. On sandy soils, heavy rainfall following application may cause leaching of Dual II Magnum, resulting in reduced weed control.

Tank Mixes:

Herbicides:

In Corn: Aatrex in both PPI and pre-emergent applications.

Fertilizers: May be applied with liquid fertilizer. May be impregnated onto dry bulk fertilizers (except nitrate fertilizers, superphosphate fertilizers or limestone).

Insecticides: None registered.

Note: The above mixes are those listed on the Dual II Magnum label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: When PPI, 0.5 inches (1.25 cm) of rain is required after application for proper activity.

Grazing: Do not graze the treated immature crop or cut for hay. In corn, immature means before ear emergence.

Recropping: In the year of treatment, seed only corn, soybeans, white beans, potatoes, snap beans, lima beans, processing peas, sweet white lupins, or (a minimum of 4.5

months after application) winter cereals. If Dual II Magnum has been applied in a tank mix with another product, consult those products' labels for additional recropping restrictions.

Aerial Application: Do not apply by air.

Storage: May be frozen.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Eye Irritant

Potential Skin Sensitizer

DyVel

Herbicide Group – 4

(Refer to page 27)

Company:

BASF Canada (PCP#16545)

Formulation:

84 g/L of dicamba and 336 g/L of MCPA K^+ formulated as a solution.

Container size - 2 x 10 L jugs, 55 L, 110 L.

Crops and Staging:

Spring wheat (including durum), barley or oats - 2 to 5 leaf stage.

Winter wheat - apply in spring when crop is 6 to 10 inches (15 to 25 cm) tall but before shot blade stage.

Note: Crop damage can occur if applications are made at other than the recommended crop stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled in the 2 to 4 leaf stage unless otherwise stated:

Ball mustard
Burdock
Cleavers (suppression only)
Cocklebur
Common ragweed
Corn spurry (2 to 3 leaf)
Cow cockle (2 to 3 leaf)
False ragweed
Lamb's-quarters
Prostrate pigweed
Redroot pigweed
Russian pigweed
Russian thistle
Shepherd's-purse
Stinkweed
Tartary buckwheat
Flixweed
Giant ragweed
Green smartweed
Hare's-ear mustard
Hemp-nettle (2 to 3 leaf)
Indian mustard
Kochia
Lady's-thumb
Tumble mustard
Wild buckwheat
Wild mustard
Wormseed mustard
Wild radish
Volunteer canola (2 to 4 leaf)
Volunteer sunflowers

Top growth control:

Canada thistle
Perennial sow-thistle

Cost (2004 suggested retail price):

$5.82 per acre.

Rate:

0.5 L/acre (one 10 L jug treats 20 acres).

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 to 45 psi (275 to 310 kPa). To reduce the risk of drift damage to sensitive non-target crops, 20 to 30 psi (150 to 200 kPa) and higher water volumes are recommended.

Nozzles: Flat fan. Select nozzles that produce a low number of fine droplets to reduce the risk of drift.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

For best weed control, apply when temperature is between 10 and 25°C. Do not treat crops under stress from excessive moisture or drought. To avoid crop injury, do not apply when temperature is expected to exceed 30°C, or when there is a risk of a severe drop in overnight temperature.

Tank Mixes:

Herbicides:

In Spring wheat:

Horizon (95 mL/acre) (wild oat control only), plus Score adjuvant

Horizon (115 mL/acre) (wild oat, green and yellow foxtail), plus Score adjuvant.

Everest – 17.4 g/acre plus non-ionic surfactant

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the DyVel label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze treated crop or cut for hay within 30 days of application.

Preharvest Interval: Leave 60 days between application and harvest.

Recropping: No restrictions the year after treatment.

Aerial Application: May be applied by air. Use a minimum of 8 L/acre water volume.

Storage: May be frozen.

Environment: Avoid drift onto sensitive plants. Leave a distance of 15 m between the outside edge of the sprayed area and sources of domestic, irrigation or natural water or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Caution Poison

DyVel DSp

Herbicide Group – 4

(Refer to page 27)

Company:

BASF Canada (PCP#27856)

Formulation:

110 g/L dicamba, 295 g/L 2,4-D amine and 80 g/L mecoprop-p formulated as a solution.

Container size - 2 x 10 L jugs, 55 L, 110 L.

Crops, Rates and Staging:

CROP	STAGE	RATE	
		(L/ACRE)	ACRES PER 10L JUG
Spring wheat (including durum)	3 to 5 leaf	0.34 to 0.45	29 to 22
Barley	2 to 3 leaf	0.34	29
Winter wheat	before crop is 12 inches (30 cm) tall in spring	0.34 to 0.45	29 to 22
Corn (field and sweet)	before corn reaches 6 inches (15 cm) in height with the top leaf extended	0.34 to 0.45	29 to 22
Native range and permanent grass pasture*	Established	1.3	7.7
Fall stubble, summerfallow	Stage according to weed	0.45 to 0.71	22 to 14

* Legumes will be severely injured by this application.

Applications outside the recommended stage may result in crop injury.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

The low registered rate for each crop will control the following weeds at the 2 to 3 leaf stage unless otherwise indicated:

Annual smartweed
Annual sow-thistle
Cocklebur
Common ragweed
Corn spurry
Hedge bindweed
Knotweed
Kochia
Lady's-thumb
Lamb's-quarters
Mustards (wild, ball, tall, wormseed, yellow)
Prostrate pigweed
Redroot pigweed
Russian thistle
Sow thistle
Stinkweed*
Volunteer canola (2 to 4 leaf, prior to bolting)
Volunteer tame buckwheat
Wild buckwheat

Use the high registered rate for each crop to control the following weeds:

Canada thistle (top growth only)***
Cleavers (1 to 2 whorls)**
Cow cockle
Field bindweed
Flixweed*
Jerusalem artichoke
Round-leaved mallow**
Shepherd's-purse

* Rosette stage in winter wheat.

** Suppression only.

***Canada thistle should be treated when 6 to 8 inches (15 to 20 cm) of new growth is present in Fall Stubble and in the early bud stage in Summerfallow.

Rates for Native Range and Pasture will control:

Alder
Bull thistle
Chicory
Goat's-beard
Poison ivy
Ragwort
Sheep laurel
White cockle

The high rate for each crop should be used for all weeds under adverse growing conditions, when weeds are at an advanced stage of growth or when weed densities are high. Guidelines are not provided for weed densities under light or heavy infestations. When in doubt as to the infestation level, use the high rate or contact the manufacturer. NOTE: *It is possible that poisonous plants such as ragworts, hemlocks and death camas could be more palatable to livestock after treatment with DyVel DSp. Suitable precautions should be taken to avoid livestock access when such plants are present.*

Cost (2004 suggested retail price):

$4.73 to $6.28 per acre in crop applications.
$6.28 to $9.88 per acre for fall stubble and summer fallow.
$18.10 per acre for pasture

Application Information:

Water Volume: Minimum 40 L per acre in cereals. Corn - 80 to 140 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crops under stress from excess moisture, drought or disease may suffer a setback when this herbicide is applied. Do not apply when temperature exceeds 27°C or when relative humidity is high. Stubble treatments for thistle control in fall should be made at least 2 weeks prior to killing frost.

Do not apply DyVel DSp at wind speed greater than 5 mph (8 km/hr).

Tank Mixes:

Herbicides:
In corn:
Atrazine.
In spring wheat (NOT including durum):
Everest (17.4 g/acre plus non-ionic surfactant).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the DyVel DSp label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not harvest for livestock feed within 30 days of application. Do not permit lactating dairy animals to graze fields within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Recropping: No restrictions the year after treatment.

Aerial Application: Do not apply by air.

Storage: May be frozen.

Environment: If there are sensitive plants within 400 m, apply only when there is a light breeze away from the sensitive area. Do not contaminate wetlands or water used for domestic or livestock consumption, irrigation or natural habitat.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

Eclipse

Herbicide Group – 4, 9
(Refer to page 27)

This product is a prepackaged tank mix of Eclipse A (Lontrel page 149) and Eclipse B (Vantage Plus pages 119 to 128). Information listed is restricted to Crop, Weeds, Rates and Cost and other important details. For other detailed information on the component products see the product pages listed above.

Company:

Dow AgroSciences

Formulation:

Eclipse A (PCP#26633): 360 g/L clopyralid formulated as a solution.

Eclipse B (PCP#26634): 360 g/L glyphosate present as an isopropylamine salt and formulated as a solution.

Container size – Each case will treat 30 acres and will consist of 3 split jugs (10 acres/jug). Each split jug will contain 1.12 L of Eclipse A and 5 L of Eclipse B.

Crops and Staging:

Glyphosate tolerant (Roundup Ready) canola varieties only in the 2 to 6 leaf stage. Some yellowing may occur when applied at the 4 to 6 leaf stage. This effect is temporary and will not influence crop growth, maturity or yield.

Weeds and Staging:

No staging is specified on the label.

Annual grasses:

Green foxtail
Wild oat
Volunteer barley
Volunteer wheat

Annual broadleaf weeds:

Chickweed
Cleavers
Corn spurry
Cow cockle
Hemp nettle
Kochia
Lady's-thumb
Lamb's-quarters
Night-flowering catchfly
Redroot pigweed
Russian thistle
Shepherd's-purse
Smartweed
Stinkweed
Wild buckwheat
Wild mustard
Volunteer canola*
Wild tomato

Perennial weeds (season long control):

Canada thistle
Dandelion less than 15 cm diameter**
Dandelion greater than 15 cm diameter***
Perennial sow-thistle**
Quackgrass

* Not including glyphosate tolerant (Roundup Ready) varieties
** Top growth only
*** Suppression only.

Cost (2004 suggested retail price):

$13.90 per acre.

Rates:

Apply Eclipse A at 0.112 L/acre and Eclipse B at 0.5 L/acre or 10 acres/jug (30 acres/case).

To prepare spray solution, add Eclipse A to the spray tank. Once it is half filled with water, add Eclipse B as the remaining water is added to the tank.

Application Information:

Water Volume: 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa)

Nozzles: Use nozzles that give even coverage without producing small droplets that are prone to drifting. Do not use with galvanized sprayer tanks since explosive hydrogen gas can be produced.

Restrictions:

Recropping: Wheat, oats, barley, rye (not underseeded to legumes such as alfalfa and clover), forage grasses, flax, canola, mustard and field peas*can be grown the year after application. Apply manure bedded with straw from treated crops only to the crops listed above excluding field peas.

**Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field peas following drought conditions in the previous year.*

Aerial application: Do not apply by air.

Edge

Herbicide Group – 3
(Refer to page 27)

Company:

Dow AgroSciences (PCP#20980)

Formulation:

Edge Granular - 5% ethalfluralin formulated as a granular.
Container size - 25 kg or 544 kg.

Crops and Staging:

Edge can be applied prior to seeding the following crops:

Seedling alfalfa (seed production only)
Canola
Caraway
Coriander
Dry beans (white or kidney)
Fababeans
Lentils (fall application only)[†]
Mustard (yellow only)
Peas
Safflower
Soybeans
Sunflowers

Weeds and Staging:

For pre-emergent control of the following weeds:

Grasses:
Barnyard grass
Crabgrass
Foxtail (Green & Yellow)
Volunteer barley*
Volunteer spring wheat
Wild oat

Broadleaf Weeds:
Cleavers*
Chickweed
Cow cockle
Corn spurry
Hemp-nettle*
Kochia
Lady's-thumb*
Lamb's-quarters
Nightshade*
Prostrate pigweed
Purslane
Redroot pigweed
Russian thistle*
Wild buckwheat

* Suppression only.

Rates:

TIME OF APPLICATION	RATE (KG/ACRE)				
	LIGHT TEXTURED SOILS		MEDIUM TO HEAVY TEXTURED SOILS		
	2-6% Organic Matter/Dark Brown-Black	6-15% Organic Matter Deep Black	2-4% Organic Matter Dark Brown	4-6% Organic Matter Black	6-15% Organic Matter Deep Black
Spring	6.9	8.9	6.9	8.9	8.9 - 11.3
Fall *	8.9	11.3	8.9	11.3	11.3

* To conserve crop residue, one incorporation may be completed in the fall and the second incorporation completed in the spring prior to planting.

Do not apply Edge to peat soils, soils containing less than 2 percent organic matter or soils containing greater than 15 percent organic matter. Application to eroded knolls or grey-wooded soils with highly variable texture or organic matter may result in reduced crop stand, delayed development or reduced yields in either the treated or rotational crop. To reduce the possibility of injury to the treated crop, use good quality certified seed. Seed shallow into a warm, moist, firm seedbed using recommended agronomic practices that will promote rapid and even crop germination and emergence.

[†] **Special instructions for lentils:**
Edge is registered for use on lentils for fall application only. One incorporation must be completed in the fall. Seeding depth is critical - do not seed more than 1.5 inches (4 cm) deep. Avoid loose seedbeds and planting into cold soils.

Cost (2004 suggested retail prices):

$14.99 to $25.00 per acre.

Application Information:

Equipment: Apply Edge Granular using a calibrated granular applicator.

Incorporation: Two incorporations are required at right angles for thorough mixing. The first incorporation must be completed within 24 hours of application. When using the granular formulation, delay the second incorporation for at least three days after the first. If applying Edge granular in the fall, it is preferred that both incorporations be done in the fall. The second incorporation may be delayed until spring to conserve trash; however, both incorporations must be done to the same depth.

Incorporate with a tandem disc, discer or field cultivator (Vibrashank type). A cultivator should have 3 to 4 rows of sweeps spaced 8 inches apart and staggered so that no soil is left unturned. Set equipment to work at a depth of 3 to 4 inches (8 to 10 cm). Operate disc implements at 4 to 6 mph (7 to 10 km/hr), cultivators at 6 to 8 mph (10 to 13 km/hr).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crops stressed by cold weather, excessive moisture or drought may be injured by Edge. Dry soil conditions between application and emergence may result in decreased weed control.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No effect once incorporated.

Grazing: Do not graze or cut treated crops for livestock feed prior to crop maturity.

Recropping: Do NOT grow - sugar beets, oats, and small-seeded annual grasses such as timothy, canaryseed and creeping red fescue in rotation following a crop treated with Edge. Do not seed wheat as a rotational crop onto land that has been treated with trifluralin and/or ethalfluralin at oilseed/special crop/barley rates for 2 consecutive crop years Thinning of crop may occur in areas that have received abnormally low amounts of precipitation or in crops that are emerging slowly.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. Edge Granular may be frozen. Do not expose to prolonged sunlight or heat.

Environment: Toxic to fish and other aquatic organisms. Do not contaminate water bodies or wetland areas.

Tank Cleaning:

Refer to page 14.

Elite

Herbicide Group – 1,4,6

(Refer to page 27)

This product is a prepackaged tank mix of Mextrol (Page 81) and Poast Ultra (Page 162). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Nufarm

Formulation:

The Elite package has 2 components:

Mextrol (PCP#25342): 200g/L bromoxynil and 200 g/L MCPA formulated as an emulsifiable concentrate.

Poast Ultra (PCP#24835): 450g/L of sethoxydim formulated as an emulsifiable concentrate.

Merge (PCP#24702): adjuvant.

Container size:
Mextrol – 11.36 L; Poast Ultra – 3.65 L; Merge – 8.1 L

One case treats 20 acres

Crops and Staging:

Flax (including Solin varieties) - 2 to 4 inches (5-10 cm)

Weeds and Staging:

Grass weeds:

Barnyard grass
Fall panicum
Green & yellow foxtail
Quackgrass*
Volunteer cereals
Wild oats
Witchgrass

Broadleaf weeds up to 4 leaf stage

American nightshade
Bluebur
Ball mustard
Canada thistle*
Cocklebur
Cow cockle
Flixweed
Green smartweed
Kochia**
Lady's-thumb
Night-flowering catchfly
Pale smartweed
Perennial sow-thistle*
Russian thistle**
Scentless chamomile***
Shepherd's-purse
Volunteer canola
Volunteer sunflower

Broadleaf weeds up to 8 leaf stage:

Common groundsel
Common ragweed
Lamb's-quarters
Stinkweed
Tame buckwheat
Tartary buckwheat
Wild buckwheat
Wild mustard
Wormseed mustard

* Suppression only
** control before plants are 2 inches high
*** spring seedlings only

Cost (2004 suggested retail price):

$21.65 per acre.

Rate:

Mextrol - 0.57 L per acre
Poast Ultra - 0.19 L per acre
Merge - 0.4 L per acre

One case treats 20 acres

Application Information:

Water Volume: 20 to 40 L per acre.

Pressure: 40 psi (275 kPa)

Nozzles: Flat fan nozzles

Eptam 8-E

Herbicide Group – 8

(Refer to page 27)

Company:

Gowan Company

Formulation:

800 g/L of EPTC formulated as an emulsifiable concentrate. Container size - 10 L.

Caution:

The level of weed control may be reduced where Eptam 8-E is used on soils that have been treated with Eradicane or Eptam 8-E the previous growing season. It is expected that the reduction in control will be greater where Eradicane or Eptam 8-E have been used repeatedly for 2 or more years.

Crops, Rates and Staging:

Eptam is applied as a preplant incorporated treatment prior to seeding the following crops:

CROP	RATE (L/ACRE)	ACRES TREATED PER 10 L CONTAINER
Dry and snap beans	1.72 - 2.23	5.8 - 4.5
Alfalfa, Bird's-foot trefoil Cicer milkvetch** Sweet clover** Sunflowers*	1.72	5.8
Potatoes	1.72 - 3.44	5.8 - 2.9
Flax*	1.42 - 1.72	7.0 - 5.8

* May also be applied in late fall prior to freeze-up
** Seed production only

NOTE: The use of Eptam 8-E on flax is not recommended in Saskatchewan because of the risk of crop injury.

Where a rate range appears, use the lower rate on light textured soils and the higher rate on heavy textured soils. Do not use on soils with less than 3 percent organic matter or more than 15 percent organic matter.

Weeds and Staging:

Must be applied prior to the emergence of the following weeds. Emerged weeds will not be controlled.

Barnyard grass
Chickweed*
Corn spurry*
Green foxtail
Hairy nightshade*
Henbit *
Lamb's-quarters*
Nettleleaf goosefoot*
Purslane*
Pigweed (prostrate, redroot, tumble)*
Quackgrass (suppression)**
Volunteer cereals (wheat, barley, oats)
Yellow foxtail
Yellow nutsedge**

* Will be controlled only if treatment is made when conditions are favourable for germination and growth.

** Roots of perennial weeds must be thoroughly chopped prior to application.

Cost (2004 suggested retail price):

$19.50 to $38.00 per acre

Application Information:

Carrier Volume: Minimum of 40 L per acre of water or liquid fertilizer (see label for liquid fertilizer compatibility).

Pressure: 30 to 40 psi (200 to 275 kPa).

Equipment and Nozzles: Since Eptam 8-E is highly volatile, the product must be incorporated immediately. This is best accomplished by mounting spraying equipment directly onto the incorporation equipment (tandem disks, field cultivators on light soil).

May also be applied to cleanly cultivated soil for potatoes, by metering into the irrigation water to achieve the recommended rate per acre ("herbigation" or "chemigation"). See label for detailed instructions.

Incorporation: All growth and stubble should be thoroughly worked into the soil before treatment. Apply to a dry soil surface. Incorporate immediately after application preferably during the spraying operation as Eptam 8-E is volatile. Set disc and cultivator implements to cut to a depth of 4 to 6 inches (10 to 15 cm). A second operation at a right angle to the first is required. The disc or cultivator must be followed with a harrow or other levelling device that extends beyond the width of the implement. Speeds in excess of 5 mph (8 km/h) will result in excessive pulverization and trash destruction leaving the field susceptible to erosion. The maximum recommended tillage depth is 4 inches (10 cm).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop injury can occur if stressful environmental conditions (cold, wet soils, drought or excessive heat) prevail after seeding. To minimize crop injury, delay seeding 10 days if these conditions prevail at the time of application, or select an alternative product. Very cold or dry soil conditions during weed emergence will reduce control.

Tank Mixes:

Herbicides:

Dry beans (white and red kidney only): Liquid formulations of Treflan and Rival.

Fertilizers: May be mixed with liquid fertilizer. Compatibility test should be conducted according to instructions on the herbicide label.

Dry bulk fertilizers, except nitrate fertilizers, may be impregnated or coated with Eptam 8-E. The impregnated fertilizer should be spread uniformly onto the field using a double overlap pattern immediately after impregnation. The impregnated fertilizer must be applied to the field when the soil surface is dry to at least 1/2 inch (1.5 cm) depth. The first incorporation must be done immediately after application.

Insecticides: Do not tank mix with insecticides.

Fungicides: None registered.

Note: The above mixes are those listed on the Eptam label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No effect once incorporated. Do not apply prior to pre-irrigation.

Grazing: Do not graze or feed treated crops to livestock in the year of application.

Recropping: Will not injure crops the year after spring application.

Aerial Application: Do not apply by air.

Storage: May be frozen.

Environment: Do not apply within 15 m of fish bearing waters or wildlife habitat.

Soil Type: Do not use on soils with less than 3 percent organic matter as crop injury will result.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Eradicane 8-E

Herbicide Group – 8

(Refer to page 27)

Company:

Gowan Company

Formulation:

800 g/L of EPTC (plus safener) formulated as an emulsifiable concentrate.

Container size - 10 L.

Caution:

The level of weed control may be reduced where Eradicane 8-E is used on soils that have been treated with Eradicane 8-E or Eptam 8-E the previous growing season. It is expected that the reduction in control will be greater where Eradicane 8-E or Eptam 8-E has been used repeatedly for 2 or more years.

Crops and Staging:

Corn (field, sweet* and silage). Eradicane is applied as a preplant incorporated treatment.

* Maximum 2.23 L/acre rate.

Weeds and Staging:

Must be applied prior to the emergence of the following weeds. Emerged weeds will not be controlled.

Barnyard grass
Chickweed
Corn spurry
Green and yellow foxtail
Hairy nightshade
Henbit
Lamb's-quarters
Pigweed (prostrate, redroot, tumble)
Purslane
Quackgrass (suppression)
Volunteer cereals (wheat, barley, oats)
Wild oats
Yellow nutsedge

Rates:

USE	RATE (L/ACRE)	ACRES TREATED PER 10 L CONTAINER
Annual weed control in light textured soils	1.72	5.8
Annual weed control in heavy textured soils	2.23	4.5
Quackgrass suppression	3.44	2.9

Cost (2004 suggested retail price):
$19.50 to $38.00 per acre.

Application Information:

Carrier Volume: Minimum of 40 L per acre of water or liquid fertilizer (See label for liquid fertilizer compatibility).

Pressure: 30 to 40 psi (200 to 275 kPa).

Equipment and Nozzles: Since Eradicane 8-E is highly volatile, the product must be incorporated immediately. This is best accomplished by mounting spraying equipment directly onto the incorporation equipment (tandem disks, field cultivators on light soil).

Incorporation:

All growth and stubble should be thoroughly worked into the soil before treatment. Apply to a dry soil surface. Incorporate immediately after application, preferably during the spraying operation, as Eradicane 8-E is volatile. Set disc and cultivator implements to cut to a depth of 4 to 6 inches (10 to 15 cm). A second operation at a right angle to the first is required. The disc or cultivator must be followed with a harrow or other levelling device that extends beyond the width of the implement.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Poor weed control may result if soils are wet during incorporation.

Crop injury may result if soil is cold and wet during germination and emergence. Hot, dry conditions during germination and early crop growth may also lead to crop injury.

Tank Mixes:

Herbicides: Atrazine.

Fertilizers: Dry bulk fertilizers, except nitrate fertilizers, may be impregnated or coated with Eradicane 8-E. The impregnated fertilizer should be spread uniformly onto the field using a double overlap pattern immediately after impregnation. The impregnated fertilizer must be applied to the field when the soil surface is dry to at least 1/2 inch (1.5 cm) depth. The first incorporation must be done immediately after application.

May also be mixed with liquid fertilizer. Conduct a compatibility test prior to mixing in tank. See label for instructions on conducting a compatibility test.

Insecticides: Do not tank mix with insecticides.

Note: The above mixes are those listed on the Eradicane 8-E label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No effect once incorporated.

Grazing: Do not graze treated crops or cut for feed prior to ear emergence.

Recropping: Will not injure crops the year after spring application.

Aerial Application: No labelled restrictions, however, since this product must be incorporated within seconds of application, aerial applications are impractical.

Storage: May be frozen.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

▽ Caution Poison

Escort

Herbicide Group – 2

(Refer to page 27)

Company:

E. I. duPont Canada (PCP#23005)

Formulation:

60% metsulfuron methyl formulated as a dry flowable. Container size - 0.25 kg.

Crops and Staging:

Pasture, rough turf, and rangeland.

No stage restrictions.

Weeds, Rates and Staging:

For seedling weeds apply to young plants up to 4 inches (10 cm) tall or wide. For established non-woody plants (biennial or perennial) apply up to the early bud stage. For western snowberry, wild rose and other woody species, apply between mid-June and mid-August after the brush has leafed out, but before the leaves begin to turn their fall colours.

Rate	Weeds Controlled	
8 g/acre	Canada thistle* Dandelion* Russian thistle Sow-thistle*	Common tansy Kochia Scentless chamomile Sweet Clover
10 g/acre	Above weeds plus: Canada thistle* Sow-thistle*	 Dandelion* Western snowberry
12 g/acre	Above weeds plus: Wild rose Canada thistle*	 Dandelion Sow-thistle*
40 g/acre**	Balsam poplar	Willow
60 g/acre**	Cherry	Trembling Aspen

At all rates add Agral 90, Agsurf, or Citowett at 0.2 L per 100 L of spray solution.

* Suppression only.

** Rangeland only. See label for detailed application instructions.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$13.76 to $20.64 per acre for weeds and up to $103 per acre for woody species.

Application Information

Water Volume: 40 to 91 L per acre for weedy growth and up to 809 L per acre applied to the point of run-off for woody species. See the label for details.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply during periods of intense rainfall or to soil saturated with water. Warm, moist conditions following treatment enhance the activity of Escort, while cold, dry conditions may reduce or delay activity. Brush hardened off by cold weather and drought stress may not be controlled.

Tank Mixes:

Herbicides: 2,4-D amine or ester at 0.79 L/acre (500 g/L formulations) plus surfactant.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: May be grazed on the day of treatment.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. May be frozen.

Environment: Leave a 15 m buffer from the down-wind edge of the spray boom to sensitive areas for rates up to 12g/acre and 45 m for rates up to 60g/acre

Tank Cleaning:

Escort can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray Escort should be flushed out immediately after use. The manufacturer recommends that sprayers used to apply this product be flushed twice with water/ammonia solution (1 L of 3% household ammonia per 100 L of water). All nozzles, screens and filters should be removed and cleaned with water/ammonia solution after applying this product. Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 14 for more information on tank cleaning.

Hazard Rating:

Caution Eye Irritant

Everest

Herbicide Group – 2
(Refer to page 27)

Company:

Arvesta (PCP#26448)

Formulation:

70% flucarbazone-sodium formulated as a water dispersible granule.

Container size - 4 x 174 g water-soluble packets.

Crops and Staging:

Spring wheat (including durum) with 1 leaf to a maximum of 4 main stem leaves plus 2 tillers (6 total leaves).

Everest MUST be tank mixed with a broadleaf partner for crop safety. Tank mix options are listed in the tank mix section.

Note: Several of the tank mix partners have more limiting staging than Everest. When tank mixing use the most restrictive application state or injury may result.

Weeds, Rates and Staging:

Green foxtail at 8.7 g per acre (one packet treats 20 acres)

Wild oat at 17.4 g per acre (one packet treats 10 acres)

Apply from 1 leaf to a maximum of 4 main stem leaves plus 2 tillers (6 total leaves).

Add non-ionic surfactant (Agral 90, Agsurf, Surf 92, Super Spreader, LI700) at 0.25 L per 100 L of spray solution.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$8.06 to $16.11 per acre, not including adjuvant and required tank mix partner.

Application Information:

Water Volume:
Spring wheat: 20 to 40 L per acre.
Note: Only Agral 90 and Ag-Surf may be used at the 20 L per acre water rate.
Durum wheat and reduced rate for green foxtail control: 40 L per acre.

Pressure: 30 to 50 psi (200 to 345 kPa)

Nozzles: Flat Fan nozzles at 80° or 110°. Use of flood-jet or controlled droplet application equipment is not recommended.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop tolerance and weed control may be reduced if applications are made to plants growing under stress. Stress includes saturated or water-logged soil, drought, extreme temperatures, low fertility or visible disease symptoms at application. Adopting practices to increase crop vigor will improve crop tolerance.

Tank Mixes:

Herbicides:
Note: All mixes must be applied with a registered surfactant unless otherwise indicated. Only one registered surfactant is required.

In spring wheat (including durum):
2,4-D Amine or Ester at recommended rates up to 0.34 L/acre (500 g/L formulation)^■
Frontline (40 acres per case)*
Spectrum (20 acres per case)*

In spring wheat (NOT including durum):
2,4-D Amine or Ester at recommended rates up to 0.45 L/acre (500 g/L formulation)^■
Ally at 2 to 3 g/acre + 2,4-D■ amine or ester up to rates recommended above*^
Attain (label rate)*^■
Buctril M (0.4 L/acre)
Curtail M (0.6 to 0.8 L/acre)*^
•DyVel (0.5 L/acre)*^
•DyVel DS (0.34 L/acre)*^■
Estaprop (0.71 L/acre)■
Express Pack (label rate)*■
Frontline 2,4-D (label rate)*^■
MCPA Amine or Ester at recommended rates up to 0.45 L/acre (500 g/L formulation)*
Pardner (0.4 L/acre)*
Prestige (labeled rate)*^
Refine Extra (8 g/acre) + 2,4-D■ (ester* or amine) at rates on this page.
•Target (0.4 to 0.6 L/acre)*^
Thumper (0.4 L/acre)■
Unity (40 acres per case)*^

* Registered for use with wild oat rate of Everest and with Agral 90 and Agsurf surfactants only
^ Apply in 40 L/acre of water only.
• Wild oat control may be reduced with this mix.
■ Mixes containing 2,4-D provide improved crop safety.

Fertilizers: Do not use liquid fertilizer or products that contain boron or free chlorine with this product or water-soluble bags will become insoluble and potentially plug sprayers.

Insecticides: None

Note: The above mixes are those listed on the Everest label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rainfall within 1 hour of application may reduce control.

Grazing: Do not graze treated fields. Mature grain or straw may be fed to livestock.

Preharvest: Leave at least 80 days from application to harvest.

Recropping Interval: Follow the chart below:

Soil Zones and Rotational Crops			
Grey-Wooded	**Black**	**Dark Brown**	**Brown**
Spring Wheat Barley Canola (all varieties) Field Peas*	Spring Wheat Barley Canola (all varieties) Field Peas* Flax Field Bean	Spring Wheat Barley Canola (all varieties) Field Peas* Flax	Spring Wheat

* Field peas may be grown the year following Everest application in fields where precipitation has been equal to or above the 10 year average during the growing season, and where organic matter content is above 4%, and pH is below 7.5. The company suggests a minimum of 100 mm (4 inches) of rain is needed in the 60 days following application for adequate breakdown to take place.

NOTE: Other rotational crops may also be affected if rainfall is less than the 10 year average for the area. Soils in the grey wooded, black and dark brown soil zones with a combination of low organic matter (less than 2%), light textured soils or high pH (greater than 7.5) (i.e. eroded knolls, sandy soils) may result in delayed growth and development in rotational crops.

Aerial Application: Do not apply by air.

Storage: Store in closed original container in a cool, dry area away from fertilizers, food or feed. Handle frozen solupaks with care since they may be fragile at low temperatures.

Environment: Leave at least 20 m from the downwind edge of the spray swath to sensitive plants like shelterbelts and woodlots and at least 35 m to well heads, sloughs, dugouts, ponds, marshes, ditches, streams, lakes, etc. Avoid drift onto sensitive crops like canola and tame oat. Do not mix or load within 10 m of wellheads, sloughs, dugouts, ponds, marshes, ditches, streams, lakes, etc.

Tank Cleaning:

Everest residues in the spray tank can cause severe injury to sensitive crops at very low concentrations. Sprayers should be cleaned out immediately before using another product.

Follow the steps below:

1. Drain the tank and thoroughly rinse the spray tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with ammonia/water solution (1 L of 3% household ammonia per 100L of water) and flush the hoses, boom and nozzles with the solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank.
3. Remove all nozzles, screens and filters and clean in a separate container using an ammonia/water solution as above.
4. Repeat #2.
5. Rinse tank, boom and hoses with clean water. And wash any residue from the outside of the tank.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk. For additional information on tank cleaning see page 14.

Hazard Rating:

Keep out of reach of children

Express Pack/Express TNG

Herbicide Group – 2
(mixed with 4 or 9)

(Refer to page 27)

Company:

E. I. duPont Canada

Formulation:

Express TNG (PCP#27532): 75% tribenuron methyl, formulated as a dry flowable. Container size – 4 x 80 g water soluble packets.

Express TNG can be purchased separately but must be used in a mix with 2,4-D Ester or glyphosate.

The Express Pack package contains 2 containers:

Express (PCP#22335): 75% tribenuron methyl, formulated as a dry flowable. Container size - 160 grams.

2,4-D herbicide (PCP#23192): 660 g/L 2,4-D LV ester, formulated as an emulsifiable concentrate. Container size - 10 L.

Crops and Staging:

Express Pack: Wheat (spring and durum), barley - 3 leaf up to emergence of the flag leaf

Summerfallow.

Express TNG + glyphosate: One day prior to seeding barley and spring wheat (including durum) and Summerfallow.

Weeds, Rates and Staging:

In-crop or fallow:

Express TNG 4 g/acre plus 0.24 L/acre 2,4-D LV Ester 700. (Express Pack case treats 40 acres) applied at up to 4 inches (10 cm) tall or across, unless otherwise indicated:

Annual sunflower
Ball mustard
Canada thistle*
Cow cockle
Flixweed^^
Hare's-ear mustard
Indian mustard
Kochia^^^
Lamb's-quarters
Narrow-leaved
  hawk's beard^^
Prickly lettuce
Redroot pigweed
Russian pigweed
Russian thistle
Shepherd's-purse^^
Stinkweed^^
Sweet clover
Thyme-leaved spurge
Tumble mustard
Wild mustard
Wild radish
Wormseed mustard
Wild buckwheat**^

Preseed prior to cereals or in fallow:

Express TNG 4 g/acre (one package of Express TNG treats 80 acres) plus Roundup Original or Transorb at 0.3 to 0.4 L/acre plus Agral 90 at 375mL per 100 L of solution:

Weeds controlled by Roundup at these rates plus:

Canada thistle rosettes**
  (0.4 L/acre rate only)
Dandelion (up to non-
  flowering rosettes)
Kochia
Lamb's-quarters
Narrow-leaved hawk's-beard**
Redroot pigweed
Russian thistle
Volunteer canola
(all varieties)

Express TNG 4 g/acre plus Roundup Transorb at 0.5 L/acre or Roundup Dry at 0.27 kg/acre:

Weeds controlled by Roundup at these rates plus:

Canada thistle rosettes**
Cow cockle
Dandelion
  (up to 6 inches)
Kochia
Volunteer canola
  (all varieties)

* Top growth control
** Suppression only
^ 1 to 3 leaf stage only
^^ Fall rosettes and spring seedlings
^^^ 2 to 10 leaf stage

Express may degrade if left in the sprayer for an extended period. Apply within 24 hours of mixing.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

Express Pack: $4.84 per acre.

Express TNG: $3.03 per acre plus cost of 2,4-D or glyphosate & adjuvant.

Application Information:

Water Volume:
Express Pack: Minimum 40 L per acre.
Express TNG + glyphosate: 22 to 40 L per acre

Pressure: 30 to 40 psi (210 to 275 kPa).

Nozzles: Flat fan nozzles with 50 mesh (or coarser) line strainers and screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply to wheat, durum or barley that are stressed by severe weather conditions (frost, drought or water saturated soil, extreme heat), as crop injury may result. Under certain conditions (such as heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures), lightening in crop colour and reduction in crop height may occur.

Tank Mixes:

Herbicides:

Do not use Express TNG alone.

In spring wheat (including durum) and barley:

Express TNG + 2,4-D Ester + Assert.
Express TNG + 2,4-D Ester + Banvel II
Express TNG + 2,4-D Ester + Puma(2) Super (green foxtail rate only)
Express TNG + MCPA Amine + Metribuzin (Sencor, Evict)

Prior to spring wheat (including durum) and barley:

Must be mixed with glyphosate (see Weeds Rates & Staging).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Express Pack label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check product labels for directions.

General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Grazing: Do not graze or feed to livestock within 7 days of application.

Preharvest: Do not harvest forage or cut for hay within 30 days of application.

Recropping: No restrictions the year after treatment. Canola, flax, lentils and alfalfa may be planted 2 months after an application of Express Pack.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. May be frozen.

Tank Cleaning:

Express can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray Express should be flushed out ***immediately*** after use. The manufacturer recommends that sprayers, including booms and nozzles, be flushed and emptied twice with a water-household ammonia rinse (1 L of 3% ammonia per 100 L of water) followed by a clean water rinse. All nozzles, screens and filters should be removed and cleaned with the above solution after applying this product. Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Hazard Rating:

Express (Pack)/Express TNG:
Warning Eye Irritant

Express Pack (2,4-D ester):

Warning Poison

FlaxMax

Herbicide Group – 1, 4
(Refer to page 27)

This product is a prepackaged tank mix of Poast Ultra (page 162) and FlaxMax (Curtail M page 86). Information listed is restricted to Crop, Weeds, Rates and Cost and other important details. For other detailed information on the component products see the product pages listed above.

Company:

BASF Canada

Formulation:

One case of FlaxMax contains 3 containers:

Poast Ultra (PCP#'25820): 450 g/L of sethoxydim formulated as an emulsifiable concentrate.
Container size - 3.65 L.

FlaxMax (PCP#25819): 50 g/L clopyralid and 280 g/L MCPA ester formulated as an emulsifiable concentrate.
Container size – 2 x 8.1 L.

Crops and Staging:

Flax – (NOT including Solin varieties) 2 to 6 inches (5 to 15 cm) in height.

Spraying at an early stage will reduce the risk of crop injury.

Weeds, Rates and Staging:

Poast Ultra at 0.13 L/acre, FlaxMax at 0.61 L/acre, and Merge at 0.3 L/acre (27 acres per case) will control the following weeds:

Annual Grasses 1 to 6 leaf stage	Annual Grasses 1 to 4 leaf stage	Annual Broadleaf Weeds 2 to 4 leaf stage	Perennial Weeds
Barnyard grass, crabgrass, fall panicum, Persian darnel, proso millet, volunteer corn, green and yellow foxtail, witch grass	wild oat, volunteer barley, volunteer wheat	flixweed, lamb's-quarters shepherd's-purse stinkweed, volunteer sunflower, wild mustard	Canada thistle*

Poast Ultra at 0.19 L/acre, FlaxMax at 0.81 L/acre, and Merge at 0.4 L/acre (20 acres per case) will control the weeds listed above plus:

Annual Grasses 1 to 6 leaf stage	Annual Grasses 1 to 4 leaf stage	Annual Broadleaf Weeds 2 to 4 leaf stage	Perennial Weeds
wild oat, volunteer barley, volunteer wheat		annual sow-thistle common groundsel, redroot pigweed, Russian pigweed, scentless chamomile, smartweed, tartary buckwheat, volunteer canola, wild buckwheat	dandelion**, perennial sow-thistle*** Canada thistle

* Apply from 10 cm until early bud stage. This treatment will provide season long control of light Canada thistle infestations with some regrowth in the fall.

** Apply at the spring rosette stage.

*** Top-growth control only.

Merge adjuvant (not included) must be used with FlaxMax. Mix to a final ratio of 0.75 to 1.0 L of Merge in 100 L of spray solution. (0.75% to 1.0% Merge volume per spray solution volume). When applying 40 L per acre of spray solution use one 8.1 L jug of Merge for every case of FlaxMax, regardless of whether the 20 or 27 acre per case rate FlaxMax is being used.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$19.28 to $26.03 per acre.

Application Information:

Water Volume: 40 L per acre.

Pressure: 35 to 40 psi (240 to 275 kPa).

Nozzles: Flat fan nozzles tilted forward at an angle of 45°.

Restrictions:

Rainfall: Within 6 hours may reduce control.

Grazing: Do not graze the treated crop or cut for livestock feed.

Preharvest Interval: Leave 60 days between spraying and harvest.

Recropping: FlaxMax residues in the soil may affect subsequent crops. Leave a minimum of 14 days between application of FlaxMax and replanting of barley, wheat, oat, and flax. Cultivating to a minimum depth of 4 inches (10 cm) is recommended before reseeding. Wheat, barley, oat, rye, corn, flax, canola or mustard may be planted the year after application. Do not seed any other crop until 2 years after application.

Aerial Application: Do not apply by air.

Storage: Store above 5°C. If frozen, bring to room temperature and agitate before use.

Hazard Rating:

◈ Warning Poison

Warning Skin and Eye Irritant

Frontier

Herbicide Group – 15
(Refer to page 27)

Company:

BASF Canada (PCP#23462)

Formulation:

900 g/L dimethanamid formulated as an emulsifiable concentrate. Container size - 10 L

Crops and Staging:

Pre-plant incorporated:
Corn (not sweet corn, popcorn, or corn grown for seed)
Dry beans (white and kidney beans only)

Pre-emergence surface:
Dry beans (white and kidney beans only).

Weeds and Staging:

Pre-emergent control of green foxtail.

Rates:

Pre-plant incorporated treatments:

Apply at 0.51 to 0.57 L/acre. Apply at the higher rate on fine-textured or high organic soils and for heavier anticipated weed problems.

Pre-emergence treatments:

SOIL TYPE	RATE (L/ACRE)		
	Less than 3% Organic Matter	3 to 6% Organic Matter	7 to 10% Organic Matter
Coarse Textured Soils	0.45	0.45	0.51
Medium and Fine Textured Soils	0.45	0.51	0.57

Cost (2004 suggested retail prices):

$18.36 to $23.25 per acre.

Application Information:

Water Volume: A minimum of 40 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan. Use 16 mesh suction screen, 50 mesh elsewhere on sprayer.

Incorporation: For pre-plant incorporated treatments, apply Frontier as a broadcast treatment and incorporate using a harrow, rolling cultivator or other implement capable of giving uniform, shallow incorporation into the top 5 cm (2 inches) of soil within 7 days of planting. Avoid deeper incorporation or reduced weed control and/or crop injury may result. Immediate incorporation after application is not necessary.

Beans must be planted at least 4 cm (1.5 inches) deep or crop injury may occur.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Rainfall is required within 7 to 10 days of application to activate and move Frontier into the soil zone. If dry conditions persist, a shallow cultivation or the use of a rotary hoe is necessary to move the herbicide into moist soil and control weed escapes. Shallow tillage is important to minimize dilution of the herbicide. If drought conditions persist after pre-plant incorporated or pre-emergence applications, weed control may not be adequate.

Tank Mixes:

Herbicides: None registered.

Fertilizers: May be applied with a liquid fertilizer carrier. Test compatibility of Frontier with liquid fertilizer by mixing a small amount of herbicide with a proportional quantity of liquid fertilizer in a jar. May also be impregnated on dry bulk fertilizers for pre-plant incorporated treatments. A minimum of 90 kg/acre of dry bulk fertilizer should be applied. Do not impregnate Frontier on nitrate fertilizers, superphosphates or limestone.

Insecticides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rainfall after application is important for good weed control.

Grazing: Do not graze or feed the treated corn crop within 40 days of application. Do not graze the treated bean crop or feed bean forage, hay or straw to livestock.

Recropping: Do not plant winter wheat within 120 days of application.

Aerial Application: Not registered.

Storage: Do not freeze. Must be stored under heated warehouse conditions.

Environment: Avoid drift onto non-target plants. Leave a buffer of 15 meters from water bodies and wetland areas.

Tank Cleaning:

The manufacturer does not provide information on tank cleaning. Generally, a mixture of water and household ammonia (1 L per 100 L of water) flushed twice through the tank and circulated through the lines and nozzles is an effective method of cleaning the sprayer tank.

Hazard Rating:

☠ Caution Poison.

Warning Eye Irritant and Potential Skin Sensitizer.

Frontline 2,4-D

Herbicide Group – 2, 4
(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

The Frontline 2,4-D package has 2 components:

Frontline 2,4-D A (PCP#'27242): 50 g/L florasulam formulated as a suspension concentrate

Frontline 2,4-D B (PCP#27243): 564 g/L 2,4-D LV ester formulated as an emulsifiable concentrate.

Container sizes – Frontline 2,4-D A - 1.6 L, Frontline 2,4-D B – 2 x 8 L.

Crops and Staging:

Spring wheat (including durum) in the 2 to 6 leaf stage.

When mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Bluebur
Burdock
Chickweed
Cleavers
Cocklebur
Dandelion*
Flixweed
Kochia**
Lady's-thumb
Lamb's-quarters
Mustards
Plantain
Prickly lettuce
Ragweed (common)
Redroot pigweed
Russian thistle
Shepherd's-purse
Smartweed
Sow-thistle (annual)
Stinkweed
Sunflower (annual)
Tartary buckwheat
Vetch
Volunteer canola***
Wild buckwheat
Wild mustard
Wild radish

Broadleaf weeds suppressed:

Canada thistle (top growth control only)
Hemp-nettle
Perennial sow-thistle (top growth control only)

* Seedlings and overwintered rosettes

** Up to 5 cm (2 inches) in height.

*** Including all herbicide-tolerant canola varieties

Cost (2004 suggested retail price):

$6.25 per acre.

Rate:

Frontline 2,4-D A: 40 mL/acre

Frontline 2,4-D B: 0.4 L (400 mL)/acre.

One case treats 40 acres.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 13 of the Guide to Crop Protection.

Application Information:

Water Volume: 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of Frontline 2,4-D. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum):
Assert 300 SC at 0.65 L/acre plus Acidulate
Everest at 17.4 g/acre plus recommended surfactant.
Puma(2) Super at 0.154 L/acre

Fertilizers: None registered.

Insecticides: None registered.

Note: The above mixes are those listed on the Frontline 2,4-D label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter. Do not harvest forage or cut hay within 30 days of application.

Pre-harvest Interval: Leave 60 days between application and harvesting mature crop.

Recropping: Wheat, barley, canola and peas may be grown the year following an application.

Aerial Application: Do not apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Environment: Leave a buffer zone of at least 30 m between the downwind edge of the boom and the closest edge of non-target areas such as shelterbelts, hedgerows and shrublands. A buffer zone of at least 15 m is required between the downwind edge of the boom and the closest edge of water or wetland areas such as ponds, streams, rivers, prairie potholes and sloughs.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◇ Warning Poison

May cause skin and eye irritation

Frontline

Herbicide Group – 2, 4

(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

The Frontline package contains 2 components:

Frontline A (PCP#27029): 50 g/L florasulam formulated as a suspension concentrate. Container size - 1.6 L

Frontline B (PCP#'27030): 500 g/L MCPA Ester formulated as an emulsifiable concentrate. Container size - 11.2 L.

Crops and Staging:

Spring wheat (including durum), barley and oats in the 2 to 6 leaf stage.

When tankmixing, always check the tankmix partner recommendations for additional staging restrictions.

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Ball Mustard
Burdock**
Chickweed
Cleavers
Flixweed
Hemp-nettleΔ
Kochia**
Lamb's-quarters
Prickly lettuce**
Ragweed (common)
Redroot pigweedΔ
Russian pigweed**
Shepherd's-purse
Smartweed
Stinkweed
Sunflower (annual)**
Volunteer canola (all varieties)*
Wild Mustard
Wild buckwheat

Broadleaf weeds suppressed:

Canada thistle Δ,†
Dandelion Δ, •
Plantain†
Stork's-bill Δ
Sow-thistle (annual)
Sow-thistle (perennial)†

* including all herbicide-tolerant canola varieties

** up to the 4 leaf stage of development

Δ for improved control of this weed add an additional 57 ml per acre of MCPA LV500 (Frontline B)

† top growth control

• seedlings and overwintered rosettes less than 15cm (6 inches)

Cost (2004 suggested retail price):

$7.40 per acre.

Rate:

Frontline A: 40 mL per acre

Frontline B: 280 mL per acre

One case treats 40 acres.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: A minimum of 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of Frontline. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley only:
Assert at 0.65 L/acre plus Acidulate.
Puma(2) Super at 155 mL/acre or 310 mL/acre.

In spring wheat (including durum) only:
Horizon at 95 mL/acre or 115 mL/acre plus Score adjuvant.
Everest at 17.4 g/acre plus Ag-Surf or Agral 90 adjuvant

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Frontline label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze treated crop or cut for feed within 7 days of application.

Preharvest Interval: Do not apply within 60 days of harvest.

Recropping: Wheat, barley, oats, canola and peas may be grown the year following an application.

Aerial Application: Do not apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Environment: Leave a buffer zone of at least 30 m between the downwind edge of the boom and the closest edge of non-target areas such as shelterbelts, hedgerows and shrublands. A buffer zone of at least 5 m is required between the downwind edge of the boom and the closest edge of water or wetland areas such as ponds, streams, rivers, prairie potholes and sloughs.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

May cause eye and skin irritation irritation

Fusion

Herbicide Group – 1

(Refer to page 27)

Company:

Bayer CropScience

Formulation:

One case of Fusion contains:

Component 1 (PCP#22845): 80.5 g/L fenoxaprop-p-ethyl. Container size - 3.7 L.

Component 2 (PCP#22850): 125 g/L fluazifop-p-butyl. Container size - 6.5 L.

Both components are formulated as emulsifiable concentrates.

Crops and Staging:

Crops are tolerant at all stages but the following preharvest restrictions apply:

CROP	STAGING
Canola	Do not apply beyond the 5 leaf stage, 80 day preharvest interval
Flax (including Solin)	Tolerant at all growth stages, 80 day preharvest interval
Field peas	Tolerant at all growth stages, 75 day preharvest interval
Lentils	Do not apply beyond the 6 node stage, 82 day preharvest interval
Mustard	Do not apply beyond the 5 leaf stage, 70 day preharvest interval

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Apply at the 1 to 6 leaf stage of the following annual grassy weeds. Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occur when weeds are controlled before tillering.

Barnyard grass
Green foxtail
Persian darnel
Volunteer barley
Volunteer wheat
Wild oats

Cost (2004 suggested retail price):

$11.95 per acre.

Rates:

Component 1: 0.19 L/acre.

Component 2: 0.32 L/acre.

One case treats 20 acres.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 23 to 45 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° stainless steel flat fan nozzles tilted forward at a 45° angle.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

If daytime temperatures before or after application are very hot and dry with low humidity, reduced control may result. Do not apply to crop that is stressed (frost, heat, drought, water saturated soil, disease, etc.) as crop injury may result.

Tank Mixes:

Herbicides:

Canola:

Lontrel (0.23 L/acre).

Muster (8 to 12 g/acre) (Argentine varieties only)

Flax (including Solin):

Buctril M (0.40 L/acre).

Flax (not including Solin):

MCPA ester or amine (up to 0.34 L/acre - 500 g/L formulations).

Under adverse conditions, tank mixes of Fusion and Buctril M or Fusion and MCPA may result in reduced grassy weed control.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: An interval of 4 days prior to, or 4 days after application of Fusion is required before any other pesticide can be applied (unless registered in a Fusion tank mix).

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Note: The above mixes are those listed on the Fusion label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: Within 2 hours may result in reduced control.
Grazing: Do not graze treated crops or cut for feed prior to crop maturity.

Recropping: If it is necessary to re-seed a crop within 60 days of applying Fusion herbicide, seed only the broadleaf crops listed on the Fusion label. If tank mixed with a broadleaf herbicide refer to both labels for recropping information. There are no recropping restrictions for Fusion herbicide applied alone the year after application.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environment: Do not apply within 15 m of non-target areas, water or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Caution Poison

Warning Eye and Skin Irritant

Experimental studies have shown that the active ingredient in Component 2 may cause birth defects in laboratory animals. Women capable of bearing children should avoid contact with this product.

glyphosate

Herbicide Group – 9
(Refer to page 27)

Company, Formulation and Cost (2004 suggested retail prices):

Product Name	Company	Salt*	Glyphosate acid content	Formulation	Container size**	Cost
Credit (PCP#25866)	Nufarm	IPA	356 g/L	Solution	1, 2, 3	$7.40 per L
Factor (PCP#27090)	IPCO	IPA	356 g/L	Solution	1, 2	$7.30 per L
Glyfos (PCP#24359)	Cheminova	IPA	360 g/L	Solution	1, 2, 3, 5	$8.95 per L
Maverick (PCP#26429)	Dow AgroSciences	IPA	356 g/L	Solution	1, 2	$7.99 per L
Renegade (PCP#20864)	Monsanto	IPA	356 g/L	Solution	1, 2	$7.99 per L
Roundup Dry (PCP#24874)	Monsanto	MA	68.5%	Soluble granule	7	$16.33 per kg
Roundup Original (PCP#13644)	Monsanto	IPA	356 g/L	Solution	1, 2, 4	$8.99 per L
Roundup Transorb (PCP#25344	Monsanto	IPA	360 g/L	Solution	1, 2, 3, 4, 6	$9.79 per L
Roundup Ultra (PCP#27764)	Monsanto	K+	540 g/L	Solution	1, 2, 3, 5	N/A
Roundup WeatherMax (PCP#27487	Monsanto	K^+	540 g/L	Solution	1, 2, 3, 5	$14.97 per L
Touchdown iQ (PCP#27912)	Syngenta	DA	360 g/L	Solution	1, 2, 3	$9.89 per L
Vantage (PCP#26172)	Dow AgroSciences	IPA	356 g/L	Solution	1, 2, 3	$8.95 per L
Vantage Plus (PCP#26171)	Dow AgroSciences	IPA	360 g/L	Solution	1, 2, 3	$9.79 per L

* Salt type: IPA = Isopropylamine, MA = Monoammonium, DA = Diammonium, K^+ = Potassium

** Container sizes available: 1) 2 x 10 L 2) 115 L 3) 450 L 4) 750 L 5) 1000 L 6) 1200 L 7) 10.4 kg

Some products may be more effective (not related to higher glyphosate content) under adverse conditions, but that benefit is reduced when applications are made under optimal conditions for activity (i.e. rapid weed growth). When selecting a glyphosate product, consult the product manufacturer or ask your retailer about differences in the level of post-sales support provided for different products.

Crops and Uses:

1. Annual weed control prior to crop emergence or in summerfallow.
2. Quackgrass control prior to seeding or after harvest.
3. Dandelion control (other than preharvest).
4. Canada thistle control in summer-fallow, shelterbelts and post-harvest.
5. Alfalfa control (other than preharvest).
6. Other perennial weeds control in summerfallow, shelterbelts and post-harvest.
7. Patch treatments of perennial weeds in cereals, corn, soybean and forages.
8. Preharvest perennial weed control.
9. For use in Roundup Ready (glyphosate tolerant) crops.
10. Tank Mixes.

1. Annual weed control prior to crop emergence or in summer-fallow:

Weeds, Rates and Staging:

PRODUCTS	RATE PER ACRE	SURFACTANT *	WEEDS CONTROLLED	WEED STAGE
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	0.30 L	0.14 L/ acre	**Grasses:** Green foxtail, volunteer cereals, wild oat (light infestations) **Broadleaves:** lady's- thumb, stinkweed, volunteer canola (NOT including Roundup Ready varieties), wild mustard	Less than 3 inches (8 cm) high. Apply at the 1 to 3 leaf stage of wild oat.
Roundup Dry	0.16 kg			
Roundup Ultra, Roundup WeatherMax	0.20 L			
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	0.40 L	0.14 L/ acre	Above weeds plus: **Grasses:** heavy infestations of wild oat **Broadleaves:** suppression of flixweed, kochia	1 to 3 leaves for wild oat. Weeds 3 to 6 inches (8 to 15 cm).
Roundup Dry	0.21 kg			
Roundup Ultra, Roundup WeatherMax	0.27 L			
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage,Vantage Plus	0.51 to 0.77 L	Not required	Above weeds plus: **Grasses:** downy brome, Persian darnel, **Broadleaves:** Canada fleabane, cleavers**, common ragweed, flixweed,hemp-nettle, lamb's-quarters,narrow-leaved hawk's-beard, redroot pigweed, Russian thistle, volunteer flax, wild buckwheat	Canada fleabane, common ragweed less than 3 inches(8 cm) high. Other weeds less than 6 inches (15 cm). Use high rate for narrow-leaved hawk's-beard 3 to 6 inches (8- 15 cm) or wild buckwheat at the 3-4 leaf stage.
Roundup Dry	0.27 to 0.40 kg			
Roundup Ultra, Roundup WeatherMax	0.33 to 0.51 L			
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	0.91	Not required	Above weeds plus: **Grasses:** annual blue grass, crabgrass, **Broadleaves:** annual sow-thistle, kochia, prickly lettuce, shepherd's purse	Less than 6 inches (15 cm) high
Roundup Dry	0.48 kg			
Roundup Ultra, Roundup WeatherMax	0.61 L			
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.42 L	Not required	Above weeds	Greater than 6 inches (15 cm) high
Roundup Dry	0.74 kg			
Roundup Ultra, Roundup WeatherMax	0.94 L			

* Unless otherwise specified, use one of the following surfactants: Agral 90, Agsurf, Companion, Enhance, Frigate or LI 700.

** Registered for control with Roundup Original, Roundup Transorb, Vantage and Vantage Plus only.

2. Quackgrass control prior to seeding or after harvest:

PRODUCT	RATE PER ACRE	QUACK GRASS STAGE
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 L	Season long control of light to moderate infestations. Apply when quack grass is 8 inches (20 cm) tall and has 3 to 4 actively growing leaves. Apply spring or fall.
Roundup Dry	0.53 kg	
Roundup Ultra, Roundup WeatherMax	0.67 L	
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 to 2.8 L	Apply when quack grass has 3-4 new leaves for long term control of heavy infestations. Use high rate for sod-bound quack grass (left undisturbed for at least 2 years).
Roundup Dry	0.53 to 1.5 kg	
Roundup Ultra, Roundup, WeatherMax	0.67 to 1.89 L	

Do not apply fall treatments if a hard frost has occurred (-5°C) or if plants are drought stressed. Spread straw to allow regrowth and good spray coverage.

Cultivation prior to application will result in reduced control. Do not cultivate between harvest and treatment when using fall applications. If using spring applications on fields which have been fall-tilled, delay application until the quack grass has reached the 4 to 5 leave stage. (This will occur 1 to 4 weeks later on fall-tilled fields than in undisturbed fields).

Cultivation after application usually will improve control of quack grass. Wait a minimum of 3 days after application before cultivating. If growing conditions are poor (cold or dry), particularly in the fall, waiting longer than 5 days may improve control.

3. Dandelion control (other than preharvest):

Apply up to and including dandelion bloom for best results.

PRODUCT	**Less than 6 inches (15 cm) diameter. Allow 3 or more days after treatment before tillage.**	**Greater than 6 inches (15 cm) diameter. Use higher rate when infestation is heavy.**	**Prior to seeding or after harvest. Use higher rate for weeds beyond 3 inches (8 cm) or for heavy infestation. Allow 7 or more days after treatment before tillage.**
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Vantage, Vantage Plus	1.0 L/acre	1.5 to 2.0 L/acre	–
Roundup Dry	0.53 kg/acre	0.80 to 1.06 kg/acre	–
Touchdown iQ	1.0 L/acre	1.5 to 2.0 L/acre	1.82 to 2.83 L/acre
Roundup Ultra, Roundup WeatherMax	0.67 L/acre	1.0 to 1.34 L/acre	–

4. Canada thistle control in summer-fallow, shelterbelts and post-harvest:

PRODUCT	RATE PER ACRE	WEED STAGING
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.9 to 2.8 L	Bud stage or beyond. Allow at least 5 days after application before tillage.
Roundup Dry	1 to 1.5 kg	
Roundup Ultra, Roundup WeatherMax	1.28 to 1.89 L	
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 L	Rosettes at least 6 inches (15 cm) in diameter, treated in late summer, following tillage in spring and early summer (up to August 1). Allow thistles to regrow for 5 weeks following last tillage. Wait a minimum of 10 days after application before tillage. Treatment after a mild frost is possible if leaves are green and pliable and plants are actively growing.
Roundup Dry	0.53 kg	
Roundup Ultra, Roundup WeatherMax	0.67 L	
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.9-2.8 L	Post-harvest stubble treatment. Allow 8-10 inches (20-25 cm) of new growth before application. must be Plants sprayed at least 2 weeks prior to killing frost. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage.
Roundup Dry	1.0-1.5 kg	
Roundup Ultra, Roundup WeatherMax	1.28 to 1.89 L	

5. Alfalfa Control (other than preharvest):

PRODUCT	RATE PER ACRE	WEED STAGING
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.5 to 2.0 L	Fall control of alfalfa in early bud to full bloom stage. Use high rate when alfalfa populations are high or when perennial grasses are present. Allow at least 5 days before tillage. See tank mix section for minimum tillage or spring applications. Apply with 23-135 L/acre water.
Roundup Dry	0.8 to 1.1 kg	
Roundup Ultra, Roundup WeatherMax	1.0 to 1.34 L	
Touchdown iQ	0.91 to 1.82 L	Prior to seeding or after harvest. Use higher rate for weeds beyond 3 inches (8 cm) in height or for heavy weed infestations. Wait 7 days after application for tillage. Apply in 23-135 L/acre water.

6. Other perennial weed control in summerfallow, shelterbelts and post-harvest:

(Refer to individual product labels for detailed application information.)

	RATES IN UNITS PER ACRE		
	Foxtail Barley Seedling to heading*	**Toadflax**	**Other Perennial weeds****
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 to 2.0 L	1.0 L (vegetative stage in summer fallow)	4.9 L
Glyfos	0.4 L (suppression)		
Roundup Dry	0.53 to 1.1 kg	0.53 kg (vegetative stage in summer fallow)	2.5 kg
Roundup Ultra, Roundup WeatherMax	0.67 to 1.34 L/acre	0.67 L (vegetative stage in summer fallow)	1.89 to 3.24 L

* Late fall applications may provide better control of established foxtail barley plants than spring applications.

** Perennial weeds such as absinthe, blue grass spp., smooth brome grass, cattail, curled dock, field bindweed (bloom stage or beyond), hemp dogbane, hoary cress, poison ivy, purple loosestrife, perennial sow-thistle, and yellow nutsedge applied at the early heading to early bud stage.

7. Patch treatments of perennial weeds in wheat, oat, barley, corn, soybean, forage legumes and forage grasses:

(refer to product label for detailed application instructions)

	RATES PER ACRE				
	Quack grass 8 in (20 cm) tall	**Canada thistle Bud or beyond**	**Milkweed Bud to bloom**	**Other perennial weeds***	**Spot treatment rates for hand held equipment**
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 to 2.8 L	1.9 to 2.8 L	4.9 L	2.8 to 4.9 L	1 to 2 L per 100 L water**
Roundup Dry	0.53 to 1.5 kg	1.0 to 1.5 kg	2.5 kg	1.5 to 2.5 kg	0.5- to 1.0 kg per 100 L water**
Roundup Ultra, Roundup WeatherMax	0.67 to 1.89 L	1.28 to 1.89 L	3.24 L	1.89 to 3.24 L	0.67 to 1.34 L per 100 L water**

* Perennial weeds such as absinthe, blue grass spp., smooth brome grass, cattail, curled dock, field bindweed (bloom stage or beyond), hemp dogbane, hoary cress, poison ivy, purple loosestrife, perennial sow-thistle, and yellow nutsedge applied at the early heading to early bud stage.

** Use the low rate for quack grass and the high rate for all other perennials.

8. Preharvest perennial weed control:

Crops registered and rates (per acre):

Product	Barley, Bean (dry), Canola, Field pea, Flax (including Solin) Lentil, Soybean Wheat.	Oat	Forages
Credit, Factor, Maverick, Renegade, Roundup Original, Roundup Transorb, Touchdown iQ, Vantage, Vantage Plus	1.0 L	1.0 L	1.0-2.0 L
Roundup Dry	0.53 kg	0.53 kg	0.53-1.06 kg
Roundup Ultra, Roundup WeatherMax	0.67 L	0.67 L	0.67 to 1.34 L
Glyfos	1.0 L	1.0 L	Not registered

Do not apply to any crops grown for seed. May be applied to barley grown for malt. Contact malt barley buyers prior to application to confirm acceptance of glyphosate-treated barley. May be applied to tame oat. Contact tame oat buyers prior to application to confirm acceptance of glyphosate treated oat.

Weeds Controlled with preharvest applications:

Product	Quack grass 4-5 green leaves	Canada thistle and perennial sow-thistle at bud stage or beyond	Common milkweed at bud to bloom stage	Toadflax at bud to full bloom stage	Dandelion from rosette to full bloomstage
Credit, Factor, Glyfos, Maverick, Renegade, Roundup Dry, Roundup Original, Roundup Transorb, Roundup Ultra, Roundup WeatherMax Touchdown iQ, Vantage, Vantage Plus	X	X	X	X	X

Crop Staging for preharvest applications:

Apply to crops (except forage) when grain moisture is less than 30%. The following chart lists visual symptoms that can be used as guidelines to when 30% grain moisture has been reached.

CROP*	VISUAL SYMPTOMS
Wheat, Barley, Oat	Hard dough stage – a thumbnail impression remains on seed.
Canola	Pods are green to yellow and most seeds are yellow to brown.
Flax (including Solin)	Majority (75 to 80% of bolls) are brown.
Lentil	Lowermost pods (bottom 15%) are brown and rattle when shaken.
Pea	Majority (75 to 80%) of pods are brown.
Soybean	Stems are green to brown in colour and pod tissue is brown and dry in appearance (80 to 90% leaf drop).
Dry Bean	Stems are green to brown in colour and pods are mature (yellow to brown) and 80 to 90% of the original leaves have dropped.
Forage	3 to 7 days prior to the last cut before rotation or forage renovation. Do not apply to forage stands that are to be maintained.

*See the crops section for a guide to the products registered on each crop.

9. For Use in Roundup Ready (glyphosate tolerant) Crops including:

Roundup Ready Canola: (Factor, Glyfos, Roundup Original, Roundup Transorb, Roundup Ultra, Roundup WeatherMax, Touchdown iQ, Vantage, Vantage Plus only)

Roundup Ready Field corn: (Factor, Roundup Original, Roundup Transorb, Roundup Ultra, Roundup WeatherMax, Vantage and Vantage Plus only)

Roundup Ready Soybean: (Factor, Glyfos, Roundup Original, Roundup Transorb, Roundup Ultra, Roundup WeatherMax, Touchdown iQ, Vantage, Vantage Plus only)

Weeds, Staging and Rates in Roundup Ready Canola:

All applications must be made within the cotyledon to 6 leaf stage of RR canola.
Temporary yellowing may occur if applied at the 4 to 6 leaf stage of the crop.

PRODUCT	RATE (L/ACRE)	WEEDS CONTROLLED
Factor Glyfos Renegade Roundup Original Roundup Transorb Touchdown iQ Vantage Vantage Plus Roundup Ultra† Roundup WeatherMax† (rates given in brackets)	0.33-0.50 (0.22 to 0.34)†	**Annual grasses:** barnyard grassΔ***, green foxtail, volunteer cereals, wild oat **Annual broadleaves:** annual smartweed spp.**, chickweed, corn spurry, cow cockle*, hemp-nettle, kochia, lamb's-quarters, night-flowering catchfly*, redroot pigweed, Russian thistle, shepherd's-purse*, stinkweed, volunteer canola (except Roundup Ready varieties), wild mustard, wild tomato.
	0.50 (0.34)†	**Annual broadleaves:** cleavers, flixweedΔ, wild buckwheat, stork's-billΔ, narrow-leaved hawk's-beardΔ **Perennial weeds suppressed:** Canada thistle, dandelion, perennial sow-thistle, and season long quack grass control.
	2nd application at 0.5 L (0.34 L)† per acre	**Annual broadleaves:** round-leaved mallowΔ **Season long control of following perennials:** Canada thistle, foxtail barleyΔ, and perennial sow-thistle.
Factor, Glyfos, Renegade, Roundup Original, Roundup Transorb, Roundup WeatherMax† (rates given in brackets) Touchdown iQ	0.75 (0.51 L)†	**Season long control of following perennials:** Canada thistle, foxtail barley, and perennial sow-thistle.

† Rates in brackets for Roundup WeatherMax and Roundup Ultra only.
* Low rates can be used only up to the 3 leaf stage of the crop otherwise use high rate.
** Low rates can be used only when annual smartweed is in 4 - 6 leaf stage.
*** Not registered for control with Roundup Ultra.
Δ Not registered for control with Vantage or Vantage Plus.

NOTE: A maximum of 1.0 L/acre (0.67 L/acre for Ultra and WeatherMax) per season is allowed in Roundup Ready Canola.

Weeds, Staging and Rates in Roundup Ready Corn and Roundup Ready Soybean:

All applications must be made within the following crop growth stages.

PRODUCT	RATE (L/ACRE)	WEEDS CONTROLLED
Factor, Glyfos[***] Renegade Roundup Original Roundup Transorb Touchdown iQ[***] Vantage Vantage Plus Round Ultra[†] Roundup WeatherMax[†] (rates given in brackets)	1.0 (0.67)[†] Apply when weeds are less than 10 inches or 25cm high to maximise weed control and crop yield.	**Grasses:** barnyard grass crabgrass spp. foxtail (green, yellow, giant) proso millet[Δ] quack grass Volunteer barley and wheat[ΔΔ] wild oats[ΔΔ] **Broadleaves:** chickweed[ΔΔ] cleavers[ΔΔ] corn spurry[ΔΔ] cocklebur cow cockle[ΔΔ] common milkweed (suppression only)[Δ] common ragweed eastern-black flowering night-shade flixweed[ΔΔ] hemp-nettle[ΔΔ] kochia[ΔΔ] lamb's-quarters narrow-leaved hawk's-beard[Δ] night-flowering catchfly[ΔΔ] pigweed (smooth, redroot) round-leaved mallow[ΔΔ] Russian thistle[ΔΔ] shepherd's-purse[ΔΔ] smartweed spp. stinkweed[ΔΔ] stork's-bill[ΔΔ] velvetleaf volunteer canola (except Roundup Ready varieties)[ΔΔ] wild mustard wild buckwheat wild tomato[ΔΔ] yellow nutsedge (suppression)[Δ]
	2nd application at 1.0 L (0.67L)[†] per acre. Late flushes of the above weeds.	**Late flushes of heavy infestations of the above weeds plus control of:** Canada thistle[Δ], common milkweed, perennial sow-thistle[Δ], yellow nutsedge

Additional Weeds, Staging and Rates in Roundup Ready Soybean only:

PRODUCT	RATE (L/ACRE)	WEEDS CONTROLLED
Factor, Glyfos Roundup Original Roundup Transorb Round Ultra[†] Roundup WeatherMax[†] (rates given in brackets) Touchdown iQ	one application of 2.0 L (1.35 L)[†] per acre	**Heavy infestations of the annual weeds listed above plus control of:** Field Bindweed, Common Milkweed, Perennial sow-thistle, Canada thistle, and yellow nutsedge

[†] Rates in brackets for Roundup WeatherMax and Roundup Ultra only.

[***] Not registered for use in Roundup Ready Corn.

[Δ] Registered for control with Vantage or Vantage Plus in Roundup Ready corn only.

[ΔΔ] Registered only with Factor, Renegade, Roundup Original, Roundup Transorb, Roundup Ultra and Roundup WeatherMax only in RR corn.

Tank mixes in Roundup Ready Crops:

Canola: Factor, Glyfos, Roundup Original, Roundup Transorb, Roundup Weathermax, Touchdown iQ, Vantage, and Vantage Plus only with Lontrel at 112 mL/acre.

Soybean: Factor, Glyfos, Roundup Original and Roundup Transorb, Roundup Weathermax only with Assure II at 154 mL/acre for volunteer RR corn control.

Corn: Roundup Transorb, and Roundup Weathermax only with Atrazine at 0.63 to 0.84 L/acre of 480 to 500 g/L formulations or 0.34 to 0.45 kg/acre of 90% dry formulations for residual control of annual broadleaf weeds.

10. Tank Mixes:

	RATE PER ACRE	Factor	Renegade, Roundup Original, Roundup Transorb, Roundup, WeatherMax†	Credit, Glyfos, Maverick, Vantage, Vantage Plus	Touchdown iQ	Roundup Dry	Roundup Ultra
Preseeding cereals••:	**2,4-DΔ (0.23 to 0.58L)***	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.5 to 0.8 L	0.27 to 0.4 kg	0.34-0.51 L
	Express Toss n Go Agral 90 (4 g/ac + 0.35 v/v)	Not registered	Not registered	Not registered	0.3 to 0.4 L	Not registered	Not registered
	Pardner(0.51 L)	0.3 to 0.4 L	0.3 to 0.4 L*** †(0.2 to 0.27 L)	0.3 to 0.4 L	0.3 to 0.4 L	0.16 to 0.21 kg	0.2 to 0.27 L
	MCPA (0.2 to 0.4 L)*	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.5 to 0.8 L	0.27 to 0.4 kg	Not registered
	Buctril M (0.2 to 0.4 L)*	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.3 to 0.4 L	0.27 to 0.4 kg	Not registered
Preseeding corn & flax:	**MCPAΔΔ (0.2 to 0.4 L)***	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.5 to 0.8 L	0.27 to 0.4 kg	Not registered
	Buctril M (0.2 to 0.4 L)*	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.3 to 0.4 L	0.27 to 0.4 kg	Not registered
Preseeding lentil & chickpea:	**MCPA AmineΔΔ (0.2 to 0.28 L)***	Not registered	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.5 to 0.8 L	0.27 to 0.4 kg	Not registered
Preseeding field pea:	**MCPA AmineΔΔ (0.2 to 0.28 L)***	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.5 to 0.8 L	0.27 to 0.4 kg	Not registered
Preseeding canaryseed & seedling forage grasses•:	**Buctril M (0.2 to 0.4 L)***	0.5 to 0.8 L	0.5 to 0.8 L †(0.34 to 0.51 L)	Not registered	0.3 to 0.4 L	0.27 to 0.4 kg	Not registered
Chem fallow:	**2,4-DΔ (0.5 L)***	0.3 to 0.4 L	0.3 to 0.4 L †(0.2 to 0.27 L)	0.3 to 0.4 L	0.3 to 0.4 L	0.16 to 0.21 kg	0.2 to 0.27 L
	Dicamba (0.12 L)*	0.3 to 0.4 L	0.3 to 0.4 L †(0.2 to 0.27 L)	0.3 to 0.4 L	0.3 to 0.4 L	0.16 to 0.21 kg	0.2 to 0.27 L
	Pardner (0.51 L)	0.3 to 0.4 L	0.3 to 0.4 L*** †(0.2 to 0.27 L)	0.3 to 0.4 L	0.3 to 0.4 L	0.16 to 0.21 kg	0.2 to 0.27 L
Canada thistle control in stubble or fallow:	**Dicamba (0.51 L)****	0.69 L	0.69 L †(0.46L)	0.69 L	0.69 L	0.36 kg	0.46 L
Alfalfa control in spring / fall:	**2,4-DΔ (0.50 to 1.0 L)***	1.0 to 2.0 L	1.0 to 2.0 L †(0.67 to 1.34 L)	1.0 to 2.0 L	1.0 to 2.0 L	0.53 to 1.1 kg	0.67 to 1.34 L

* Roundup Ready (RR) canola control: Tank mixes of 2,4-D* at 0.23 to 0.34 L/acre, MCPA and Buctril M will control volunteer RR canola up to the 4 leaf stage and 2,4-D at 0.45 to 0.68 L/acre will give control up to the 6 leaf stage. Earlier application will result in more consistent control. Dicamba at 0.12 L/acre will not control RR canola.

** See recropping restrictions for Dicamba with fall applications.

*** Roundup Transorb registered for 0.4 L/acre rate of glyphosate only with Pardner tankmixes.

Δ Rates based on 500 g/L formulations. All formulations are registered for tankmixes.

ΔΔ Use only amine formulations of MCPA prior to corn lentil, chickpea and field peas

• Forage grasses include brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, Timothy, Orchard grass, creeping red fescue, meadow fescue, meadow foxtail, tall fescue, meadow bromegrass, streambank wheatgrass and reed canary grass.

•• 2,4-D tankmixes in cereals are registered for winter wheat, wheat, barley and rye; Pardner tank-mixes in cereals are registered on wheat oats and barley; Buctril M and MCPA tank-mixes registered on cereals include wheat, barley oat and rye; and Express Toss n Go tank-mixes include wheat, durum and barley.

† WeatherMax rates given in brackets.

Note: The above mixes are those listed on the glyphosate labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. See the general guidelines for mixing pesticides for more information.

Application Information:

Water Volume: Do not use 'hard' or dirty water or weed control may be reduced.

Using clean water at 23-45 L per acre will provide acceptable weed control in most situations; use of the lower volume will improve control.

Refer to specific weed control situations or labels for more information on water volumes and adjuvants.

Pressure: 30 to 40 psi (200 to 275 kPa)

Nozzles: Flat fan

How it Works:

Refer to table 2 on page 29.

Effects of Growing Conditions:

Best results are achieved when temperatures are near 20°C and when weeds are actively growing. Frost which kills more than 40% of the above ground tissue will reduce control. Control will also be reduced if foliage is heavily covered with dust.

Restrictions:

Rainfall: Do not apply if rainfall is forecast for the time of application, as weed control may be reduced. Consult manufacturer for more information.

Grazing Interval: Only grain and forages treated with Touchdown iQ may be fed to livestock. All portions of forage and crops treated with other products may be fed to livestock.

Recropping Interval: No restrictions.

Aerial application: Roundup Original, Roundup Transorb Vantage and Vantage Plus, are now registered for aerial application of preharvest treatments only to all registered preharvest crops (not grown for seed) excluding forages. Do not apply any other brands of glyphosate by air. Consult manufacturer for current aerial preharvest registration status.

Storage: May be stored below 0° C.

Equipment: Do not mix, store or apply this product in galvanized steel or unlined steel (except stainless steel) containers or spraytanks.

Environment: Glyphosate is very toxic to non-target plants.

Gound applications - maintain a 15 m buffer zone from the downwind edge of the spray boom to non-target areas to minimize drift damage.

Preharvest aerial applications - maintain a 25 m buffer zone from water and wetland areas, and a 55 m buffer from shelterbelts, woodlots and other cover on the edge of treated fields.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

▽ Caution Poison (Roundup WeatherMax)

Warning Eye and Skin Irritant (Roundup Dry, Roundup WeatherMax)

Danger Eye and Skin Irritant (Roundup Ultra)

Danger Corrosive to eyes (Roundup Dry)

Caution Skin and Eye Irritant (all other products)

Gramoxone

Herbicide Group – 22
(Refer to page 27)

Company:

Syngenta (PCP#8661)

Formulation:

200 g/L paraquat. Container size - 5 L, 20 L.

Crops and Staging:

Stale seedbed - Non-selective weed control applied 3 days prior to crop emergence in beans, corn, potatoes, peas, soybeans.

Non-selective weed control applied as a directed spray between rows in row crops. Avoid contact with crop foliage

Control of weed seedlings in established alfalfa and bird's-foot trefoil for hay - apply 5 days after first cut.

Control of weed seedlings in bird's-foot trefoil for seed - apply in spring when bird's-foot trefoil shoots are 3 to 6 inches (7.5 to 15 cm) long

Non-selective weed control applied as a directed spray in and around shelterbelt trees or woodlot plantings. Avoid contact with foliage

Weeds and Staging:

Annual weed burn-off. Best control when weeds are less than 2 inches (5 cm) in height or diameter.

Cost (2004 suggested retail price):

$25.55 to $51.10 per acre.

Rates:

If weeds are less than 2 inches (5 cm) in height, apply 1.1 L per acre (5 L treats 4.5 acres). If weeds are taller than 2 inches (5 cm), increase the rate of Gramoxone to 2.2 L per acre (5 L treats 2.2 acres).

Application Information:

Water Volumes: 135 to 500 L per acre. Use the higher water volumes within the range if weed growth is dense. Good coverage is critical for good control.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Best results on cloudy days or just prior to darkness.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 1 hour will reduce weed control.

Grazing: Do not graze or harvest treated foliage. Regrowth from treated alfalfa or bird's-foot trefoil may be fed to livestock.

Recropping Interval: No restrictions

Aerial Application: No restrictions on label, however, since this product requires a minimum of 30 gallons per acre of water, applications by air are impractical.

Storage: Do not freeze.

Environment: Maintain a buffer zone of 15 m between the downwind edge of the spray boom and non-target plants and 45 m for wetland areas. Do not spray when conditions are dead clam or when wind is gusty or blowing faster than 16 km/hr.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger - Poison (may be fatal if swallowed)

Danger – Corrosive to eyes

Re-entry Period: Do not re-enter treated fields for 24 hours following application. If necessary, workers may re-enter field after 4 hours if wearing protective clothing. See label for details.

Grazon

Herbicide Group – 4
(Refer to page 27)

Company:

Dow AgroSciences (PCP#26649)

Formulation:

65 g/L picloram + 240 g/L 2,4-D formulated as an emulsifiable concentrate.

Container size – 2 x 10 L.

Note: Available only through selected retail outlets.

Crops and Staging:

Permanent grass pasture and rangeland. Apply in spring or early summer.

Weeds, Rates and Staging:

Broadleaf weeds:

Apply at 1.5 L/acre:

Canada thistle
Dandelion
Common Yarrow

Apply at 2.8 L per acre:

Burdock
Clovers (red, sweet)
Common ragweed
Dock
Fleabane
Goldenrod
Plantain
Prickly lettuce
Vetch
Wild carrot

Woody plants:

Apply at 7.3 to 10.1 L per acre by ground and 10.1 to 14.2 L per acre by air:

Alder
AspenΔ
Birch
Cedar (red cedar*)
CherryΔ
LocustΔ
Maple*
Oak*Δ
Pine*
Poplar
Spruce*
Sumac Δ
WillowΔ

Δ Aerial application only

* High rate by air when species growing on heavy clay soils.

Use an approved drift control additive or system.

Cost (2004 suggested retail price):

$14.25 per L.

Application Information:

Water volume:

Ground application: 80 L per acre.
Aerial application: 60 L per acre

Pressure: Use pressures that will give the lowest amount of drift while maintaining good pattern and coverage.

Nozzles: Use an approved drift control system that produces coarse droplets in a uniform pattern.

Drift of even small amounts of Grazon into sensitive plants or areas where sensitive crops may be grown can cause injury. Do not apply under conditions prone to drift (i.e. high winds and temperature inversions).

How it Works:

Refer to table 2 on page 29.

IMPORTANT: Picloram is a very persistent and water-soluble herbicide. Treated soil should not be moved from the treated area. Do not apply to soils that are permeable, have sinkholes, or lie over limestone bedrock. Do not apply to soils whose surfaces are composed of fractured rock or unconsolidated gravel. Application to these sites may allow the movement of herbicide to underlying water sources or aquifers. When applying Grazon over sandy soils ensure that aquifers are not within 1.8 m of the soil surface. If shallow aquifers are present, do not apply Grazon. Grazon must not be applied on range and pasture acres that are irrigated. Do no compost or mulch clippings or manure from grass treated with Grazon unless being reapplied to the treated area.

Effects of Growing Conditions:

Nothing listed on label.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Do not apply if rainfall is forecast.

Grazing: Do not allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter. Do not harvest forage or cut hay within 30 days of application. **See restrictions in "How it Works" section above.**

Recropping: Legumes may not be established in a pasture for several years after treatment. If legumes are essential in a pasture, do not use Grazon. Do not break up treated pasture and plant to sensitive broadleaf crops for at least 5 years after application of Grazon.

Storage: Store product in original containers in a secure, dry, cool area. Do not freeze.

Environment: This product is moderately toxic to fish. Apply this product well back from any water bodies to avoid contamination. Heavy rains can move this product from its application site down slope toward sensitive areas. Do not load or mix near wells, dugouts or other water bodies. Contact the provincial environment department for permits to apply near water.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

May cause skin and eye irritation

Harmony Total

Herbicide Group – 1, 2
(Refer to page 27)

This product is a prepackaged tank mix of Refine Extra (page 176) and Horizon (page 135). Information listed is restricted to Crop, Weeds, Rates, Cost and Tank mixes. For other detailed information on the component products see the product pages listed above.

Company:

E. I. duPont Canada

Formulation:

Refine Extra (PCP#22575): 50% thifensulfuron methyl + 25% tribenuron methyl; formulated as a water dispersible granule.
Container size: 4 x 80 g water soluble pouches (320 g total).

Horizon WO (PCP#25477): 240g/ L clodinafop-propargyl formulated as an emulsifiable concentrate.
Container size: 1 x 3.68 L

Score adjuvant (PCP#28476): Container size - 2 x 6.4 L

Crops and Staging:

Spring wheat (including durum) Prior to the emergence of the 4th tiller.

Weeds and Staging:

Grass Weeds:

Wild oat -1 to 6 leaves up to the emergence of the 4th tiller.
Green foxtail -1 to 5 leaves up to the emergence of the 3rd tiller.

Broadleaf Weeds:

Unless otherwise noted below, apply to young and actively growing weeds that are less than 4 inches (10 cm) in height or width.

Controlled:

Ball mustard
Chickweed (1 to 6 leaf)
Common groundsel
Corn spurry
Cow cockle
Flixweed
Green smartweed
Hemp-nettle
Kochia
Lady's-thumb
Lamb's-quarters
Narrow-leaved hawk's beard
Redroot pigweed
Russian thistle
Shepherd's-purse
Stinkweed
Tartary buckwheat
Volunteer canola
 (CLEARFIELD canola only with MCPA Ester tank mix)
Volunteer sunflowers
Wild buckwheat
 (up to 3 leaf)
Wild mustard

Suppressed:
Cleavers - 1 to 3 whorls
Stork's-bill, round-leaved mallow - 2 to 6 leaves
Canada thistle, perennial sow-thistle - under 6" (15 cm) tall, prior to budding
Scentless chamomile
Toadflax - less than 6" (15 cm) tall

For optimum control apply before the annual grasses tiller. Harmony Total will also control or suppress the broadleaf weeds listed for Refine Extra alone at 8 g/acre.

Cost (2004 suggested retail price):

$23.11 per acre.

Rates:

Refine Extra - 8 g/acre
Horizon WO - 93 mL/acre
Score adjuvant - 0.8 L per 100 L of spray solution
One case treats 40 acres (16 ha)

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kpa).

Nozzles: Flat fan nozzles with 50 mesh line strainers and screens.

Tank Mixes:

Do not tank mix with substances that contain boron or that release chlorine.

Herbicides:
MCPA Ester (0.38 to 0.45 L/acre of 500 g/L formulation).
Banvel II (44.5 mL/acre). Not for use in durum.

Hoe-Grass 284

Herbicide Group – 1
(Refer to page 27)

Company:

Bayer CropScience (PCP#18042)

Formulation:

284 g/L diclofop-methyl formulated as an emulsifiable concentrate. Container size - 10 L.

Crops, Rates and Staging:

Unless specified, there are no leaf stage restrictions but do not apply within 60 days of harvest.

Apply 1.42 L/acre (7 acres per jug) to the following crops:
Dry beans (white, black, pinto)
Fababeans
Potatoes
Soybeans

Apply 1.01 to 1.13 L per acre (7.75 to 10 acres per jug) to the following crops:
Barley (1 to 4 leaves, before tillering)
Flax (NOT including Solin varieties)
Lentils
Mustard
Peas (field and processing)
Rapeseed (canola)
Rye (spring and fall)
Sunflowers (except Corona)
Tame buckwheat
Triticale
Wheat (spring, durum and winter)

Seedling Forage grasses:
bromegrass, creeping red fescue, wheatgrass (crested, intermediate) , Russian wild ryegrass

Seedling legumes (forage or seed production):
alfalfa, red clover and sweet clover

Seedling legumes (seed production only):
Sainfoin and alsike clover

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Wild oats, green and yellow foxtail and barnyard grass - 1 to 4 leaf stage. Although registered for application up to the 5 leaf stage of wild oats, results at this stage are inconsistent and this use is not recommended. For best results, apply when the majority of weeds are in the 2 to 3 leaf stage.

Persian darnel - 1 to 3 leaf stage.

Volunteer corn - 6 to 10 inches (15 to 25 cm) in height.

Cost (2004 suggested retail price):

$13.43 to $19.18 per acre.

Application Information:

Water Volume:

Ground equipment: 45 L per acre.
Aerial equipment: A minimum of 14 L per acre.

Pressure:

Ground equipment: 40 psi (275 kPa).
Aircraft: 45 psi (300 kPa).

Nozzles:

Ground equipment: 80° or 110° stainless steel flat fan nozzles tilted forward at an angle of 45°.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Application under hot (above 28°C) or droughty conditions will result in reduced weed control. Hoe-Grass 284 provides optimum results under cool, moist conditions.

Under certain environmental conditions, yellow blotches may appear on barley leaves. Under certain environmental conditions, especially hot, humid weather, leaf scorch on cereals and leaf cupping and burn on broadleaf crops may result from Hoe-Grass 284 application.

Tank Mixes:

Herbicides:

CROP	HERBICIDE TANK MIX
Wheat (spring and durum), Flax (NOT including Solin varieties), barley. Use 1.13 L/acre of Hoe-Grass 284.	Pardner - 0.4 L/acre. Will not control wild mustard beyond the 3 leaf stage.
Canola. Use 1.13 L/acre of Hoe-Grass 284.	Lontrel - 0.17 L/acre.

Fertilizers: None registered.

Insecticides: Hoe-Grass 284 may be tank mixed with Decis 5EC on wheat (spring, winter and durum), barley, flax (NOT including Solin varieties), rapeseed (canola), mustard and potatoes. See Decis 5EC label for recommended rates and cautions.

Fungicides: None registered.

Do not tank mix Hoe-Grass 284 with any pesticide for aerial application. Do not tank mix Hoe-Grass 284 under stress conditions such as prolonged periods of drought, excessively hot temperatures or low humidity as reduced control of grassy weeds may result. Apply other pesticides, fertilizers or chemical additives at least 4 days after application of Hoe-Grass 284.

Note: The above mixes are those listed on the HoeGrass 284 label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour of application will reduce control.

Grazing: Do not graze or cut for feed treated cereal crops prior to crop maturity. Do not feed treated forages in the year of treatment. Do not feed sainfoin or alsike clover for the life of the stand.

Preharvest Interval: Do not harvest within 60 days of treatment.

Recropping: No restrictions the year after application.

Aerial Application: May be applied by aircraft for weed control in wheat, barley, flax (excluding Solin varieties) or rapeseed (canola) only. Apply in a minimum 14 L per acre water volume. Do not tank mix with any other products.

Storage: Do not freeze. Shake well before using.

Environment: HoeGrass 284 is toxic to fish. Avoid drift or other contamination of water bodies, wetlands and other non-target areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger Corrosive

Hoe-Grass II

Herbicide Group – 1,6
(Refer to page 27)

Company:

Bayer CropScience (PCP#17507)

Formulation:

230 g/L diclofop methyl and 80 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 10 L.

Crops and Staging:

Wheat (including durum), triticale and spring rye - no leaf stage restrictions, but do not apply within 60 days of harvest.

Barley - 1 to 4 leaf stage and before tillering.

Flax (NOT including Solin varieties) - 2 to 4 inches (5 to 10 cm) in height.

The following seedling forage grasses grown for seed in the 2 to 5 leaf stage:

- bromegrass
- creeping red fescue
- crested wheatgrass
- intermediate wheatgrass
- Russian wild ryegrass

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Grasses:

WEED	STAGE
Wild oats, green & yellow foxtail, barnyard grass	1 to 4 leaf
Persian darnel	1 to 3 leaf
Volunteer corn	6 to 10 inches (15 to 25 cm) in height

Broadleaves:

Apply up to the early 4 leaf stage unless otherwise indicated

- Common groundsel
- Cow cockle
- Green smartweed
- Kochia
- Knawel
- Lady's-thumb
- Lamb's-quarters
- Night-flowering catchfly
- Redroot pigweed
- Russian thistle - up to 2 in. (5 cm) tall
- Scentless chamomile
- Stinkweed
- Tartary buckwheat
- Wild buckwheat
- Wild mustard

Best results are obtained from early applications (2 to 3 leaf stage of grassy and broadleaf weeds).

Cost (2004 suggested retail price):

$19.78 per acre.

Rates:

1.4 L per acre (one container treats 7 acres).

Application Information:

Water Volume: 45 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° stainless steel flat fan nozzles tilted forward at an angle of 45°.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Applications of Hoe-Grass II during periods of crop stress, when plants are not actively growing because of hot temperatures (above 28°C) or under very dry conditions or low humidity, may result in substantially reduced weed control. Under certain environmental conditions yellow blotches may appear on crop leaves. During periods of stress (above 28°C) or high humidity, flax may show leaf burn, retarded growth and a slight delay in maturity. Avoid spraying flax under these conditions.

Tank Mixes:

Herbicides: Hoe-Grass II may be tank mixed with MCPA amine (500 g/L) at the reduced rate of 28 mL/acre to improve wild mustard control in wheat (spring and durum), barley, spring rye and triticale.

Exceeding the recommended rate of MCPA amine 500 will result in a severe reduction in grassy weed control. Although registered, this tank mix is not recommended because of inconsistent results and possible antagonism of grassy weed control.

Fertilizers: None registered.

Insecticides: Hoe-Grass II may be tank mixed with Decis 5EC on wheat (spring and durum), barley and flax (NOT including Solin varieties). Allow a 4 day interval between application of Hoe-Grass II and other herbicides.

Fungicides: None registered.

Note: The above mixes are those listed on the HoeGrass II label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour will reduce control.

Grazing: Do not graze treated crops or cut for hay.

Preharvest Interval: Do not apply within 60 days of harvest.

Recropping: No restrictions the year after application.

Aerial Application: Do not apply by air.

Storage: Do not freeze. Shake well before using.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Horizon

Herbicide Group – 1

(Refer to page 27)

Company:

Syngenta

Formulation:

Horizon (PCP#24076): 240 g/L clodinafop-propargyl formulated as an emulsifiable concentrate.
Container size – 3.68 L.

Score adjuvant (PCP#12200): 2 x 6.4 L.

Crops and Staging:

Spring wheat (including durum) - up to the emergence of the 4th tiller.

When tank mixing, check broadleaf product description for additional restrictions.

Weeds, Rates and Staging:

Apply at 95 mL per acre (40 acres per case) for:

WEED	STAGE
Barnyard grass	1 to 5 leaf prior to tillering
Green and yellow foxtail	1 to 5 leaf stage, prior to emergence of 3rd tiller
Volunteer canaryseed, wild oats	1 to 6 leaf, maximum 3 tillers
Volunteer oats	3 to 6 leaf, maximum 3 tillers

Apply at 115 mL per acre (32 acres per case) to control the above weeds plus

WEED	STAGE
Persian darnel	1 to 5 leaf prior to tillering

Apply at the 2 to 3 leaf stage for optimum control.

Optimum weed control and yield response occurs when weeds are controlled before tillering.

Add Score adjuvant at a rate of 1 L per 100 L spray solution.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$17.61 to $22.01 per acre.

Application Information:

Water Volume: Minimum 40 L per acre.

Pressure: 40 to 45 psi (275 to 310 kPa).

Nozzles: 80° or 110° flat fan stainless steel nozzles tilted forward at an angle of 45°.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

For optimum results, apply to actively growing weeds. Do not apply to crops or weeds that are stressed by hot or cool conditions, frost, drought, low fertility, water-saturated soil, disease or insect damage as crop injury and poor weed control may result.

Tank Mixes:

Mixes provide control of wild oat, green foxtail, and weeds/insects controlled by the tank mix partner unless otherwise noted.

Herbicides:

In spring wheat (including durum except where indicated):
2,4-D amine (0.34 to 0.45 L/acre)
Ally (3 g/acre)
Attain (40 acres/case) (NOT in durum wheat)
Bromoxynil+MCPA (Buctril M*** or Mextrol only) (label rates)
Curtail M (0.81 L/acre)
Dichlorprop + 2,4-D (0.71 L/acre)**
DyVel (0.50 L/acre)
Lontrel (0.17 to 0.34 L/acre)
Lontrel (0.11 to 0.17 L/acre) + MCPA ester (0.45 L/acre)
Lontrel (85 mL/acre) + Refine Extra (8 g/acre)
MCPA amine or ester (0.34 to 0.45 L/acre)
MCPA Sodium Salt (0.48 to 1.09 L/acre)*
Pardner (0.40 L/acre)
Prestige (20 acres/case) (NOT in durum wheat)
Refine Extra (8 g/acre)
Target (0.4 to 0.6 L/acre)**
Thumper (0.40 L/acre)
Trophy (20 acres per case) (NOT in durum wheat)
Unity (40 acres/case)

* Rate above 0.81 L/acre may cause crop injury.
** Barnyard grass also controlled
*** Barnyard grass and Persian darnel also controlled

Refer to the broadleaf herbicide label for crop staging, and other information. When tank mixing, always add the broadleaf herbicide first, followed by Horizon, with the Score adjuvant added last. Reductions in green foxtail and wild oat control may be observed when tank mixed with 2,4-D amine and MCPA amine.

Fertilizers: None registered.

Insecticides: Horizon can be tank mixed with:
Decis 5F (32 to 48 mL/acre).
Matador (25 to 33 mL/acre).

Fungicides:
Tilt (0.2 L/acre)

Note: The above mixes are those listed on the Horizon label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13

Restrictions:

Rainfall: Rainfall within 30 minutes may reduce control.

Grazing: Do not graze or harvest treated crops for forage within 3 days of application.

Preharvest: Leave at least 60 days from application to harvest.

Recropping: No restrictions.

Storage: May be frozen.

Aerial Application: Do not apply by air.

Environment: Do not apply within 15 m of rivers, lakes, ponds, wetlands, sloughs, irrigation water, vegetated ditch banks, or water used for livestock or domestic purposes.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Warning Eye and Skin Irritant

Horizon BTM

Herbicide Group – 1, 4, 6
(Refer to page 27)

This product is a prepackaged tank mix of Horizon (page 135) and Buctril M (page 81). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Syngenta

Formulation:

The Horizon BTM package has three components:

Horizon 240 EC (PCP#24067): 240g/L clodinafop - propargyl formulated as an emulsifiable concentrate. Container size - 1.84 L

Buctril M (PCP#18022): 280g/L bromoxynil and 280g/L MCPA ester formulated as an emulsifiable concentrate. Container size -8 L

Score (PCP#12200). Container size: 6.4 L

Each case treats 20 acres.

Crops and Staging:

Spring wheat (including durum) from the 2 leaf stage up to the emergence of 4th tiller.

Weeds and Staging:

Grass Weeds:

Wild oat	1 to 6 leaves to the emergence of the 4th tiller
Green foxtail	1 to 5 leaves to the emergence of the 3rd tiller.

Broadleaf Weeds:

Weeds controlled up to 4 leaf stage:

American nightshade
Bluebur
Ball mustard
Cocklebur
Cow cockle
Flixweed
Green smartweed
Kochia**
Lady's-thumb
Night-flowering catchfly
Pale smartweed
Redroot pigweed*
Russian thistle**
Scentless chamomile***
Shepherd's-purse
Volunteer canola
Volunteer sunflower

* May not be controlled in flax.
** Control before plants are 2 inches tall.
*** Spring seedlings only.

Weed controlled up to 6 leaf stage:

Wild tomato

Weeds controlled up to 8 leaf stage:

Common groundsel
Common ragweed
Lamb's-quarters
Stinkweed
Tame buckwheat
Tartary buckwheat
Wild buckwheat
Wild mustard
Wormseed mustard

Weeds with top growth controlled:

Canada thistle
Perennial sow-thistle

Cost (2004 suggested retail price):

$23.64 per acre.

Rates:

0.40 L/acre Buctril M
92 mL/acre Horizon
0.32 L/acre Horizon BTM adjuvant

One case treats 20 acres.

Add Buctril M first, followed by Horizon 240 EC with the Horizon BTM adjuvant added last.

Application Information:

Water Volume: 40 L per acre

K2

Herbicide Group – 2
(Refer to page 27)

This product is a prepackaged tank mix of Everest (page 107) and Refine Extra (page 176). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

E. I. duPont Canada Inc.

Formulation:

Refine Extra (PCP#22575): 50% thifensulfuron methyl plus 25% tribenuron methyl formulated as a dry flowable.
Container size: 4 water-soluble bags (80 g each, total 320 g).

Everest (PCP#26448): 70% flucarbazone-sodium formulated as a water dispersible granule.
Container size: 4 x 174g water-soluble packets.

Crops and Staging:

Spring wheat (NOT including durum) - Apply from 1 (expanded) to 4 main stem leaves up to two tillers*. Injury may result if applied outside the recommended stages.

* **Caution:** K2 MUST be applied in a mix with with 2,4-D ester (not supplied) at 0.35 L to 0.45 L/acre (based on 500g/L formulation) plus 0.25% v/v of a recommended surfactant. See 2,4-D for additional staging and application restrictions. 2,4-D staging is 4 leaf to early flag leaf.

Weeds and Staging:

Grasses:
Wild oats and green foxtail at the 1 to 6 leaf stage (4 leaves on the main stem plus 2 tillers)

Broadleaves:
Unless otherwise indicated, apply to young, actively growing broadleaf weeds that are up to 4 inches (10 cm) tall or wide.

Weeds Controlled (when tank mixed with 2,4-D):

Burdock (seedlings)
Chickweed
Cocklebur
Common ragweed
Common plantain
Common groundsel
Corn spurry
Cow cockle
False ragweed
Flixweed
Mustard (ball, hare's ear, Indian, tumble, wild, wormseed)
Narrow-leaved hawk's-beard
Pigweed (redroot, Russian)
Prickly lettuce
Russian thistle
Shepherd's-purse
Stinkweed
Sweet clover
Giant ragweed
Green smartweed
Hemp-nettle
Kochia
Lady's-thumb
Lamb's-quarters
Tartary buckwheat
Volunteer canola
Volunteer sunflower
Wild Buckwheat
Wild radish

Weeds Suppressed:

Canada thistle (less than 6 inches (15 cm) tall or across and prior to budding)
Cleavers (1 to 3 whorls)
Round-leaved mallow (2 to 6 leaf)
Scentless chamomile
Sow-thistle
Stork's-bill (2-6 leaf)
Toadflax (less than 15cm tall)

Cost (2004 suggested retail price):

$21.46 per acre not including cost of 2,4-D ester and adjuvant.

Rates:

Refine Extra: 8 g per acre

Everest: 17.4 g per acre. (One case treats 40 acres)

2,4-D ester (not supplied) must be mixed with K2 at 0.34 to 0.45 L/acre (based on 500g/ L formulation) for improved crop tolerance.

Agral 90 or Ag-Surf at.0.25 L per 100L of spray solution

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application information:

Water Volume: 40 L per acre.

Pressure: 30 to 50 psi (200 to 345 kPa).

Nozzles: Flat Fan nozzles at 80° or 110°. Use of flood-jet or controlled droplet application equipment is not recommended.

Tank Cleaning:

Both components of K2 can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray K2 should be flushed out immediately before using another product. Follow the steps below:

1. Drain the tank and thoroughly rinse the spray tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with clean water and 1 L of 3% household ammonia per 100L of water. Flush the hoses, boom and nozzles with the solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank.
3. All nozzles, screens and filters should be removed and cleaned after applying this product. Clean nozzles and screens in a separate container using the ammonia and water.
4. Repeat #2.
5. Rinse tank, boom and hoses with clean water.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

For additional information on tank cleaning see page 14.

Kerb 50-WSP

Herbicide Group – 15

(Refer to page 27)

Company:

Dow AgroSciences (PCP#25595)

Formulation:

50% propyzamide formulated as a wettable powder. Container size - 1.36 kg (3 x 454 g water soluble pouches).

Crops and Staging:

Apply to the following established crops between October 1 and freeze-up or very early spring*:

Seedling and established alfalfa, bird's-foot trefoil, and established pastures**.

* Seed alfalfa only.

** Severe stand thinning may occur to pastures consisting primarily of crested wheatgrass, meadow fescue and timothy. Some thinning (10 to 15%) may occur with tall fescue and creeping red fescue.

Weeds, Rates and Staging:

Apply in late fall or very early spring (seed alfalfa only) prior to the emergence of weeds.

Established grass or grass/legume pastures for control of foxtail barley:
Brown, dark brown or gray wooded soils - 0.36 kg/acre.
Black soils - 0.45 kg/acre.

Alfalfa and bird's foot trefoil:
Annual grasses, volunteer cereals, wild oat - 0.71 kg/acre.
Quackgrass, orchardgrass, timothy, chickweed - 0.91 to 1.32 kg/acre*.
Dodder control (fall application only) - 1.3 kg/acre. Note that complete control may not be achieved.

* Maximum 0.91 kg per acre with spring application. Low temperatures and adequate moisture following application are needed for efficacy.

Caution: Do not use on soils with more than 6% organic matter. Do not apply to soils prone to flooding. Do not apply to pastures that contain high proportions of timothy, crested wheat grass or meadow fescue. Consult the manufacturer for other forage grass species sensitivities to Kerb 50-WSP.

Cost (2004 suggested retail price):

$91.78 per kg.

Application Information:

Water Volume: 120 to 200 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Dry soil conditions at time of weed emergence may result in reduced control. Approximately 3 inches of total precipitation is required for adequate activation. Best results when soil temperatures are low but above freezing.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Surface applications are most effective if followed by 0.5 to 1 inch (1.25 to 2.5 cm) of rain within 2 days of application.

Grazing: Do not graze or harvest for livestock feed within 90 days of the 1.32 kg/acre rate and 60 days of application for lower rates.

Recropping: Do not replant to crops within 1 year of treatment.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. Do not freeze

Environment: Do not contaminate domestic or natural water sources or wetlands.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Caution Poison

Laddok

Herbicide Groups – 5,6
(Refer to page 27)

Company:

BASF Canada (PCP#16641)

Formulation:

200 g/L bentazon and 200 g/L atrazine formulated as a solution.
Container size - 10 L.

Crops and Staging:

Corn (grain, silage and seed) at the 1 to 5 leaf stage. Seed corn producers should consult their seed corn company regarding the tolerance of seed production lines to Laddok and oil concentrates.

Weeds, Staging and Rates:

Add Assist at 1.0 L per 100 L of spray solution.

ANNUAL WEEDS	1.21 L/ACRE		1.62 L/ACRE	
	inches	(cm)	inches	(cm)
Black nightshade			0.2 - 0.8	(0.5 - 2.0)
Buttercup			2 - 4	(5 - 10)
Cocklebur	3 - 7	(7.5 - 17.5)	7 - 12	(17.5 - 30.0)
Common chickweed			Apply 1 to 3 weeks after emergence	
Common groundsel			2 - 4	(5 - 10)
Common ragweed	1 - 1.5	(2.5 - 4.0)	1.5 - 3	(4.0 - 7.5)
Corn spurry			1 - 4	(2.5 - 10.0)
Giant ragweed			2 - 6	(5 - 15)
Hairy galinsoga			2 - 3	(5.0 - 7.5)
Lady's-thumb (smartweed)	1 - 3	(2.5 - 7.5)	3 - 8	(7.5 - 20.0)
Lamb's-quarters	1 - 2	(2.5 - 5.0)	2 - 4	(5 - 10)
Purslane			1 - 2	(2.5 - 5.0)
Redroot pigweed	0.7 - 1.2	(1.5 - 3.0)	1.2 - 1.5	(3 - 4)
Russian thistle			1 - 3	(2.5 - 7.5)
Venice mallow	1 - 2	(2.5 - 5.0)	2 - 4	(5 - 10)
Wild mustard	1 - 5	(2.5 - 12.5)	5 - 10	(12.5 - 25.0)
PERENNIAL WEEDS	Rate: 1.21 - 1.62 L/acre			
Yellow nutsedge	4 - 6	(10 - 15)	(repeat 7 - 10 days if necessary)	

Cost (2004 suggested retail price):

$15.97 to $21.38 per acre.

Application Information:

Water Volume: 80 to 160 L per acre. The higher recommended water volumes and pressures should be used for dense weed infestations or for weeds approaching the upper limit of their recommended stage for treatment.

Pressure: 40 to 45 psi (275 to 300 kPa).

Nozzles: Flat fan nozzles capable of delivering high water volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Crop injury may occur if applications are made when crop is under stress because of prolonged cold weather, poor fertility or other factors; or when corn is wet and succulent from recent rainfall.

Best weed control is under warm, humid conditions.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 6 to 8 hours of application will reduce control.

Grazing: Do not graze the treated crop or cut for hay prior to crop maturity.

Recropping: No restrictions the year following treatment when label rates for annual weeds have been used. Use of the double application for yellow nutsedge control may cause injury to rotational crops because of carryover of the Atrazine component of the product. Under particularly dry soil conditions, there may be greater carryover of the atrazine component of Laddok that could damage a number of different special crops. Refer to the "Atrazine" product description for information.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environment: Avoid conditions that will risk drift to non-target plant, water sources or wetlands.

Tank Cleaning:

The manufacturer does not provide information on tank cleaning.

Refer to Basagran and Atrazine recommendations and page 14.

Hazard Rating:

☠ Caution Poison

Liberty

Herbicide Group – 10

(Refer to page 27)

Company:

Bayer CropScience (PCP#26694)

Formulation:

150 g/L glufosinate ammonium formulated as a solution. Container sizes - 13.5 L, 108 L.

Crops and Staging:

Liberty Link Canola - cotyledon to early bolting stage. Temporary crop discolouration (bronzing) may be observed after application.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

For Desiccation use only:

Alfalfa (seed production only) - 50 to 75% pod turn (brown)
Lentils* - 40 to 60% pod turn (yellow to brown)
Potatoes* - 14 to 21 days prior to harvest.
*Not for crops grown for seed production.

Weeds, Rates and Staging:

WEED	WEED STAGE (from emergence to stage)	RATE (L/ACRE)	ACRES PER 13.5 L
Cow cockle	4 leaf	0.54	25
Green foxtail	6 leaf (max. 3 tillers)		
Barnyard grass, volunteer canola	4 leaf	0.81	16.6
Wild mustard	5 leaf		
Lamb's-quarters, smartweed (lady's-thumb)	6 leaf		
Stinkweed	8 leaf		
Volunteer flax	2.5 inches (6 cm)		
Russian thistle	3 inches (8 cm)		
Wild buckwheat	3 leaf	1.1	12.4
Redroot pigweed, round-leaved mallow, quackgrass*	4 leaf		
Light to moderate infestationsΔ of volunteer wheat, volunteer barley*	4 leaf (max. 2 tillers)		
Hemp-nettle (1 to 3 leaf pairs), shepherd's purse	6 leaf		
Common chickweed (max. 4 leaf pairs), Perennial sow-thistle	8 leaf		
Kochia	3 inches (8 cm)		
Canada thistle*, scentless chamomile	4 inches (10 cm)		
Cleavers	2 whorls (nodes)	1.35	10
Stork's-bill and heavy populations of wild buckwheat	3 leaf		
Quackgrass (light to moderate** or heavy infestations*)Δ, volunteer wheat, volunteer barley*, wild oat	4 leaf (max. 2 tillers except quackgrass)		
Hemp-nettle	8 leaf (1 to 4 leaf pairs)		
Dandelion rosettes	6 in. (15 cm) across		
Flixweed, Canada thistle*	4 inches (10 cm)		
Quackgrass***	4 leaf	1.6	8.3
Canada thistle**	4 inches (10 cm)		

* Temporary top growth control. Plants may return from surviving growing points.

** Extended Top growth control.

***Season long control

Δ The company does not provide guidelines for weed densities. When in doubt as to the infestation level, use the high rate or contact the manufacturer

A second application can be made to fields that were treated initially with up to 1.35 L/acre. Apply Liberty at a rate of 0.81 to 1.35 L/acre in the second application.

Desiccation Rates:

Alfalfa (seed production only) - 1.09 L per acre
Lentil* - 0.81 to 1.09 L per acre
Potato* - 1.21 L per acre
* Not for crops grown for seed production.

Cost (2004 suggested retail price):

$14.04 per L.

Application Information:

Water Volume:
Ground applications: 45 L per acre.
Aerial applications:
In Liberty Link canola use 22 L per acre.
For desiccation, 13 to 22 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles:
Ground Application: 80° or 110° stainless steel flat fan nozzles tilted forward at an angle of 45°.
Aerial applications: Do not use raindrop nozzles. Use a combination of nozzles and pressure to provide a coarse droplet size distribution (mean diameter of 350 microns).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Liberty activity is influenced by environmental conditions. Cool temperatures (less than 10°C), drought, and low humidity conditions slow weed growth. Applications made under these stressed conditions may result in reduced weed control.

Tank Mixes:

Herbicides:
In Liberty Link canola:
Liberty may be tank mixed at 1.1 -1.35 L/acre with Select at 25.5 mL/acre for enhanced control of wild oat and volunteer barley. Consult Liberty label for exact weed control options. Add Amigo to the tank first at 0.5 L per 100 L spray solution followed by Liberty and then Select. Consult label for specific mixing instructions.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Liberty label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Grazing: Do not graze the treated crop or cut for feed.

Preharvest Interval: Do not harvest desiccated potato or lentil crops within 9 days of treatment.

Recropping: No restrictions.

Aerial Application: May be applied by air.

Storage: Do not freeze.

Environment: Do not apply within 15 m of environmentally sensitive areas using ground equipment or within 30 m of environmentally sensitive areas when applying by air.

Tank Cleaning:

Refer to tank cleaning section on page 14.

Hazard Rating:

☠ Warning Poison

Skin and Eye Irritant

Liberty 200 SN

(Not for use in Saskatchewan)

Herbicide Group – 10
(Refer to page 27)

Company:

Bayer CropScience (PCP#25337)

Formulation:

200 g/L of glufosinate ammonium formulated as a solution
Container sizes- 10 L
One case treats 20 acres

Crops and Staging:

For use on LIBERTY 200SN tolerant Corn only. Apply from 1-8 leaf stage of the crop, refer to product label for appropriate method of determining crop leaf stage.

Weeds Rates and Staging:

Weeds controlled with 0.61 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Green foxtail	5 leaf
Proso millet	5 leaf
Chickweed	8 leaf
Cocklebur	4 leaf
Ragweed	5 leaf
Redroot pigweed	6 leaf
Shepherd's purse	6 leaf

Weeds controlled with 0.81 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Barnyard grass	5 leaf
Wild oat	4 leaf
Yellow foxtail	4 leaf
Eastern black nightshade	5 leaf
Lady's thumb	6 leaf
Lamb's quarters	6 leaf
Perennial sowthistle	4 leaf
Ragweed	7 leaf
Stinkweed	8 leaf
Wild buckwheat	4 leaf
Wild mustard	4 leaf
Wormseed mustard	6 leaf
Field bindweed*	6 leaf
Canada thistle*	6 leaf

* season long suppression

Weeds controlled with 1.0 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Quackgrass **	4 leaf

** season long suppression, apply with ammonium sulphate, 2.4 L/acre (49% solution) or 1.2 kg/acre (99%).

Second Application:
A second application of LIBERTY 200SN may be made to fields treated initially with up to 1 L/acre, if weeds are present. Apply when new weed growth is in the correct leaf staging, and before maximum allowable leaf staging for crop. Do not apply more than 1.8 L/acre LIBERTY 200SN to a crop in a single season.

Split Application Program:
For season long control of the above listed weeds a split application of Liberty 200SN may be employed. The first application must be a minimum of 0.81 L/acre LIBERTY 200SN made at the correct weed staging. For the second application of LIBERTY 200SN a 0.51 L/acre rate may be used. The second application timing must be made as soon as the second flush of weeds occurs and before the maximum leaf stage for the crop.

Cost: (2004 sugested retail price)

Not available

Application Information:

Water Volume: A minimum of 45 L per acre

Pressure: 175-275 kPa (25-40 psi)

Speed: 6-10 km/h

Nozzles: 80° or 110° flat fan nozzles tilted forward at an angle of 45°.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Liberty 200SN activity is influenced by environmental conditions. Cool temperatures (less than 10°C), drought and low humidity conditions slow weed growth. Applications made under these stress conditions may result in reduced weed control.

Weed control may also be reduced when heavy dew, fog, or mist are present at the time of application.

Tank Mixes:

None registered.

Restrictions:

For sale and use in Manitoba on glufosinate-ammonium tolerant field corn.

Rainfall: Rainfall within 4 hours of application may reduce weed control

Grazing: Do not graze treated fields within 20 days of application.

Pre-harvest Interval: Do not apply within 86 days of harvesting corn grain.

Recropping: No recropping restrictions the year after treatment.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environmental: Do not contaminate water supplies, ponds, lakes, streams, and irrigation ditches by direct applications, spray drift, or when cleaning and rinsing spray equipment or containers. Leave a 15 meter buffer between the edge of the treated field and adjacent crops not tolerant to LIBERTY 200SN or environmentally sensitive areas (e.g. wetlands, ponds, rivers, or other open bodies of water, shelter belts or wildlife habitat).

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Warning Eye and Skin Irritant

Linuron

Herbicide Group – 7

(Refer to page 27)

Company/Products:

E. I. duPont Canada (Lorox Toss-N-Go – PCP#21353, Lorox L – PCP#16279)

United Agri Products (Linuron 400 – PCP#15544)

The following recommendations are a blend of recommendations of all Linuron products. Consult the individual product labels for specific recommendations.

Formulations:

Linuron 400: 400 g/L linuron formulated as a flowable. Container size - 10 L.

Lorox L: 480 g/L linuron formulated as a flowable. Container size - 10 L.

Lorox Toss-N-Go: 50% linuron formulated as a dry flowable. Container size - 10 kg).

*Linuron 400 may be in short supply in the distribution system for 2005 but is expected to be produced in the future.

Crops, Rates and Stages:

Post-emergent applications only:

CROP	STAGE	LINURON 400 (L/acre)	LOROX L (L/acre)	LOROX TNG (kg/acre)
Spring wheat (including durum), oats and barley*	2 to 4 leaf stage	0.20 - 0.26	0.17 - 0.22	0.17 - 0.22
Field corn (post-emergent** directed spray)	Apply when corn is at least 15 inches (38 cm) high (highest leaf on free standing plant)	1.16 - 2.18	0.97 - 1.82	0.93 - 1.74
Shelterbelts (caragana, green ash, Siberian and American elm, Manitoba maple, poplar, willow, white spruce, Colorado spruce, Scots pine)	Apply as an overall spray to dormant stock or as a directed spray if buds have broken.	2.18	1.82	1.7

* Only when tank mixed with MCPA amine at 0.45 L/acre

** Use lower rate when weeds do not exceed 2 inches (5 cm) and higher rate for weeds up to 8 inches (20 cm) in height, preferably before they are 5 inches (13 cm) high. Requires the addition of a mineral oil surfactant blend at 1 to 2 L per 100 L or spray solution or spray oil at 1 to 2 L per 10 L of spray solution. See oil labels for directions. DO NOT apply if linuron has been applied pre-emergent.

Pre-emergent surface applications for use on loam to clay soils only:

PRODUCT	LINURON 400 (L/acre)		LOROX L (L/acre)		LOROX TNG (kg/acre)	
Soil organic matter	less than 2%	from 2 - 5%	less than 2%	from 2 - 5%	less than 2%	from 2 - 5%
Field corn	1.09*	1.58	0.91**	1.31**	0.30**	0.61**
Soybeans	1.09 - 1.58	1.58 - 2.18	0.91 - 1.31	1.31 - 1.82	0.91 - 1.31	1.31 - 1.82
Sweet white lupins	1.01	1.50	0.85	1.25	0.81	1.21
Potatoes	1.11 - 1.72	1.72 - 2.22	0.91	1.82	0.89	1.74

* Must be tank mixed with Atrazine 80W. Refer to label for tank mixes with Dual II and/or Primextra.

** Must be tank mixed with Dual II Magnum + Atrazine or Primextra.

If used on sandy soils, severe crop injury may result.
Seed the crop at least 2 inches (5 cm) deep.Make only one application per year to field crops.
When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Banded Applications:
This product may also be applied in a narrow band directly over the row in wide rowed crops if another method is to be used weed control in between the rows. For band treatment, use proportionately less; for example, for 10 inch band on 30 inch row, use 1/3 of the broadcast rate.

Weeds and Staging:

Post-Emergence

When tank mixed with MCPA amine in cereals, the following weeds are controlled:

Chickweed
Corn Spurry
Cow cockle
Flixweed
Green foxtail
(suppression possible)
Green smartweed
Hemp-nettle
Lady's-thumb
Lamb's-quarters
Ragweed (Common, giant)
Redroot pigweed
Shepherd's-purse
Stinkweed
Stork's-bill
Tartary buckwheat
Wild buckwheat

Apply when annual broadleaf weeds are in the 2 to 4 leaf stage and when green foxtail is in the 1 to 3 leaf stage.

In shelterbelts, apply when weeds are less than 4 inches (10 cm) tall.

Pre-emergent surface treatments and Post-emergent applications in corn and shelterbelts:
Sufficient moisture (1 to 2 inches or 3 to 5 cm) in the form of rainfall or irrigation is necessary within 7 to 10 days of a pre-emergence application or poor weed control will result.

Barnyard grass*
Common chickweed
Common Groundsel■
Corn Spurry■
Crabgrass*
Dandelion (seedlings only)■
Foxtail (green and yellow)*
Goosefoot
Knotweed
Kochia■
Lamb's-quarters
Pigweed (Prostrate■, Redroot)
Plantain (seedlings only)■
Purslane
Ragweed (Common)
Shepherd's-purse
Smartweed (Annual)
Sow-thistle (Annual, perennial■ seedlings only)
Stinkweed▫
Wild buckwheat
Wild Radish■
Witchgrass
Wormseed mustard

* partial control
■ Not registered with Lorox products.
▫ Not registered with Linuron 400.

Costs (2004 suggested retail prices):

Lorox - $33.38 per kg
Lorox L - $25.28 per L
Linuron 400 - $26.50 per L

Application Information:

Water Volume:
Post-emergent: Cereals - 40 L per acre.
Field Corn directed spray – 69 to 138 L per acre.
Pre-emergent surface: 81 to 162 L per acre.
Shelterbelts: 90 to 180 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Stainless steel, flat fan nozzles with 50-mesh (or coarser) line strainers and screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

In post-emergent applications the best weed control occurs when temperatures are moderate, when relative humidity is high and when soil moisture is adequate. Injury to cereals (crop lightening) will occur when the crop is under stress because of drought or disease. This injury is worse when the product is applied at advanced leaf stages. In pre-emergent surface treatments, rainfall or irrigation (1 to 2 inches or 3 to 5 cm) is required to move linuron into the root zone of germinating seeds. Insufficient moisture will result in poor weed control. Drought conditions after application will result in little to no weed control. If rainfall does not occur within 7 to 10 days of application and prior to crop emergence, a shallow rotary hoeing (0.75 to 1.5 inches/ 2 to 4 cm) should be made to mix the top layer of soil to help activation. Avoid covering treated ground with untreated soil. If unusually heavy rain follows application, severe crop injury may result from herbicide in the root zone of the crop. Do not use on sandy soils or severe crop injury will result.

Tank Mixes:

Herbicides: For post-emergent applications in cereals, linuron must be tank mixed with MCPA amine. Do not tank mix with other herbicides.

Fertilizers: None registered.

Insecticides: None registered.

Note: The above mixes are those listed on the linuron labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Pre-emergent applications require rainfall for activation. Contact manufacturer for more information.

Grazing: Do not graze treated crops or cut for feed prior to crop maturity.

Preharvest Interval:
Sweet Corn: Do not harvest within 50 days of treatment.
Field Corn: Do not harvest within 60 days of treatment or until after tassle emergence.

Recropping: If the intended crop fails, fields treated with Pre-emergent surface applications of linuron, may be seeded back to corn, soybeans, sweet white lupins, or potatoes. Till the soil thoroughly before reseeding. No restrictions 1 year after treatment.

Aerial Application: Do not apply by air.

Storage: Do not store liquid linuron formulations at temperatures below 5°C. Lorox may be frozen. Avoid handling Toss-N-Go bags under freezing temperatures.

Tank Cleaning:

Refer to page 14.

Lontrel

Herbicide Group – 4
(Refer to page 27)

Company:

Dow AgroSciences (PCP#23545)

Formulation:

360 g/L clopyralid formulated as a solution.
Container size - 4.45 L.

Crops Rates and Staging:

Applications of 0.17 to 0.23 L/acre (26.2 to 19.3 acres/jug):
Barley, spring wheat (NOT including durum), oat - 3 to flag leaf emergence stage.

Applications of 0.23 to 0.34 L/acre (19.3 to 13.1 acres/jug):
Flax (not including Solin) - 2 to 4 inches (5 to 10 cm) in height.

Applications of 0.17 to 0.34 L/acre (26.2 to 13.1 acres/jug):
Canola - 2 to 6 leaf stage.
Seedling forage grasses - 2 to 4 leaf stage.
Established grasses - at the shot blade stage, or in the fall after harvest or in early spring.
Seedling and established grasses for forage and seed production include:

Altai wild ryegrass
Creeping red fescue
Crested wheatgrass
Intermediate wheatgrass
Kentucky bluegrass
Meadow fescue
Meadow foxtail
Orchardgrass
Reed canarygrass
Slender wheatgrass
Smooth bromegrass
Streambank wheatgrass
Tall fescue
Tall wheatgrass
  (for forage only)
Timothy
Russian wild ryegrass

Applications of 0.34 L per acre (13.1 acres per jug):
Summerfallow – Stage according to weeds.
Shelterbelts containing villosa lilac, acute willow, Colorado spruce, white spruce, buffaloberry and chokecherry.*

* Since this product is registered under User Requested Minor Use Label Expansion program, the manufacturer assumes no responsibility for herbicide performance. *Users of this product for these uses do so at their own risk.*

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

The following weeds will be controlled when young and actively growing:

Alsike clover.
Common groundsel
Perennial sow-thistle
  (top growth only)
Scentless chamomile
Wild buckwheat
Volunteer alfalfa - 2 to 20
  inches (5 to 50 cm) in
  height.

Canada thistle - after all thistles have emerged and when the majority are in the rosette to pre-bud stage;

0.17 L/acre - Provides top growth control of Canada thistle for 6 to 8 weeks.

0.23 L/acre - Provides season long control of Canada thistle. Not all root stalks will be killed and some regrowth may occur by the end of the growing season.

0.34 L/acre - Provides season long control of Canada thistle with suppression into the following year.

Cost (2004 suggested retail price):

$24.91 to $49.82 per acre.

Application Information:

Water Volume: 40 to 89 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to table 2 on page 29.

Effects of Growing Conditions:

Poor control may occur under dry conditions. Injury to flax may occur when tank mixing with MCPA. To reduce the risk of crop injury, do not apply tank mixes if temperature exceeds 27°C.

Tank Mixes:

Herbicides:

Lontrel applications following applications of products containing bromoxynil (Badge, Buctril M, Koril, Mextrol, Pardner, Thumper, Hoegrass II) should be delayed by 14 days to allow the Canada thistle to recover from leaf burn.

Recommended rates of Lontrel may be used for each crop unless otherwise indicated.

In Canola:
Poast Ultra plus Merge adjuvant.
Fusion (Lontrel at 0.23 L/acre)
Hoegrass 284
Roundup Original* /Transorb* (Lontrel at 0.112 L/acre)
Odyssey** (Lontrel at 0.17 to 0.23 L/acre)
Select plus Amigo adjuvant (Lontrel at 0.17 to 0.34 L/acre)
Vantage* /Vantage Plus* (Lontrel at 0.112 L/acre)

* Roundup Ready varieties only
** CLEARFIELD varieties only

In Flax:
MCPA amine or ester - 0.34 to 0.45 L/acre
(Lontrel at 0.17 L/acre)
Poast Ultra plus Merge adjuvant.
Poast Ultra + MCPA Ester (rates as above)
Select plus Amigo adjuvant.

In Spring wheat (NOT including durum) and barley:
Lontrel at 85 mL/acre for the following mixes:
Refine Extra (8 g/acre) + 2,4-D (0.28 L/acre - 600g/L formulation) plus non-ionic surfactant.
Refine Extra (8 g/acre) + MCPA (0.34 L/acre - 500g/L formulation) plus non-ionic surfactant.
Triumph Plus - 20 acres per case
Lontrel at 0.11 to 0.17 L/acre for the following mix:
2,4-D or MCPA (amine or ester) at label rates

In Oat:
MCPA amine or ester (Lontrel at 0.11 to 0.17 L/acre)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Lontrel label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions.

General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Crops or areas treated with this product may be grazed immediately following treatment.

Recropping: Lontrel residues in the soil may affect succeeding crops. The year after application, replant to wheat, barley, oats, rye, flax, forage grasses, mustard or canola.

Do not use manure from animals fed or bedded with Lontrel-treated straw except on fields that are to be sown to Lontrel-tolerant crops.

Aerial Application: Do not apply by air.

Storage: Store in heated storage. If product is frozen, bring to room temperature and agitate before use.

Environment: Avoid contamination of non-target land, water or irrigation ditches.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

▽ Caution Poison

MCPA

Herbicide Group – 4

(Refer to page 27)

Company, Formulation and Cost (2004 suggested retail prices):

	Price per L (PCP number)							
	500 Ester**		500 Amine*		400 (K) Potassium salt*		300 (Na) Sodium salt*	
IPCO	$9.65	(20307)	$8.35	(20308)	$8.90	(20305)	$8.50	(20306)
NuFarm	$8.60	(27054)	$7.40	(14730)	$6.55	(17564)	$5.85	(14718)
United Agri Products	$8.71	(22199)	$7.83	(9516)	$6.15	(14188)	$6.12	(9858)

* formulated as a solution.

** formulated as an emulsifiable concentrate.

Crops, weeds and rates and other application details may differ with different product labels. Consult specific product labels for more information.

Crops, Rates and Staging:

The maximum safe rates for various crops are given below. Higher rates used for harder to control weeds (see "Weeds, Rates and Staging") may cause crop injury. Application rates for individual products may vary from those listed. Refer to the label for product specific use rates. Rates greater than these for harder to control weeds may cause crop injury. When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	STAGE	RATE (L/ACRE)			
		AMINE	ESTER	K SALT	NA SALT
Wheat (spring and durum), barley, oats	3 leaf to early flag leaf.	0.45	0.45	0.71	0.81
Spring rye	2 leaf to early flag leaf.	0.45	0.45	**NR**	0.81
Flax	2 inches (5 cm) in height to prebud stage. Apply at 2 to 4 inches (5 to 10 cm) in height for maximum crop tolerance.	0.4	0.34	0.65	0.81
Winter wheat, fall rye	In spring, apply from the time growth commences until the early flag leaf stage. Winter wheat may be treated in the fall for winter annual weed control (caution: this is not a registered treatment, any such use is at the risk of the user).	0.45	0.45	**NR**	0.81
Corn	As a broadcast spray up to 6 – 7 in. (15 – 18 cm) tall or six leaf stage. Up to 3 weeks before tassling as a directed spray using drop nozzles.	0.45	**NR**	**NR**	0.74
Peas	Vines 4 to 7 inches (10 to 18 cm) long.For short-statured, determinate flowering peas, apply at the early stages within this range.	0.22*	**NR**	**NR**	0.36*
Cereals underseeded to alfalfa (not Flemish varieties)	Apply when the majority of seedling legumes are in the 1 to 3 trifoliate leaf stage.	0.23	**NR**	**NR**	0.4

Crops, Rates and Staging continued on next page.

Crops, Rates and Staging *continued*

CROP	STAGE	RATE (L/ACRE) AMINE	 ESTER	 K SALT	 NA SALT
Underseeded alsike, ladino and red clover	Apply when the majority of seedling legumes are in the 1 to 3 trifoliate leaf stage.	0.28	NR	NR	0.4
Red Clover^ Seedling (seed and forage) Established^ (seed only)	*Seedlings:* 1 - 3 trifoliate stage. Do not feed to livestock in the first year. *Established:* Apply at the breaking of dormancy in the spring up to 7.5 cm.	0.23	NR	NR	NR
Grass pastures	Spring or fall.	1.42	1.38	NR	2.8
Seedling forage** grasses (not for seed)	Apply from the 3 leaf stage to the shot blade stage.	0.45**	NR	NR	NR
Established forage** grasses(not for seed)	Apply in the spring up to the shot blade stage or in the fall after harvest.	0.45**	NR	NR	NR

NR = Not Registered

* The rates given are lower than the registered rates for peas. Less than the maximum label rates are recommended because of crop injury concerns.

** MCPA is NOT registered for use on forage grasses, any such use is entirely at the risk of the user.

^ NuFarm MCPA Amine only

Formulation Characteristics:

Formulation	Risk of Vapour Drift	Activity on Weeds	Risk of Crop Injury
LV Ester	Medium	Fast	Medium
Amine	Very Low	Medium	Low
Salts	Very Low	Slow	Very Low

Weeds, Rates and Staging:

Apply at lower rates when weeds are small (2 to 4 leaf stage) and actively growing. Higher rates are needed when weeds are larger, in heavy populations, or growing under stressful conditions (excessively cold, hot, dry or wet). Lower rates may be applied in late fall to control winter annual weeds.

NOTE: The following rates are a general range for all products. Rate ranges for individual products may differ slightly. Consult the product label for specific rates for each application.

+ Not controlled by MCPA K salt formulation

++ Not controlled by MCPA K or Na salt formulations

Susceptible weeds:

Amine and Ester formulations – 0.28 to 0.45 L/ac
K formulations – 0.61 to 0.71 L/ac
Na formulations – 0.5 to 0.81 L/ac

- Burdock
- Cocklebur
- Flixweed (late fall applications or small seedlings)*
- Kochia
- Lamb's-quarters
- Mustards (except dog and tansy)
- Prickly lettuce+
- Ragweed
- Russian pigweed
- Shepherd's purse*
- Stinkweed*
- Wild Radish
- Wild Sunflower+

* Winter annual weeds

Harder to control weeds:

Amine & Ester formulations – 0.45 to 0.71 L/acre
K formulations – 0.71 to 0.81 L/acre
Na formulations –0.81 to 1.1 L/acre

Annual sow thistle+
Biennial wormwood+
Bluebur+
Common peppergrass+
Curled dock+
Flixweed (overwintered rosettes prior to bolting)+
Goat's-beard+
Goosefoot+
Hemp-nettle (suppression only)
Mustard (dog and tansy)+
Plantain+
Purslane+
Redroot pigweed+
Smartweed (annual)+
Tumble pigweed+

Top growth control only (rates as for harder to control weeds):

Blue lettuce+
Canada thistle
Dandelion++
Gumweed+
Field bindweed+
Hedge bindweed+
Hoary Cress+
Leafy spurge+
Perennial sow thistle+
Russian thistle++

Application Information:

Water Volumes:

Cereals, flax, pastures, forage grasses: 40 to 81 L per acre.

Peas: Minimum 61 L per acre.

Cereals Underseeded to Forage Legumes: 61 to 81 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Best weed control occurs when temperatures are above 21°C (daytime) or 10°C (night time) and humidity is above 70 percent. Do not apply if temperature exceeds 27°C. If applying to flax, injury and a delay in maturity may result from application under hot or humid conditions. Extremely hard water may reduce performance or cause problems in spraying the product.

Tank Mixes:

Herbicides:

In Wheat and barley:
Avenge (ester only),
Banvel II and Pardner (amine and K salt only),
Linuron and Sencor (amine only)

In Oats:
Banvel II and Pardner (amine and K salt only)
Linuron and Sencor (amine only)

Not all brands are labelled for tank mixing. Check the product label prior to use for registered mixes and rates. Follow all precautions and restrictions on both labels.

Fertilizers: Liquid nitrogen (28-0-0) may be used in place of water as a carrier with certain amine formulations for application in spring to winter wheat or fall rye.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the MCPA labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours of MCPA Na salt or MCPA K salt, 4 hours of MCPA amine, or 2 hours of MCPA ester application will reduce control.

Grazing: Do not graze within 7 days of application.

Recropping: No restrictions the year after application.

Aerial Application: Some products may be applied by air to specific crops. Check the label for detailed instructions.

Storage: MCPA ester may be frozen. Do not freeze MCPA amine, MCPA sodium salt and MCPA K.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

MCPB + MCPA

Herbicide Group – 4
(Refer to page 27)

Company:

IPCO (Clovitox Plus – PCP#24336)
Nufarm (Tropotox Plus – PCP#8211)
UAP (Topside – PCP#22003)

Formulation:

375 g/L MCPB, 25 g/L MCPA present as sodium salts and formulated as a solution. Container size - 10 L.

Crops, Rates and Staging:

Registered for all products:
Apply 1.11 to 1.72 L per acre. Apply only that needed to control the target weeds.

CROP	STAGE
Peas	3 to 6 expanded leaves
Clover (Alsike, Ladino, Red, White Dutch, Wild white)	Monofoliate to 3 trifoliate leaf stage (with or without a cover crop).
Oats, wheat, rye or barley (alone or as a companion crop)	2 leaf to flag leaf stage
Field Corn	45 cm high to the start of tassling – use drop nozzles
Established pasture	After grazing or cutting when weeds have regrown to a susceptible stage

Seedling Forage Grasses:
Apply at 1.11 to 1.42 L per acre from the 2 to 4 leaf stage:

Bromegrass (smooth, meadow)	Timothy
Fescue (altai, creeping red, meadow, tall)	Wheatgrass (crested, intermediate, northern, pubescent, slender, streambank, tall, western)
Green needlegrass	
Reed canarygrass	Wild ryegrass (altai, Russian)

Registered for Tropotox Plus, and Clovitox Plus only:
Seedling alfalfa for seed production* at the 3 to 6 trifoliate stage.

Note: Seedling alfalfa vigour may be reduced in the year of treatment, however, the crop recovers and yield will not normally be affected.

* Since these uses are registered under the User Requested Minor Use Label Expansion program, the manufacturers assume no responsibility for herbicide performance. *Users of this use do so at their own risk.*

Weeds, Rates and Staging:

WEEDS	RATE STAGE	(L/ACRE)
Ball mustard, Lamb's-quarters, Stinkweed, Wild mustard, Wormseed mustard	Seedlings	1.11
Annual sow-thistle*, Hempnettle*, Redroot pigweed, Ragweed, Shepherd's-purse, Volunteer rapeseed (including canola), Wild Radish*	Seedlings	1.72
Curled dock, Perennial sow-thistle**, Plantain	Rosette	1.72
Bull thistle	Rosette to early bud	1.72
Buttercup (Creeping, Tall), Field Bindweed	In spring during rapid growth	1.72
Canada thistle	6 inches (15 cm) to early bud	1.72
Horsetail	6 inches (15 cm)	1.72 *

* Suppression only
**Topgrowth control only

Cost (2004 suggested retail prices):
Clovitox Plus: $13.25 per L
Topside: $14.00 per L
Tropotox Plus: $13.25 per L

Application Information:

Water Volume:
Clovitox Plus: 71 to 91 L per acre.
Tropotox Plus, Topside: 61 to 81 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Damage to peas or seedling forage legumes may occur if the crop is sprayed when under drought or disease stress. Under extremely hot or humid conditions, crop injury may be severe. Do not apply when temperatures are over 27 degrees C. Best activity on weeds occurs in warm weather.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze crop treated with Topside or cut for hay in the year of establishment. Cereals or pastures treated with Tropotox or Clovitox may be used for grazing or cut for greenfeed or hay 30 days after application. Forage legumes and peas treated with Clovitox may be used for animal feed 30 days after application.

Recropping: No restrictions listed. Phenoxy herbicides can persist in soils for weeks, particularly if dry or cool weather persists.
Do not seed sensitive crops immediately after spraying.

Aerial Application: Clovitox may be applied by air to established pasture and **cereal crops (not underseeded to clover).**

Storage: Do not freeze.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger Poison - Clovitox

Caution Poison - Tropotox Plus & Topside

Mecoprop

Herbicide Group – 4
(Refer to page 27)

Company:

United Agri-Products (Mecoprop – PCP#10915)
Nufarm (Compitox – PCP#7363)

Formulation:

150 g/L mecoprop present as potassium salt. Container size - 10 L.

Crops and Staging:

Spring wheat (including durum), barley and oats - 3 leaf to flag leaf stage.

Weeds and Staging:

Apply to weeds from the 2 to 4 leaf stage.

- Black medic
- Canada thistle (top growth control only)
- Chickweed
- Cleavers
- Clover (volunteer)
- Corn spurry
- Lamb's-quarters
- Plantain
- Wild mustard

Cost (2004 suggested retail price):

Compitox: $6.81 per L.
Mecoprop: $7.41 per L.

Rates:

2.2 to 2.8 L/acre.

Use the high rate for weeds in an advanced stage of growth.

Application Information:

Water Volume: 81 to 121 L per acre.

Pressure: 40 psi (275 kPa).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Apply in warm weather under good growing conditions. Avoid spraying in very hot weather or in drought conditions.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze or feed treated crop to livestock prior to crop maturity.

Recropping: No restrictions the year after application.

Aerial application: Not registered.

Storage: Do not freeze.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

Muster Gold II

Herbicide Group – 1, 2
(Refer to page 27)

This product is a prepackaged tank mix of Muster (page 157) and Assure II (page 65). Information listed is restricted to Crop, Weeds, Rates, and Cost. For other detailed information on the component products see the product pages listed above.

Company:

E. I. duPont Canada

Formulations:

Muster Gold II contains 3 components:

Muster (PCP#23569): 75% ethametsulfuron-methyl as a water dispersible granule. Container size - 4 x 80 g water soluble bags.

Assure II (PCP#25462): 96 g/L quizalofop-p-ethyl as an emulsifiable concentrate. Container size - 8 L.

Merge adjuvant (PCP#24702). Container size - 8 L

Crops and Staging:

Canola, including all herbicide tolerant canola varieties.

2 leaf stage to the beginning of bolting. If application is made prior to this stage, severe crop injury may occur.

Weeds and Staging:

Grasses:

Apply at the 2 leaf to early tillering stage unless otherwise noted. For best results on volunteer cereals & wild oats apply before tillering begins

Barnyard grass
Fall pannicum
Green and yellow foxtail
Old witchgrass
Proso millet
Quackgrass* (2 to 6 leaf)
Volunteer cereals
Volunteer corn
Wild oat

Broadleaf weeds:

Apply at the cotyledon to 6-leaf stage unless otherwise noted.

Flixweed
Hemp-nettle
Redroot pigweed*
Stinkweed (1 to 4 leaf)
Wild mustard
Smartweed/lady's-thumb

*Suppression only

Cost (2004 suggested retail price):

$21.25 per acre

Rates:

Muster: 8 g per acre.

Assure II: 0.2 L per acre

Merge adjuvant: 0.2 L per acre (as packaged)

A case of Muster Gold II will treat 40 acres (16 ha).

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 30 to 40 psi (210 to 275 kPa).

Nozzles: Flat fan nozzles with 50 mesh line strainers and screens.

Muster Toss-N-Go

Herbicide Group – 2
(Refer to page 27)

Company:

E. I. duPont Canada (PCP#23569)

Formulation:

75% ethametsulfuron-methyl formulated as a dry flowable.

Container sizes – 320 g package (containing 4 x 80 g water soluble bags).

Crops and Staging:

Canola: From the 2 leaf stage (main stem leaves) to the start of bolting. Do not apply prior to this stage as severe crop injury can occur.

Condiment mustard (brown & oriental only): From the 4 leaf stage but prior to bolting. Do not apply prior to this stage as severe crop injury can occur. Spraying must take place at least 60 days prior to harvest.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Apply from the 2 to 6 leaf stage. Stinkweed must be sprayed in the 1 to 4 leaf stage

At the 8 g/acre rate (one 320 g package treats 40 acres):

Flixweed *
Green smartweed
Hemp-nettle
Stinkweed **
Wild mustard

The 12 g/acre rate (one 320 g package treats 26.7 acres)*** controls above weeds plus:

Stinkweed
Redroot pigweed **

* Spring seedlings only

** Suppression with Muster alone but control with Assure II plus Sure-Mix or a Poast Ultra + Merge tank mix.

*** High rate is for use in canola only. DO NOT use high rate on condiment mustard crops.

Muster applied alone requires the addition of Agral 90, Agsurf, or Citowett at 0.2 L per 100 L of spray solution. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$16.27 to $24.37 per acre.

Application Information:

Water Volume: 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Equipment and Nozzles: Flat fan nozzles. Sprayer must be equipped with continuous agitation. Use 50 mesh or coarser strainers and screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not use on crops that are stressed because of drought or flooding. Less than acceptable control will occur in fields where high weed populations exist and where stressful environmental conditions prevail (drought, cold weather). Heavy rainfall soon after application may result in visual crop injury or possible yield reduction. Thin crop stands or application prior to the 2 leaf stage, sandy soils or soils with low organic matter may increase the severity of the injury.

Tank Mixes:

Do not mix with substances that contain boron or that release chlorine.

Herbicides:

Canola only: Assure II, or Poast Ultra. The adjuvants used with these products are all that are required for tank mixes.

Fertilizers: None registered. Do NOT mix soluble bags with liquid fertilizers.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Muster label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 4 to 6 hours may reduce control.

Grazing: Do not graze or feed crop to livestock within 60 days of application.

Preharvest: Leave at least 60 days from application to harvest.

Recropping: Do not sow wheat, barley, oats or flax within 10 months of application. Do not sow canola, lentils, peas, fababeans, tame mustard, alfalfa, canaryseed, dry beans, fescues or red clover within 22 months of application. All other crops must not be sown until a "field bioassay" is performed at 22 months (or more) after application. Growers may experience reduced yields if other crops (such as corn) are grown without following these guidelines.

Aerial Application: Do not apply by air.

Storage: May be frozen.

Environment: Leave a 15 m buffer between the last sprayer pass and wetland areas or waterways. Leave a 5 m buffer between the last spray pass and woodlots, shelterbelts, ditch bank vegetation and the borders of dry sloughs.

Tank Cleaning:

Sprayers used to spray Muster should be flushed out immediately after Muster is used. The manufacturer recommends that sprayers used to apply this product be flushed twice with a water/household ammonia rinse (1L of 3 % ammonia per 100 L water). All nozzles, screens and filters should be removed and cleaned after applying this product.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 14 for additional information.

Hazard Rating:

▽ Caution Poison

Odyssey

Herbicide Group – 2
(Refer to page 27)

Company:

BASF Canada (PCP#25111)

Formulation:

35% imazamox and 35% imazethapyr formulated as a dispersible granule.

Container size - 4 x 692 g water soluble bags

Crops and Staging:

Field peas - 1 to 6 above-ground nodes (1 to 6 true leaves).
CLEARFIELD canola - 2 to 6 leaf stage.

Temporary crop yellowing may be observed shortly after application in field peas and CLEARFIELD canola.

Forage Crops (seed production only)
Alfalfa (seedling or established) – 1 to 4 leaf stage

Weeds, Rates and Staging:

At 12 g/acre (60 acres per case), Odyssey will control:

Redroot pigweed
Wild mustard
Volunteer tame mustard

At 17 g/acre (40 acres per case), Odyssey will control:

Grasses - 1 to 4 main stem leaves, until tillers are visible:

Barnyard grass
Green foxtail
Persian darnel
Volunteer cereals (wheat, barley, oats)
Wild oat

Broadleaf Weeds - cotyledon to 4 leaf stage unless otherwise indicated:

Chickweed
Cleavers (up to 4 whorls)
Flixweed
Green smartweed
Hemp-nettle*
Kochia**
Lamb's-quarters**
Redroot pigweed
Russian thistle
Shepherd's-purse
Stinkweed
Stork's-bill
Volunteer canola (not CLEARFIELD varieties)
Volunteer tame mustard
Wild buckwheat*
Wild mustard

* Suppression only in field peas.
** Suppression only in field peas and CLEARFIELD canola.

Merge surfactant (not included) must be used with Odyssey.

CLEARFIELD Canola and Peas: Use only Merge surfactant. Mix to a final ratio of 0.5 L of Merge in 100 L of spray solution. (0.5% Merge volume per spray solution volume).

Alfalfa: Apply at 40 L/acre of water with Merge or Amigo adjuvants alone at a rate of 0.5 L adjuvant per 100L spray solution (0.5% adjuvant volume per spray solution volume) or with a non-ionic surfactant (Agral 90, Agsurf or Surf92) and a fertilizer solution (28-0-0 or 32-0-0) at a rate of 0.25 L non-ionic surfactant per 100 L spray solution plus 0.8 L/acre fertilizer.

Do not apply Odyssey more than once per year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Cost (2004 suggested retail price):

$16.61 to $24.92 per acre (including Merge).

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan. Use 50 mesh or coarser filter screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

No information provided on the product label. Plants are under stress when conditions before or after application are very hot or cold, or excessively dry or wet. During periods of stress, plants are not actively growing and reduced weed control or crop injury may result.

Tank Mixes:

Herbicides:
In CLEARFIELD canola only:
Lontrel (0.17 - 0.23 L/acre).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Odyssey label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rainfall within 3 hours of application may reduce control.

Grazing: Do not graze treated crop or cut for feed prior to crop maturity. Field peas may be fed to livestock 30 days after application. Do not harvest forage or cut for hay.

Preharvest Interval: Do not apply within 60 days of harvest.

Recropping: Field peas, lentils, CLEARFIELD canola, canaryseed, oat, barley and spring wheat (including durum) may be seeded the first spring after application. Flax may be seeded the second spring after application. The company recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed above.

NOTE: Breakdown of Odyssey may be slowed or delayed by environmental conditions such as drought, excessive cold and/or acid soils (pH less than 6.5) resulting in an increased risk of injury to rotational crops. The most tolerant crops are CLEARFIELD canola and legume crops, then cereals. Contact manufacturer for additional information on recropping intervals.

Aerial Application: Do not apply by air.

Storage: Do not freeze. Store in a cool, dry place above 5° C.

Environment: Avoid spraying in situations where drift may occur. Leave a buffer zone of at least 14 m between the outside boundary of the sprayed area and sensitive areas such as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Warning Eye and Skin Irritant

Pardner/Koril 235

See Bromoxynil section on page 79

Pea Pack*

Herbicide Group – 4, 5

(Refer to page 27)

This product is a prepackaged tank mix of Sencor (page 184) and MCPA Na (page 151). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Bayer CropScience

Formulation:

The Pea Pack* case contains two components:

Sencor 75DF (PCP#25840): 75% metribuzin formulated as a dispersible graule. Container size - 3 kg.

MCPA sodium salt (PCP#14718): 300 g/L MCPA sodium salt formulated as a solution. Container size - 7.6 L.

*** Pea Pack is no longer manufactured but quantities still remain in the distribution system. This product may be removed from this guide in future editions.**

Crops and Staging:

Field peas before vines are 6 inches (15 cm) long.

DO NOT USE ON LENTILS OR CHICKPEAS.

Do not use on peas seeded less than 2 inches (5 cm) deep or on soils with less than 4 percent organic matter.

Weeds and Staging:

Apply when weeds are small i.e. less than 2 inches (5 cm) in height or diameter.

Lamb's-quarters
Redroot pigweed
Stinkweed
Volunteer canola (including all herbicide-tolerant canola cultivars),
Wild mustard.

Cost (2004 suggested retail price):

$5.95 per acre.

Rate:

77 g Sencor DF and 189 mL MCPA sodium salt per acre. Each case will treat 40 acres.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 70 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa)

Nozzles: Flat fan nozzles with an opening no smaller than 8002 or TK2 with 50 mesh screen. Angle nozzles 45° forward.

Pinnacle

Herbicide Group – 2
(Refer to page 27)

Company:

E. I. duPont Canada (PCP#22002)

Formulation:

75% thifensulfuron methyl as a water dispersible granule. Container size – 8 x 8 g water soluble pouches

Crops and Staging:

Soybeans - First fully expanded trifoliate leaf to flower initiation.

Weeds and Staging:

Apply up to 4 inches (10 cm) tall or wide:

Lady's-thumb
Lamb's-quarters*
Redroot pigweed
Wild mustard
Velvetleaf*

* High rate of application only

Cost (2004 suggested retail price):

$ 9.47 to $13.50 per acre.

Rates:

2.2 to 3.2 g/acre (one container treats 20 to 28.5 acres).

Requires the addition of a non-ionic surfactant such as Agral 90, Agsurf, or Citowett at 1 L per 1000 L of spray solution. Oil surfactant blends such as Assist, CanPlus 411 or Sure-Mix may be used as adjuvants (check label for use rates).

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum of 45 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat Fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Pinnacle applied to crops that have been under stress before application may result in crop injury. Stress conditions within 3 days after application may also result in crop injury.

Weeds under stress conditions at the time of application may not be adequately controlled.

Stress conditions are severe weather conditions, frost, low fertility, drought, water-saturated soils, and disease or insect damage.

Injury symptoms can be crop discoloration (yellowing, purpling or reddening), or stunting.

Tank Mixes:

Herbicides:

Assure II (0.2 L/acre) plus Sure-Mix.

Basagran or Basagran Forte (0.71 to 0.91 L/acre) plus Assist adjuvant.

Assure II (0.25 L/acre) plus Basagran Forte (0.71 to 0.91 L/acre) plus Sure-Mix or CanPlus 411 adjuvant.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Pinnacle label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: If rainfall occurs soon after application control may be reduced. Several hours of dry weather are needed after application to allow uptake by the plants.

Grazing: Do not graze or cut for feed.

Pre-harvest: Leave at least 60 days from application to harvest.

Recropping Interval: Do not plant any crop other than wheat or barley for 30 days after application.

Aerial Application: Do not apply by air.

Storage: Store in closed original container in a dry area away from food or feed.

Environment: Do not mix, load or apply within 15 m of water bodies, wetlands or sensitive plants. Do not apply to irrigated land where the drainage water is also used to irrigate cropland.

Tank Cleaning:

Pinnacle can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray Pinnacle should be flushed out immediately after use. The manufacturer recommends that sprayers used to apply this product be flushed twice with a water/household ammonia rinse (1 L of 3 percent ammonia per 100 L of water). All nozzles, screens and filters should be removed and clean separately during the cleaning process. Do a clean water rinse of the tank, booms, hoses and nozzles to complete the procedure.

Refer to page 14 for additional information.

Hazard Rating:

Warning – Eye and Skin Irritant

Poast Ultra

Herbicide Group – 1

(Refer to page 27)

Company:

BASF Canada (PCP#24835)

Formulation:

450 g/L sethoxydim formulated as an emulsifiable concentrate.

Container size – 2 x 7.7 L jug of Poast Ultra.

Crops, Rates and Staging:

Crops are tolerant at all growth stages. However, the preharvest interval outlined in the "Restrictions:" section must be followed to avoid unacceptable residues of sethoxydim in harvested crops.

To a maximum rate of 0.45 L per acre:

Alfalfa
Alsike clover*
Canola
Chickling vetch
Cicer milkvetch*
Creeping red fescue (for seed only)
Dry beans (adzuki, kidney, lima, mung, pinto, white)
Dry field peas
Fababeans
Flax (not including Solin)
Lentil
Mustard
Potatoes
Sainfoin*
Shelterbelts
Soybeans
Sunflower
Sweet clover*

To a maximum rate of 0.26 L per acre:

Alsike clover**
Caraway
Cicer milkvetch**
Coriander
Dill
Safflower
Sainfoin**
Solin (low linolenic flax)
Sweet clover**

To a maximum rate of 0.23 L per acre:

Tame buckwheat

To a maximum rate of 0.19 L per acre:

Chickpea

* Established stands
** Seedling stands

Weeds, Rates and Staging:

Optimum yield response occurs when weeds are controlled early.

WEEDS AND STAGES	RATE (L/ACRE)	ACRES TREATED PER 7.7 L CONTAINER
Green or yellow foxtail, barnyard grass, volunteer corn, Persian darnel, proso millet, witchgrass, large crabgrass and fall panicum - 1 to 6 leaf stage	0.13	60
Wild oats, Volunteer wheat, oats and barley - 1 to 6 leaf stage except for low rate (See footnote)	0.13* or 0.19	60 or 40
Quackgrass suppression - 1 to 3 leaf stage	0.19	40
Quackgrass (season long control) - 1 to 3 leaf stage	0.45	17
Foxtail barley suppression - prior to tillering.	0.45	17

*Use the low rate in canola, flax and peas when wild oat, volunteer wheat and volunteer barley are from 1 to 4 leaves (best results prior to tillering) and under ideal growing conditions (adequate moisture, good fertility and moderate temperatures 15 to 28°C). Do not apply under stress conditions.

A surfactant must be used with Poast Ultra when applied alone or in tank-mix with a broadleaf herbicide. Use only Merge surfactant. Mix to a final ratio of one jug of MERGE in 1000 L of diluted product solution (or 0.5 to 1.0 % Merge volume/spray solution volume).

Add one 8.1 L jug of Merge and apply in 40 L per acre of spray solution for 20 acres to 40 acres.

Add one 8.1 L jug of Merge and apply in 20 L per acre of spray solution for 40 acres to 80 acres.

When applying at water rates greater than 40 L per acre, mix and apply 0.4 L of MERGE per acre.

When applying to areas where quackgrass and/or foxtail barley is present:, mix and apply 0.4 L to 0.8 L MERGE* per acre. An 8.1 L jug of MERGE* will cover 10 to 20 acres.

Cost (2004 suggested retail prices):

$11.03 to $38.94 per acre.

Application Information:

Water Volume: Apply at water rates of 20 to 40 L/acre Use a minimum of 40 L/acre if crop or weed growth is dense, and when spraying quackgrass.

Use 20 to 40 L/acre when using the 0.19 L/acre rate for wild oats, volunteer wheat and volunteer barley.

Pressure: 40 to 45 psi (275 to 300 kPa).

Nozzles: Stainless steel 80° or 110° flat fan nozzles tilted forward at an angle of 45°.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Most effective control is achieved when grasses are actively growing. Weeds stressed by drought, flooding, hot or prolonged cool temperatures (<15°C) and poor fertility are more difficult to control. Use the higher of the recommended rates for grasses stressed for less than 20 days. Do not apply to grasses stressed more than 20 days because of lack of moisture. Control may be reduced if temperatures are below 15°C. Retillering may occur under stress conditions or if fertility is low.

Tank Mixes:

Herbicides: The following tank mixes can be applied with 0.13 to 0.19 L/acre of Poast Ultra. Include Merge at 0.3 to 0.4 L/acre with all mixes.

In Flax:
Buctril M (0.405 L/acre) (including Solin)
Lontrel 360 (0.23 to 0.34 L/acre)
Lontrel 360 (0.23 to 0.34 L/acre) + MCPA Ester (0.34 to 0.45 L/acre of 500 g/L formulations)
MCPA Ester (up to 0.45 L/acre of 500 g/L formulations)
The above tank mixes may reduce grass control, especially under adverse weather conditions.

In Canola:
Lontrel (0.17 to 0.34 L/acre)
Muster (8 to 12 g/acre)
Lontrel (0.17 L/acre) + Muster (8 g/acre)

In Liberty Link canola only:
Poast Ultra (0.09 L/acre) + Liberty (1.08 L/acre)

In Field peas and CLEARFIELD canola:
Poast Ultra (0.19 L/acre) plus Merge (0.4 L/acre) may be tank mixed with the following herbicides:

Pursuit (40 mL/acre) to control: Chickweed, stinkweed, cleavers, volunteer canola (non-CLEARFIELD varieties), hemp-nettle (peas only), redroot pigweed (light infestations only), wild buckwheat (light infestations only) smartweed, wild mustard.

The company does not provide guidelines for weed densities under light infestations. When in doubt, use the higher rate below or contact the manufacturer.

Pursuit (85 mL/acre) for all weeds on the Pursuit label. Check label directions for mixing order and additional timing restrictions for broadleaf partners.

Allow 4 days between application of Poast Ultra and application of herbicides other than those registered for tank mixing. Allow 5 days between application of Sencor and Poast Ultra. Allow 14 days for regrowth when applied in sequence with a grass control herbicide.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Poast Ultra label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Grazing: Do not graze the treated crop or cut for feed prior to crop maturity. Forage legumes may be cut after the specified preharvest interval.

Preharvest Interval: Forage legumes - 30 days; Dry peas & flax – 60 days; Lentil & chickpea - 65 days; Canola, chickling vetch & alfalfa – 70 days; Mustard – 76 days; Potatoes, dry beans, soybeans, & fababeans – 80 days; Buckwheat - 85 days; Solin – 86 days; Safflower – 90 days; Sunflower 105 days.

Recropping: Do not sow cereal or grass crops within 14 days of Poast Ultra application.

Aerial Application: May be applied by air in 10 to 20 L/acre of water. Merge should be added at rates of 0.10 to 0.20 L/acre.

Storage: May be frozen.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Caution Poison

PrePass

Herbicide Group – 2,9
(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

Each case of PrePass contains 2 components:

PrePass A (PCP#27395): 50 g/L florasulam formulated as a suspension concentrate.

PrePass B (PCP#27394): 360 g/L glyphosate IPA salt formulated as a solution.

Container sizes: PrePass A - 1.6 L, PrePass B - 2 x 10 L.

Crops and Staging:

Prior to seeding of spring wheat (including durum), barley and oats.

Weeds Controlled and Staging:

Grasses at the 1 to 4 leaf stage:

Downy brome
Green foxtail
Persian darnel
Volunteer barley
Volunteer wheat
Wild oat

Broadleaf weeds controlled at the 1 to 4 leaf stage:

Canada fleabane**
Common chickweed
Cleavers
Dandelion seedlings
Flixweed
Hemp-nettle
Lady's-thumb
Lamb's-quarters
Narrow-leaved hawk's-beard**
Ragweed (common)**
Redroot pigweed
Russian thistle
Shepherd's-purse
Stinkweed
Volunteer canola (all varieties)
Volunteer flax
Wild buckwheat*
Wild mustard

Broadleaf weeds suppressed:

Annual sow-thistle
Kochia
Perennial sow-thistle***

* up to 5 leaf stage
** up to 8 cm in height
*** better results from earlier applications

Cost (2004 suggested retail price):

$8.45 per acre.

Rate:

PrePass A: 0.04 L (40 mL)/acre
PrePass B: 0.5 L/acre

One case treats 40 acres.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13 of the Guide to Crop Protection.

Application Information:

Water Volume: 20 to 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

PrePass A: Warm, moist growing conditions promote active weed growth and enhance activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur.

PrePass B: Best results are achieved when temperatures are near 20°C and when weeds are actively growing. Frost that kills more than 40% of above ground tissue will reduce control. Heavy dust layer on leaves will also reduce control.

Tank Mixes:

None registered.

Restrictions:

Rainfall:
PrePass A: No rainfast period is specified for this product; required interval may be up to 8 hours.
PrePass B: Heavy rainfall immediately after application may wash the chemical off the foliage. Do not apply if rainfall is forecast for the time of application. Contact manufacturer for more information.

Grazing: Do not graze treated crop or cut for feed within 7 days of application.

Pre-harvest: Leave 60 days between application and harvest.

Recropping: Spring wheat (including durum), barley and oat, may be seeded after application or the field may be fallowed. Barley, canola, field peas and wheat, may be grown the following season (10 months following an application).

Aerial Application: Do not apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Environment: Leave a buffer zone of at least 30 m between the downwind edge of the boom and the closest edge of sensitive terrestrial areas such as shelterbelts, hedgerows and shrub lands. A buffer zone of at least 5 m is required between the downwind edge of the boom and the closest edge of aquatic areas such as ponds, streams, rivers, prairie potholes and sloughs.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

May cause eye irritation

Prestige

Herbicide Group – 4

(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

Prestige A (PCP#25465): 180 g/L fluroxypyr. Container size – 6.4 L.

Prestige B (PCP#25464): 50 g/L clopyralid and 280 g/L MCPA ester. Container size – 2 x 8.0 L.

Both are formulated as an emulsifiable concentrate.

Crops and Staging:

Spring wheat (including durum), barley and canaryseed* from the 3 leaf stage up to the time just prior to the emergence of the flag leaf, tall fescue (established)* grown for seed at 4 leaf to flag leaf stage.*

* *Since these uses of Prestige are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance.* ***Users of this product on tall fescue and canary seed do so at their own risk.***

Weeds and Staging:

Unless otherwise stated, the following weeds will be controlled if sprayed in the 2 to 4 leaf stage.

Annual sow-thistle
Canada thistle*
Chickweed**
Cleavers (1-4 whorls)
Common groundsel
Dandelion***
Flixweed***
Hemp-nettle (2-6 leaf stage)**
Kochia
Lamb's-quarters
Perennial sow-thistle*
Redroot pigweed
Round leaved mallow (1 to 6 leaf)
Russian pigweed
Scentless chamomile
Shepherd's-purse
Smartweed
Stinkweed
Stork's-bill (1-8 leaf)
Tartary buckwheat
Volunteer canola
Volunteer flax (1-12 cm)
Volunteer sunflower
Wild buckwheat (1-4 leaf)
Wild mustard

* Spray when 4 to 6 inches (10 to 15 cm) high. Season long control, with some regrowth in the fall.

** Suppression only. Spray as late as possible (within recommended stages) to ensure the majority of weed flushes have emerged.

*** Spring rosettes only.

Cost (2004 suggested retail price):

$14.70 per acre.

Rates:

Prestige A: Apply at 0.32 L per acre

Prestige B: Apply at 0.8 L per acre

One case of Prestige will treat 20 acres.

Application Information:

Water Volume: 40 L per acre

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan nozzles tilted forward at a 45° angle.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

The activity of the Prestige is influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought or heat stress) or if heavy infestations exist.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Achieve Liquid (200 mL/acre) plus Turbocharge adjuvant
Assert (0.53 to 0.65 L/acre) plus acidifier
Puma® Super (0.16 to 0.32 L/acre)

In spring wheat (including durum):

Horizon (95 to 115 mL/acre) plus Score adjuvant
Everest (17.4 g/acre) plus recommended surfactant

Fertilizers: None registered.

Insecticides: None registered.

Note: The above mixes are those listed on the Prestige label only. For other possible mixes see the blue fold out chart inside back cover.

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours of post-emergent application may result in reduced weed control.

Grazing: Do not cut or graze treated fields of wheat or barley for 7 days after application. Do not cut treated tall fescue fields for hay or forage. Do not graze treated tall fescue fields.

Preharvest Interval: Do not harvest crop within 60 days of application.

Recropping: Wheat, oats, barley, rye (not under-seeded to forage legumes, clover or alfalfa), flax, canola, field peas* and mustard may be seeded the season following Prestige application.

*NOTE: Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Do not seed legume forages or crops other than those listed above until the second season following the Prestige application.

Aerial Application: Do not apply by air.

Storage: Store product in original containers in a secure, dry, heated area. If the product is frozen, bring to room temperature and agitate before use.

Environment: This product contains elements that are toxic to aquatic organisms and can damage susceptible non-target plants. Leave a 15 metre buffer between the nearest sprayer pass to non-target areas and domestic, irrigation or natural water or other wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Danger Poison.

Warning Eye Irritant.

Caution Skin Irritant.

Prevail Liquid

Herbicide Group – 1, 4
(Refer to page 27)

This product is a prepackaged tank mix of Prevail A (Achieve page 54) and Prevail B (Curtail M page 86). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products see the product pages listed above.

Company:

Dow AgroSciences.

Formulation:

The Prevail package contains 3 components:

Prevail A (PCP#27579): 400 g/L tralkoxydim formulated as a suspension concentrate. Container size - 4 L.

Prevail B (PCP#23526): 50 g/L clopyralid plus 280 g/L MCPA ester formulated as an emulsifiable concentrate. Container size - 8 L.

Prevail C (PCP#23527): adjuvant (Turbocharge). Container size - 4 L.

Crops and Staging:

Spring wheat (including durum) and barley at the 3 to 6 leaf stage.

Weeds and Staging:

Grasses:

Wild oats - 1 to 6 total leaves,to a maximum of 2 tillers.
Green foxtail - 1 to 5 total leaves, to a maximum of 1 tiller.
Persian darnel - 1 to 4 total leaves including tillers.
Optimum weed control and yield response occurs when weeds are removed before tillering.

Broadleaves:

Apply at the 1 to 4 leaf stage unless otherwise indicated

Annual sow-thistle
Canada thistle*
Common groundsel
Dandelion***
Flixweed***
Kochia (suppression)[1]
Lamb's-quarters
Perennial sow-thistle**
Redroot pigweed
Russian pigweed
Scentless chamomile[1]
Shepherd's-purse***
Smartweed
Stinkweed***
Tartary buckwheat
Wild buckwheat
Wild mustard
Volunteer canola

* Season long control, some regrowth may occur in the fall. Apply after all the thistles have emerged and are between 4 inches (10 cm) and 6 inches (15 cm) tall but prior to early bud stage.
** Top growth control only.
*** Spring rosettes only.
[1] 2 to 4 leaf stage.

Cost (2004 suggested retail price):

$25.95 per acre.

Rate:

The Prevail package will treat 20 acres.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° flat fan nozzles. All strainer and nozzle screens must be 50 mesh or coarser.

Hazard Rating:

Caution Poison

Warning : Eye and Skin Irritant

Primextra II Magnum

Herbicide Group – 15, 5

(Refer to page 27)

Company:

Syngenta (PCP#25730)

Formulation:

400 g/L of s-metolachlor + 313 g/L of atrazine formulated as a liquid. Container size - 14 L.

Crops and Staging:

Corn - Preplant incorporated (PPI) or Pre-emergence if irrigated within 10 days of application.

Weeds and Staging:

Apply prior to the emergence of weeds. Weeds that have emerged prior to application will not be controlled.

American nightshade
Barnyard grass
Buckwheat
Eastern black nightshade
Green foxtail
Lady's-thumb
Prostrate pigweed
Purslane
Ragweed
Redroot pigweed
Smartweed
Wild mustard
Yellow foxtail
Yellow nutsedge*

* Herbicide must be incorporated for best control.

Cost (2004 suggested retail price):

$22.03 to $29.29 per acre.

Rates:

WEED POPULATIONS	RATE (L/ACRE)	ACRES TREATED PER 14 L CONTAINER
Light infestations	1.2	11.7
Medium infestations	1.4	10
Heavy infestations	1.6	8.8

Application Information:

Water Volume: 61 L per acre

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan with 50-mesh nozzle screens.

Incorporation:

Incorporate using S-tine or C-tine cultivators or tandem disk. Do not incorporate deeper than 4 inches (10 cm).

To ensure that the product remains in the top 2 inches (5 cm) of soil, apply to a firm seedbed free of large clods or lumps. If using tandem disks, set disks to work the soil at a depth of 4 inches (10 cm) and operate at a speed of 4 mph (6 km/hr). If using an S-tine cultivator, set the implement to work the soil to a depth of 4 inches (10 cm) and operate at a speed of 6 mph (10 km/hr).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Extended periods of dry soil conditions may result in reduced weed control. Moderate rainfall after application will enhance activity. Heavy rainfall following application of Primextra II Magnum may dilute the metolachlor deeper than 2 inches (5 cm) and result in reduced weed control, particularly on light textured soils.

Tank Mixes:

Herbicides: None registered

Fertilizers: May be tank mixed with liquid fertilizer for preplant incorporated applications. Conduct a compatibility test by performing a jar test prior to mixing the products in the tank. May be impregnated onto dry bulk fertilizers (except nitrate or superphosphate fertilizers).

Insecticides: None registered.

Note: The above mixes are those listed on the Primextra label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Moderate rainfall shortly after application will enhance activity. Heavy rainfall reduces weed control by leaching the chemical out of the top few centimeters of soil. Inadequate rainfall after application (within 10 days) will cause reduced weed control.

Grazing: Do not graze or cut corn for feed before ear emergence.

Recropping: This product contains Atrazine. All crops except corn and triazine-tolerant canola may be affected the year following the use of Atrazine. Other more sensitive crops may be affected two or more growing seasons after application.

Aerial Application: Do not apply by air.

Storage: Store in a dry place.

Environment: Do not mix or load this product within 30 m, or apply within 29 m, of any wells, lakes, streams, ponds, dugouts or sink holes.

Tank Cleaning:

Refer to page 14.

Hazard Rating

Keep out of reach of children.

Prism

Herbicide Group – 2

(Refer to page 27)

Company:

E. I. duPont Canada Inc.

Formulation:

25% rimsulfuron formulated as a dry flowable.
Container size - 480 g.

Crops and Staging:

Irrigated potatoes prior to flower initiation.

Potato tolerance to Prism differs by variety. Limit first use of Prism to a small area of each variety prior to adoption as a field practice. Delay cultivation for 7 to 10 days after application.

Weeds and Staging:

WEED	STAGE
Barnyard grass Green foxtail Yellow foxtail Witchgrass	1 to 6 leaf stage, maximum 2 tillers
Quackgrass	3 to 6 leaf stage (less than 10 inches or 25 cm leaf extended).
Redroot pigweed Lamb's-quarters (suppression only)	4 to 6 leaf stage (less than 4 inches or 10 cm tall or across).

Cost (2004 suggested retail price):

$20.31 per acre (plus adjuvant).

Rates:

24 g/acre (one package treats 20 acres).

Add a recommended non-ionic surfactant such as Citowett Plus, Agsurf, or Agral 90 at 0.2 L per 100 L spray solution. Make only one application per growing season.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 25 to 40 psi (175 to 275 kPa).

Nozzles: Flat fan. 50 mesh screens or coarser.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Apply Prism when the temperature 24 hours before and after application is between 5°C and 28°C. Temperatures beyond this range increase the potential for crop injury. Rapid fluctuations in temperature will stress the crop (greater than a 20°C difference within 24 to 36 hours). Allow 48 to 72 hours for the crop to acclimatize before spraying Prism if severe temperature fluctuations occur. Crop injury may result if applications are made when potatoes are stressed by abnormally hot, humid, or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If potatoes have been injured by frost, wait 48 to 72 hours after normal growing conditions have resumed before applying Prism.

Warm, moist conditions after application promote good weed control with Prism while cool and/or dry conditions may reduce or delay activity. Weeds hardened off by cold weather or drought stress may not be controlled.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 2 to 4 hours of application may reduce weed control.

Preharvest: Leave at least 30 days from application to harvest.

Recropping: Spring barley, soybeans, white beans, red clover, sorghum, potatoes and field corn may be planted the year after application. Winter wheat may be planted 4 months after application. For all other crops, a field bioassay is recommended before planting.

Aerial application: Do not apply by air.

Storage: May be frozen.

Environment: Do not apply in areas where surface water from the treated area can run off to adjacent cropland, streams, irrigation water or wells. Leave a buffer of 10 m from the last spray pass and wetland areas or bodies of water. Leave a buffer of 5 m from the last spray pass and woodlots or shelterbelts.

Tank Cleaning:

Do not clean equipment near desirable trees or other plants. Do not contaminate water sources. Drain tank, hose down tank interior and flush clean water through hoses for a minimum of 5 minutes. Create a water-ammonia cleaning solution (1 L of minimum 3% household ammonia per 100 L water) in spray tank. Flush hoses and nozzles, and add water to fill tank. Circulate cleaning solution for 15 minutes, and flush through boom again. Wash nozzles and screens in cleaning solution in pail. Repeat above tank and hose cleaning procedure. Rinse system with clean water for 5 minutes. Prior to using sprayer for next application, flush system with fresh water for 5 minutes.

Refer to page 14 for additional information.

Hazard Rating:

Warning Eye Irritant

Puma^{120} Super

Herbicide Group – 1

(Refer to page 27)

Company:

Bayer CropScience (PCP#25864)

Formulation:

120 g/L fenoxaprop-p-ethyl formulated as an emulsifiable concentrate. Container sizes – 6.2 L, 99.3L

Crops and Staging:

Spring wheat (including durum) with 1 to 6 leaves on the main stem plus 3 tillers

Barley[†] with 1 to 5 leaves on the main stem plus 2 tillers.

[†] **Apply to barley only when tank mixed with a registered broadleaf product.**

Durum wheat and barely may experience some initial, temporary stunting and yellowing that rarely results in yield loss. Injury is more likely under stress conditions (see Effect of Growing Conditions).

Treatment at the 3 to 4 leaf stage of crops and weeds will maximize crop tolerance and weed control. Temporary crop injury such as shortening or discolouration may be observed after application. Such injury is more likely to occur in barley and also when Puma® Super is applied outside recommended stages.

Weeds, Rates and Staging:

WEEDS	RATE (L/ACRE)	AREA (ACRES) PER 6.2 L	STAGE†
Green foxtail only	0.154	40	1 to 6 leaf stage up to emergence of 3rd tiller
Low wild oat infestations*	0.268	23.1	
Moderate-heavy wild oat infestations, barnyard grass, green and yellow foxtail	0.31	20	

† Apply at the 3 to 4 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

* Low wild oat rate for use on Wheat and Durum only, and when applied alone and not in a tank-mix.

Cost (2004 suggested retail price):

$8.46 to $16.92 per acre

Application Information:

Water Volume:

Ground application: 23 to 45 L per acre.

Aerial application: A minimum of 14 L per acre.

Pressure:

Ground application: 40 psi (275 kPa)

Aerial application: 42 psi (300 kPa)

Nozzles:

Ground application: 80° or 110° flat fan nozzles tilted forward at a 45° angle. Do not use flood jet nozzles, controlled droplet application equipment or Sprafoil equipment.

Aerial application: Use a combination of nozzles and pressure for a coarse droplet size distribution (400-800 microns).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

DO NOT apply Puma® Super 2-3 days prior to, or following, temperatures of 3°C or lower as crop injury may occur. Under stressful conditions (hot/dry, waterlogging, disease or insect damage) or heavy crop canopy, early application will improve weed control.

Do not apply by air when both the temperature is greater than 25°C and the relative humidity is less than 30%.

Tank Mixes:

Herbicides:

In Spring Wheat, Durum Wheat and Barley:

NOTE: Do NOT apply Puma® Super alone in barley. ALWAYS tank mix with a registered broadleaf herbicide.

2,4-D Ester LV 600 (0.28 L/acre), LV 700 (0.24 L/acre)
Ally (2 to 3 g/acre)
Buctril M (0.4 L/acre)
Buctril M (0.40 L/acre) + Refine Extra (2.7 g/acre)
Curtail M (0.6*** to 0.8 L/acre)
Dichlorprop + 2,4-D (0.71 L/acre)
DyVel** (0.5 L/acre)
Express Pack** (40 acres/case rate)
Frontline** (40 acres/case rate)
MCPA Amine or Ester (0.34 L/acre)
Prestige (20 acres/case rate)
Refine Extra (8 g/acre)
Refine Extra (8 g/acre) + MCPA Ester (0.34 L/acre)
Spectrum** (20 acres/case rate)
Thumper (0.40 L/acre)
Trophy* (20 acres per case)
Unity **** (40 acres/case rate)

In Spring and durum wheat only:

Attain (40 acres/case rate)
DyVel** DS (0.44 L/acre)
Lontrel 360 (0.17 L/acre)
Lontrel 360 (0.17 L/acre) + MCPA Ester (0.34 L/acre)*
Lontrel 360 (0.112 L/acre) + MCPA Ester (0.34 – 0.45 L/acre)*
Mecoprop (2.2 to 2.8 L/acre)*

* Use only at the high rate of Puma® Super
** Use only at the green foxtail rate of Puma® Super
*** Low Curtail M tank-mix rate for use in Barley only.
**** Not registered for use in Manitoba.

Fertilizers: None registered

Insecticides: None registered

Fungicides: None registered.

Note: The above mixes are those listed on the Puma^{120} Super label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Leave an interval of 7 days prior to application or 4 days after application of Puma^{120} Super, when applying any pesticide that is not registered as a tank mix.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Grazing: Do not graze or cut for hay, within 25 days of application

Preharvest Interval: Do not harvest within 65 days of application.

Recropping: No restrictions in the year after application. Only one application may be made per year.

Aerial Application: May be applied by air.

Storage: Do not freeze.

Environment: Leave 3 metres from the downwind edge of the spray boom to water or wetland areas by ground or air. Leave 10 metres by ground and 33 meters by air from the downwind edge of the boom to sensitive non-target land areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison.

Eye and Skin Irritant.

Pursuit

Herbicide Group – 2

(Refer to page 27)

Company:

BASF Canada (PCP#23844)

Formulation:

240 g/L imazethapyr formulated as a liquid.

Container size - 2 x 3.3 L jugs per case.

Crops and Staging:

CROP	STAGE
CLEARFIELD canola	Apply after the first leaf stage.
Dry beans (pinto, pink and red varieties only)	Up to and including the second trifoliate leaf stage
Soybeans (Manitoba only)	Up to and including the third trifoliate leaf stage
Field peas	May be applied up to the sixth above-ground node stage (6 true leaves).
Seedling alfalfa (forage or seed production)[A]	Apply after the first trifoliate leaf stage.
Established alfalfa (seed production only)[A,A]	Apply before alfalfa reaches 12 inches (30 cm) in height.
Chickling vetch grown for seed	Apply at the 5 - 7 leaf stage.

Δ Apply only to seedling alfalfa that will remain in production for at least 3 years following application. Apply only once during the life of the alfalfa stand.

ΔΔ Do not apply in the last year of established alfalfa stands.

Do **NOT** use in the brown or dark brown soil zones (except for use in dry beans and alfalfa under irrigated brown soils); rotational crops may be severely injured due to carryover in these soils.

Weeds and Staging:

In field peas and CLEARFIELD canola varieties. Apply up to the 4 leaf stage, unless otherwise indicated:

Chickweed
Cleavers
Green foxtail
Hemp-nettle*
Lamb's-quartersΔ
Redroot pigweed
Shepherd's purse*
Smartweed
Stinkweed
Volunteer canola (not CLEARFIELD varieties)
Volunteer barley**
Volunteer wheat**
Wild buckwheat
Wild mustard
Wild oats (2 to 4 leaf stage)

In seedling and established alfalfa:

Common groundsel***Δ
Green foxtailΔ
Green smartweed ***
Lamb's quarters***Δ
Redroot pigweed
Shepherd's purse***Δ
Stinkweed
Volunteer canola (not CLEARFIELD varieties)
Wild mustard

In dry beans:

Hairy nightshade (up to 6 leaf stage)

* In field peas only.
** Suppression in CLEARFIELD canola only.
*** Seedling alfalfa only.
Δ Suppression

Cost (2004 suggested retail price):

$22.64 per acre.

Rates:

85 mL/acre.

A non-ionic surfactant with at least 80% active ingredient (Agral 90, Agsurf, Surf 92) should be added at a rate of 0.25 L per 100L of spray solution. Do not over apply Pursuit, as crop injury may result. Only one application may be made during the growing season.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 to 160 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Plants are under stress when conditions before or after application are very hot or cold, excessively dry or wet. During periods of stress, plants are not actively growing and reduced weed control or crop injury may result.

Tank Mixes:

None registered.

Note: There are no tank mixes listed on the Pursuit label. To check for other possible tank mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze or harvest seedling alfalfa within 14 days of treatment. Do not graze or harvest field peas for feed within 30 days. Do not graze other treated crops or cut for feed prior to crop maturity.

Preharvest Interval: Do not apply within 60 days of harvesting field peas or chickling vetch, within 70 days of harvesting CLEARFIELD canola, within 75 days of harvesting dry beans, or within 85 days of harvesting soybeans.

Recropping: Rotate to barley, spring wheat (not durum), lentils, alfalfa, field peas or CLEARFIELD canola the year following application. The manufacturer recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crop other than those listed above. However, yield losses within the test strips may not be noticed unless the yield can be compared to an untreated area seeded adjacent to the Pursuit-treated strip. In case of crop failure, only field peas or CLEARFIELD canola may be replanted in the year of application.

Aerial application: Do not apply by air.

Storage: Do not freeze. If the product is exposed to temperatures below 0°C, thaw the product completely and shake the container vigorously prior to use.

Environment: Do not apply within 15 m of shelterbelts, water bodies, wetlands, and woodlots.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Keep out of reach of children.

Pursuit Ultra

Herbicide Group – 1, 2
(Refer to page 27)

Pursuit Ultra is a prepackaged tank mix of Pursuit (Page 173) at 0.04 L/acre and Poast Ultra (Page 162). Information listed is restricted to Crop, Weeds, Rates and Cost. For other detailed information on the component products including restrictions see the product pages listed above.

Company:

BASF Canada

Formulation:

The Pursuit Ultra package contains:

Poast Ultra (PCP#24835): 450 g/L sethoxydim formulated as an emulsifiable concentrate.

Pursuit (PCP#'23844): 240g/L imazethapyr formulated as a liquid.

Container size – 2 x (1.65L Pursuit + 7.7L Poast Ultra). Jugs are split internally and contain both Poast Ultra and Pursuit. Each split jug treats 40 acres.

Crops and Staging:

CLEARFIELD canola - after the 1 leaf stage.

Field pea - up to the 6th above ground node stage (six true leaves).

Do NOT use in the brown or dark brown soil zones as rotational crops may be severely injured due to carryover in these soils.

Weeds and Staging:

Grasses: at the 1 to 6 leaf stage:

Barnyard grass
Green and yellow foxtail
Quack grass*
Persian darnel
Proso millet
Volunteer barley
Volunteer oat
Volunteer corn
Volunteer wheat
Witchgrass
Wild oat

Broadleaf weeds up to the 4 leaf stage:

Chickweed
Cleavers
Hemp nettle (peas only)
Redroot pigweed**
Smartweed
Stinkweed
Volunteer canola (non-CLEARFIELD varieties)
Wild buckwheat**
Wild mustard

* Suppression
** Light infestations only

Rates:

Poast Ultra - 0.19 L per acre;

Pursuit - 40 mL per acre

(One split jug treats 40 acres).

Merge surfactant must be used with Pursuit Ultra.

Mix to a final ratio of 1.0 L of Merge in 100 L of spray solution. (1.0% Merge volume per spray solution volume).

Cost (2004 suggested retail price):

$26.34 per acre.

Application Information:

Water Volume: 40 L per acre

Pressure: 40 psi (275 kPa)

Nozzles: Stainless Steel 80° or 110° flat fan nozzles tilted forward at a 45° angle.

Refine Extra Toss-N-Go

Herbicide Group – 2
(Refer to page 27)

Company:

E. I. duPont Canada (PCP#22575)

Formulation:

50% thifensulfuron methyl plus 25% tribnuron methyl formulated as a dry flowable.

Container size - 4 water soluble bags (80 g each, total 320 g).

Crops and Staging:

Apply from 2 leaf to the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions

Cereals:

Barley
Oats
Wheat (including durum, spring and winter)

Seedling or established forage grasses for forage* or seed production:*

Bromegrass (meadow, smooth)
Creeping red fescue
Kentucky bluegrass**
Orchardgrass
Tall fescue
Wheatgrass (crested, intermediate, northern, pubescent, slender, streambank, tall, western)

Since the use of this product on forage grasses is registered under the User Requested Minor Use registration system, the manufacturer assumes no responsibility for herbicide performance.* ***Users of this product on forage grass do so at their own risk.*

** Established stands only.

Weeds and Staging:

Unless otherwise noted below, apply to young and actively growing weeds that are less than 4 inches (10 cm) in height or width.

Weeds Controlled:

Annual smartweed (green smartweed, lady's-thumb)
Ball mustard
Chickweed (1 to 6 leaf)
Common groundsel
Corn spurry
Cow cockle
Flixweed
Hemp-nettle
Kochia
Lamb's-quarters
Narrow-leaved hawk's-beard
Redroot pigweed
Russian thistle
Shepherd's-purse
Stinkweed
Tartary buckwheat
Volunteer canola (CLEARFIELD varieties with 2,4-D or MCPA mixes only)
Volunteer sunflowers
Wild buckwheat (1 to 3 leaf)
Wild mustard

Weeds Suppressed:

Canada thistle, sow-thistle (less than 6 inches (15 cm) tall or across and prior to budding)
Cleavers (1 to 3 whorls)
Round-leaved mallow (2 to 6 leaf)
Scentless chamomile
Stork's-bill (2 to 6 leaves)
Toadflax (less than 6 inches or 15 cm tall)

Cost (2004 suggested retail price):

$6.27 per acre.

Rate:

8 g/acre (one container treats 40 acres). Add Agral 90, Agsurf, or Citowett Plus at 0.2 L per 100 L of spray solution.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 22 L per acre.

Pressure: 30 to 40 psi (210 to 275 kPa).

Nozzles: Flat fan nozzles with 50 mesh line strainers and screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply to wheat, barley or oats that are stressed by severe weather conditions (frost, drought or water saturated soil) as crop injury may result.

Under certain conditions (heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) lightening in crop colour and reduction in crop height may occur.

Kochia control may be reduced during stress conditions or if extremely heavy infestations exist.

Tank Mixes:

Do not mix with substances that contain boron or that release chlorine.

Herbicides:

Tank mix Partner	CROPS			
	Spring wheat	Durum	Barley	Oats
2,4-D amine or ester (0.34 to 0.45 L/acre)*[§]	✓	✓	✓	
Assert (0.53 to 0.65 L/acre)	✓	✓	✓	
Assert (0.53 to 0.65 L/acre) + MCPA ester (0.28 to 0.45 L/acre)	✓	✓	✓	
Attain (80 acres per case)[§]	✓		✓	
Banvel II (44.5 mL/acre)[§]	✓		✓	
Curtail M (0.61 L/acre)[§]	✓		✓	
Everest (17.4 g/acre) + 2,4-D amine or ester (up to 0.45 L/acre)*[§]	✓			
Horizon (95 to 115 mL/acre) plus Score adjuvant	✓	✓		
Horizon (95 mL/acre) + Banvel II (44.5 mL/acre) plus Score adjuvant	✓			
Horizon (95 to 115 mL/acre) + MCPA ester (0.34 to 0.45 L/acre) plus Score adjuvant	✓	✓		
Lontrel 360 (85 mL/acre)[§]	✓		✓	
Lontrel 360 (85 mL/acre) + 2,4-D or MCPA ester (0.34 L/ acre)[§]	✓		✓	
MCPA amine or ester (0.28 to 0.45 L/acre)[§]	✓	✓	✓	✓

* 500 g/L formulation

[§] tank mix requires the addition of a non-ionic surfactant

Check the above tank mix partner(s) respective labels for additional staging and varietal restrictions.

Fertilizers: None registered. Do not mix soluble bags with liquid fertilizers.

Note: The above mixes are those listed on the Refine Extra label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Grazing: Must not be grazed or fed to livestock for 7 days after treatment.

Recropping: None the year after treatment. Canola, flax, lentils and alfalfa may be planted 2 months after an application of Refine Extra.

Aerial Application: Do not apply by air.

Storage: Store in a cool, dry place. May be frozen.

Tank Cleaning:

Refine Extra can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray Refine Extra should be flushed out immediately after Refine Extra is used. The manufacturer recommends that sprayers used to apply this product be flushed twice with a water/ammonia rinse (1L of 3% ammonia per 100 L of water). All nozzles, screens and filters should be removed and cleaned after applying this product. Rinse the tank, booms, hoses, and nozzles with clean water to complete the process.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 14 for additional information.

Hazard Rating:

Warning - Eye and Skin Irritant

Reflex*

*(For use only in the Red River Valley of Manitoba)

Herbicide Group – 14
(Refer to page 27)

Company:

Syngenta (PCP#24779)

Formulation:

240 g/L fomesafen formulated as a solution.

Container size- 10L.

Crops and Staging:

Dry beans[^] in the 1 to 2 trifoliate leaf stage. For use in the Red River Valley of Manitoba only.

[^] NOTE - Since applications to beans in the Red River Valley has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. ***Application to corn is at the risk of the user.***

Weeds and Staging:

Broadleaf weeds controlled by Basagran at the 0.7 L/acre rate plus improved control of the following weeds up to the 4-leaf stage:

Cocklebur
Eastern black nightshade
Lady's thumb
Lamb's-quarters
Ragweed (common)
Redroot pigweed
Velvetleaf*
Wild Mustard

* Suppression only

Cost (2004 suggested retail prices):

$7.79 per acre, not including cost of Basagran or adjuvant.

Rate:

Reflex is registered in the Red River Valley of Manitoba only as a tank mix at a rate of 235 mL/acre Reflex plus 0.71 L/acre Basagran plus Agral 90 at 1 L per 1000L of spray solution.

Application Information:

Water Volume: Minimum 81 L per acre. Increase water volume to 142 L per acre for fields with heavy weed densities or with weeds at the upper limit of their recommended stage.

Pressure: 275 kPa (40 psi). Increase pressure to 420 kPa (60 psi) for fields with heavy weed densities or with weeds at the upper limit of their recommended stage.

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Weed control and crop tolerance may be reduced under certain stress conditions such as cold temperatures, excess moisture, drought and injury from hail or previous herbicide applications.

Tank Mixes:

Herbicides: Reflex is only registered for use in a Basagran tank mix. See Rates.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Restrictions:

Rainfall: Reflex is unaffected by rain falling 4 hours after application.

Grazing: Do not graze treated crop or cut for hay. There is insufficient data to support such use.

Preharvest: Leave at least 84 days from application to harvest.

Recropping: Winter wheat may be sown 4 months after application. Spring wheat, dry beans and field corn may be grown the year following an application.

Do not apply Reflex to any field more often than once every two years.

These recropping restrictions refer only to the Red River Valley of Manitoba. Use outside this is region is not registered as recropping options have not been determined.

Aerial Application: Do not apply by air.

Storage: No information on label.

Environment: Leave a buffer zone of at least 15 m between the last spray swath and the edge of sensitive terrestrial areas such as shelterbelts, hedgerows and shrublands as well as aquatic areas such as ponds, streams, rivers, prairie potholes and sloughs. Do not apply when winds are greater than 15 km/hr.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

▽ Caution Poison

Warning- skin and eye irritant; potential skin sensitizer

Reglone Desiccant

Herbicide Group – 22

(Refer to page 27)

Company:

Syngenta (PCP#26396)

Formulation:

240 g/L diquat formulated as a solution.
Container size - 10 L.

Crops and Staging:

Reglone is used to dry immature green material at top of indeterminent crops and green weeds to facilitate harvest. Reglone will not speed maturity of green crops. Treatment before the recommended stage can result in reduced yield and quality. Add 0.1 L of Agral 90 or Agsurf per 100 L of spray solution for all applications except those indicated.

CROP	STAGE	RATE (L/ACRE)	
		Ground	Aerial
Dry beans, soybeans	Crop has lost 80 to 90 percent of leaves and 80 percent of pods are yellow.	0.5 to 0.7 (use high rate for dense crop, heavy weed infestations)	0.7 to 0.9 (use high rate for dense crop, heavy weed infestations)
Flax, Solin	75 percent of bolls brown.		
Lentils	Lowest pods are light brown and rattle when shaken.		
Mustard	75 percent of seed has turned brown		
Canola*	60 to 75 percent of seed has turned brown.		
Peas	Bottom pods are ripe and dry, seeds detached from pods.		
Chickpeas (No aerial application on chickpeas)	Plants yellow, pods mature, seeds changed colour and detached from pods.		
Sunflowers	Backs of sunflower heads and bracts are turning yellow and seed moisture is 20 to 50 percent.		
Potatoes (some top growth and/or some weeds)	Two weeks prior to harvest	0.7 to 0.9**	2 applications required. First application – 0.7 to 0.9 L/acre**. Second application (4 to 5 days later) at 0.5 L/acre
Potatoes (dense crop, heavy weed infestations)		1.4**	
Alfalfa, bird's -foot trefoil, red and white clover (for seed production only).***	Pods are ripe but before shattering. Harvest within 7 days.	0.7 to 1.1 (use high rates for dense crops, heavy weed infestations)	0.7 to 1.1 (use high rates for dense crops, heavy weed infestations)

* This product can cause shattering losses in canola. It should only be used on Argentine canola to assist in the harvest of a severely lodged crop.

** Use high rate for dense or immature vines.

*** Do not use on forage legumes that have been treated with a residual herbicide in the previous 12 months.

Cost (2004 suggested retail prices):

$22.11 per L plus surfactant.

Application Information:

Water Volumes:

Ground applications: 91 to 222 L per acre. Use 222 to 445 L per acre on potatoes.

Aerial applications: 18 L per acre. Use the highest water volumes when crop canopy is heavy or if weed growth is dense.

Pressure:

Ground application: 200 to 275 kPa (30 to 40 psi).

Aerial applications: 150 to 200 kPa (22 to 30 psi).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Best results under cloudy conditions or in evening.

Shattering losses can increase if heavy winds, rain or hail occur after the crop has dried down.

Tank Mixes:

Herbicides: None registered.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: Fungicides may be added when applying Reglone to potatoes for vine killing.

Restrictions:

Rainfall: Within 15 minutes may reduce effectiveness.

Grazing: Crop residues remaining after harvest may be fed to livestock.

Preharvest Interval: Wait 4 to 7 days before harvesting lentil. Do not exceed 7 days to harvest when treating forage legumes. Wait at least 7 days before harvesting Lupins. Wait 7 to 10 days before combining canola and mustard, but do not wait longer than 14 days. Wait 15 to 20 days for sunflowers. Harvest flax and peas when sample tests dry.

Recropping: No restrictions the year after application.

Aerial Application: May be applied by air in a minimum of 18 L per acre water volume. No aerial application on chickpeas.

Storage: Do not freeze.

Environment: Do not apply by air where wetlands or other good wildlife cover are present. Leave a 15 m border around the edges of sloughs and other wildlife cover. Apply when wind is between 3.5 and 9 km/hr.

Tank Cleaning:

When finished spraying Reglone, rinse the sprayer out with clean water. Run through pump, lines and nozzles. Drain tank by spraying out on an untreated portion of a crop on which the product is registered, or by spraying on uncropped land. Refill sprayer with water and Agral 90, Agsurf at 0.6 L per 1,000 L spray solution. Run the solution through lines and boom. Spray out, then refill with clean water. Leave equipment standing overnight, then drain water out.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members are likely to frequent or walk.

Refer to page 14 for additional information.

Hazard Rating:

◈ Warning Poison

Danger Eye Irritant

Warning Skin Irritant

Remedy EC

Herbicide Group – 4
(Refer to page 27)

Company:

Dow AgroSciences (PCP#26420)

Formulation:

480 g/L of triclopyr acid formulated as an emulsifiable concentrate. Container size - 2 X 10 L jugs

Crops and Staging:

Pastures and Rangeland. Staging is based on weed stage.

Weeds and Staging:

For best results, apply to the following weeds when leaves are fully expanded and plants are actively growing, until autumn coloration begins to appear. Single stem applications should not be used on woody species greater than 8 ft. (2.5 m) tall.

Woody plants controlled:

Alder
Ash
Aspen
Basswood
Beech
Birch
Blackberry
Buckthorn
Cherry*
Chokecherry*
Cottonwood
Dogwood
Elderberry
Elm*
Hawthorn
Maples
Oaks*
Poison oak
Pines*
Poplar
Red maple*
Raspberry*
Sumac
Tamarack
Wild rose
Willow

* These species may require treatment at the higher rate and may need to be retreated the following year, particularly if the original treatment was made at the lower rate.

Non-woody annual and perennial broadleaved weeds controlled are:

Burdock
Chicory
Curled dock
Dandelion
Field bindweed
Lamb's-quarters
Ragweed
Smartweed spp.
Smooth bedstraw
Vetch
Wild lettuce

Cost (2004 suggested retail price):

$28.00 per L

Rate:

Ground application:
Broadcast foliar - Woody species: 1.6 to 3.2 L/acre, Non-woody species: 0.4 to 1.62 L/acre
Single stem: mix 0.04 to 0.08 L per 10 L of spray solution.
Backpack (low volume foliar): 0.1 to 0.5 L per 10 L of spray solution.

Aerial application: 1.6 to 3.2 L/acre

Application Information:

Water Volume:
Broadcast foliar: Minimum 80 L per acre. Proper coverage is important for effective control. Even, uniform coverage, without gaps or pooling of droplets is preferred.
Single stem: Apply to the point of runoff.
Backpack (low volume foliar): Thoroughly wet foliage but not to runoff.
Larger Woody Species: For control of woody species larger than 8 ft. (2.5 m) the basal bark method of control can be used to apply small amounts of Remedy mixed in oil carrier (mineral oil or vegetable oil) directly to the bark of the trunk. Several methods can be used. See the label for directions on this use.

Aerial application: A minimum of 12 L per acre.

Pressure:
Broadcast foliar: Maximum 275 kPa (40 psi).
Nozzles: Use application equipment that provides even coverage of large droplets that are not prone to drift.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Avoid spraying if temperatures exceed 28°C.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfree period is specified on the label; required interval may be up to 8 hours. Contact the manufacturer for more information.

Grazing:

For lactating dairy animals:

Rates below 1.9 L/acre - treated areas may be grazed or harvested for greenfeed 14 days after treatment.

Label rates greater than 1.9 L/acre - restrict grazing or greenfeed for 60 days following treatment. Do not harvest treated area for hay for 60 days after treatment.

For other livestock:

Rates below 1.9 L/acre - There is no restriction on grazing or harvesting treated areas for greenfeed but do not harvest treated areas for hay for 7 days after treatment.

Label rates greater than 1.9 L/acre - restrict grazing and harvest for greenfeed or hay for 14 days following treatment.

If less than 25% of grazed area is treated, there is no grazing restriction for other livestock.

Remove all animals from grazing treated grass or consumption of treated hay at least 3 days prior to slaughter.

Aerial Application: May be applied by air.

Storage: Store above -2°C. If frozen agitate container before use.

Environment: This product is highly toxic to fish, aquatic crustaceans, and plants. Do not apply to water. Do not spray at the edge of wetlands, lakes, ponds, streams, rivers, and wetland wildlife habitats. Buffer zones are dependent on rate of application, method of application and boom height. Consult the product label for specific buffer zone requirements.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison.

Reward

Herbicide Group – 22
(Refer to page 27)

Company:

Syngenta (PCP#26271)

Formulation:

Reward - 240 g/L diquat formulated as a solution.

Container sizes - 4 x 3.78L.

Use:

Control of water weeds such as coontail, duckweed, Canada water weed, pond weeds, and water milfoil in farm dugouts. Offers temporary control of certain species of algae.

Cost (2004 suggested retail prices):

$46.17 per L

Timing:

Mid-May through late June.

Rates:

Dugouts less than 5 feet (1.5 m) deep: Apply Reward at 7.4 L per acre.

At this rate, 2.2 L of Reward will treat a dugout that is 160 feet by 80 feet (49 m x 24.4 m).

Dugouts more than 5 feet (1.5 m) deep: Apply Reward 10 to 11.8 L per acre.

At these rates, a dugout that is 160 feet by 80 feet (49 m x 24.4 m) will require 3.0 to 3.5 L of Reward.

Milfoil can be controlled in early stages by 3.7 L per acre.

Application:

Dilute 1 part Reward with 4 parts clean water.

Spray over the water surface.

How it Works:

Refer to Table 2 on page 29.

Restrictions:

Grazing: Do not use water for animal consumption for 24 hours after application.

Irrigation: Do not use water for irrigation for 5 days after application.

Domestic Use: Do not use water for human consumption for 5 days after application. Do not swim in water for 24 hours after treatment.

Storage: Do not freeze.

Environment: If weed growth is dense, protect fish by not treating more than one-fourth of dugout at a time.

Equipment Clean Out:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

Danger Eye Irritant
Warning Skin Irritant

Rustler

Herbicide Group – 4, 9

(Refer to page 27)

Company:

Monsanto Canada Inc. (PCP#27200)

Formulation:

194 g/L glyphosate present as the isopropylamine salt. 46 g/L dicamba present as a solution.

Container sizes - 10 L, 115 L.

Crops and Staging:

Summerfallow.

Preseeding in spring on fields to be sown to wheat, barley, oats and rye.

May also be applied prior to sowing field corn in fields with more than 2.5% organic matter (do not use on sandy or sandy loam soils).

Rustler **SHOULD NOT** be applied prior to broadleaf crops such as lentils, peas, canola and flax due to the risk of injury.

Weeds, Rates and Staging:

Application should be made to emerged, actively growing weeds. Application at early growth stages generally provides the best results.

Annual grasses - Apply 1 L per acre between emergence and heading. For wild oat apply at the 1 to 3 leaf stage.

Downy brome
Persian darnel
Green foxtail
Volunteer cereals
Wild oats

Annual broadleaves - Apply 1 L per acre up to 6 inches (15 cm) height unless otherwise indicated.

Cow cockle
Flixweed
Kochia
Lady's-thumb
Lamb's-quarters
Redroot pigweed
Russian thistle
Smartweed
Stinkweed
Volunteer canola*
Wild buckwheat
  (1 to 4 leaf)
Wild mustard

* NOT including Roundup Ready varieties.

Foxtail barley suppression- Apply 1.26 L per acre before initiation of the seed head.

Cost (2004 suggested retail price):

$6.24 to $7.86 per acre.

Application Information:

Water Volume: 20 to 40 L per acre water. Use of the lower water volume will improve control. Avoid the use of extremely hard water (greater than 700 ppm calcium or magnesium or high levels of iron).

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Reduced effectiveness may result if application is made to weeds that are drought-stressed, damaged by disease or insects. Poor control under cool, cloudy weather can occur. Dust on foliage can also cause reduction in control.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 6 hours may reduce weed control. Heavy rainfall within 2 hours of application may require a repeat treatment.

Grazing: Do not graze treated areas or cut for feed.

Recropping: No restrictions in the season following treatment. Do not apply in fall or spring prior to broadleaf crops such as lentils, peas, canola and flax due to the risk of injury.

Aerial Application: Do not apply by air.

Storage: Store above 5°C.

Equipment: Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.

Environment: Maintain a minimum buffer zone of 15 m between the downwind edge of the spray boom and non-target plants or wetland areas or other open water. Apply only in light wind conditions. Do not spray when conditions are dead calm or when wind is gusty.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger Corrosive to eyes

May cause irreversible eye damage.

Select

See clethodim on page 84.

Sencor

Herbicide Group – 5

(Refer to page 27)

Company:

Bayer CropScience

Sencor Solupak 75 DF - (PCP#20968)
Sencor 500F - (PCP#14867)
Sencor 480F - (PCP#26280)
Sencor 75 DF - (PCP#17242)

Formulations:

Sencor Solupak 75 DF - 75% metribuzin formulated as a dispersible granule. Container size - 2.5 kg (5 water soluble bags, 500 g each).

Sencor 500F -500 g/L metribuzin formulated as a flowable. Container size - 5 L.

Sencor 480F -480 g/L metribuzin formulated as a flowable. Container size - 5 L.

Sencor 75 DF - 75% metribuzin formulated as a dispersible granule. Container size - 2.5 and 5 kg.

Crops, Rates and Staging:

CROP	APPLICATION TIMING	RATES (kg or L/acre)
Barley	2 to 5 leaf stage.	Sencor 500F/480F - 0.111 to 0.223 Sencor 75DF - 0.08 to 0.152
Spring wheat (including durum)	2 to 5 leaf stage.	Sencor 500F/480F - 0.111 to 0.172 Sencor 75DF - 0.08 to 0.111
Winter wheat (Norstar only)	In late fall after tillers have developed (past the 3 leaf stage).	Sencor 500F/480F - 0.344 to 0.45
Peas (field and processing)*	Preplant incorporated (when tank mixed with Edge DC, Rival or Treflan EC)	**Spring:** Sencor 500F/480F - 0.223 to 0.283 Sencor 75DF - 0.152 to 0.192 **Fall:** Sencor 500F/480F - 0.280 to 0.343 Sencor 75DF - 0.190 to 0.223
Peas (field only)*	Postemergence - up to 6 inches (15 cm) of vine length. For short-statured, determinate flowering peas, apply at the early stages within this range.	Sencor 500F/480F - 0.172 to 0.223 Sencor 75DF - 0.111 to 0.152
	Split post-emergent applications**	**First application of:** Sencor 500F/480F - 0.08 to 0.11 Sencor 75DF - 0.06 to 0.08 Second application 7 to 10 days later with rates within these ranges.
Lentils*	Single or split applications** to plants up to 6 inches (15 cm) of vine length. For maximum crop tolerance, apply at the 1 to 4 above ground node stage.	Sencor 500F/480F - 0.172 Sencor 75DF - 0.111
		First split application of: Sencor 500F/480F - 0.08 to 0.11 Sencor 75DF - 0.06 to 0.08 Second split application 7 to 10 days later with rates within these ranges.
Chickpeas*	Up to 2.5 inches (6 cm) in height, when vines have 1 - 3 above ground nodes. Note application past recommended growth stage may result in severe crop injury.	Sencor 500F/480F - 0.167 Sencor 75 DF - 0.111
Potatoes (except Belleisle or Tobique)***	Preplant incorporated (with Eptam).	Sencor 500F/480F - 0.222 to 0.343 Sencor 75DF - 0.152 to 0.223
	Pre-emergence in sprinkler irrigation systems (apply only in a tank mix with Eptam 8-E)	Sencor 500F/480F - 0.222 to 0.343 Sencor 75DF - 0.152 to 0.223
Potatoes (except Atlantic, Belleisle, Eramosa, Tobique and red-skinned or early maturing varieties)***	Early postemergence (up to 4 inches or 10 cm in height).	Sencor 500F/480F - 0.222 Sencor 75DF - 0.151
Soybeans***	Preplant incorporated (tank mixed with Treflan EC).	Sencor 500F/480F - 0.172 to 0.343 Sencor 75DF - 0.111 to 0.223
Fababeans	Preplant incorporated (tank mixed with Treflan EC)	**Spring:** Sencor 500F/480F - 0.172 to 0.343 Sencor 75DF - 0.111 to 0.223 **Fall:** Sencor 500F/480F - 0.220 to 0.343 Sencor 75DF - 0.190 to 0.223

* DO NOT use on lentils, peas or chickpeas seeded less than 2 inches (5 cm) deep or in soils with less than 4 percent organic matter.

** Under certain field or weather conditions a split application may provide better weed control and crop tolerance than single applications. The first application should be made at the cotyledon to 2 leaf stage of the weeds. The second application should be made when a second flush of weeds have emerged or if weeds which were more advanced at the time of the first application have started to show regrowth. The split applications are normally 7 to 10 days apart.

*** Consult manufacturer or seed supplier for varietal tolerances to Sencor applications in soybean and potato.

Note: When Sencor is tank mixed with Edge or Treflan in peas, fababeans, and soybeans, refer to product label for maximum rates that can be applied on light textured soils.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Postemergence applications should be made when weeds are small – 2 inches (5 cm) in height or diameter.

Split applications (postemergence on lentils and peas) – 1st application at cotyledon to 2 leaf stage of weeds. 2nd application (if necessary) 7 to 10 days after the first.

Post-emergence Sencor 75 DF at 81 g/acre or Sencor 500F 121 mL/acre:

Weeds controlled in spring wheat, barley, field pea and suppressed in lentil and chickpea:

Chickweed
Green smartweed
Hemp-nettle*
Lamb's quarters
Stinkweed
Volunteer canola
Wild mustard

Additional weeds controlled in spring wheat and barley only:

lady's-thumb
redroot pigweed

Postemergence with Sencor 75 DF at 111 g/acre or Sencor 500F at 172 mL/acre:

Weeds controlled in spring wheat, barley, potato, field pea, and suppressed in lentil and chickpea:

Weeds above plus:
Ball mustard
Corn spurry
Hemp-nettle
Tartary buckwheat

Additional weeds controlled in spring wheat and barley only:

Common Groundsel
Night-flowering catchfly
Wormseed mustard

Post Emergence with Sencor 75 DF at 152 g/acre or Sencor 500F at 222 mL/acre in spring wheat and barley only:

Weeds above plus:
Henbit
Russian thistle

Post Emergence with Sencor 75 DF at 152 g/acre or Sencor 500F at 222 mL/acre in potatoes only:

Weeds listed for peas above plus:
Lady's-thumb
Redroot pigweed
Shepherd's-purse

Post Emergence in winter wheat with Sencor 500F at 344 to 450 mL/acre in controls:

Downy brome
Flixweed
Shepherd's-purse
Stinkweed

Preplant Incorporated in fababeans, lentils, field pea, soybean:

Must be applied in tank mix with Treflan EC or Rival DF):
Chickweed
Corn spurry
Green smartweed
Hemp-nettle
Lamb's quarters
Stinkweed
Volunteer canola
Wild mustard
Plus weeds controlled by Treflan EC.

* Use the high rate for best control.

** Suppressed only in lentils and chickpeas.

Cost (2004 suggested retail price):

Sencor 75 DF & Sencor Solupak 75 DF - $72.40 per kg

Sencor 500F/Sencor 480F - $43.06 per L

Application Information:

Water Volume:
Preplant incorporated: 40 L per acre.
Postemergence applications:
Cereals - 40 L per acre.
Lentils, peas, chickpeas - 70 L per acre.

Pressure: 30 to 40 psi (200 - 275 kPa).

Nozzles: Flat fan nozzles with opening no smaller than 8002 or TK2 with 50 mesh screens. For lentils, peas and chickpeas use nozzles no smaller than 8003 or TK3. Angle nozzles 45° forward.

Incorporation: All plant growth and stubble should be thoroughly worked into the soil before treatment. Apply directly to the soil surface. Two incorporations are required at right angles for thorough mixing. The first incorporation must be made within 24 hours of spraying. For fall applications, it is preferred that both incorporations be done in the fall. The second incorporation may be delayed until spring to conserve trash; however, both incorporations must be done the recommended depth.

Incorporate with a tandem disc, discer or field cultivator (Vibrashank type). Set equipment to work at a depth of 3 to 4 inches (8 to 10 cm). Operate disc implements at 4 to 6 mph (7 to 10 km/hr), cultivators at 6 to 8 mph (10 to 13 km/hr).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions

Crop height reductions or yellowing may occur if high temperatures occur within 48 hours of application. Cold, cloudy weather or frost within 3 days of application will also aggravate injury. If frost occurs, allow 4 to 5 days for crop to recover prior to applying Sencor. Heavy rainfall soon after application to peas, lentils and chickpeas can result in stand reduction on soils with less than 4 percent organic matter.

Tank Mixes:

Herbicides:

In spring wheat or barley:
Dicamba, Target, MCPA amine or 2,4-D amine.

In potatoes (preplant incorporated):
Eptam 8-E (Required).

In fababeans (preplant incorporated):
Treflan EC (Required).

In soybeans (preplant incorporated):
Treflan EC (Required).

In peas:
Treflan (PPI)
Rival (PPI).
Sencor 500F/480F at 0.113 L/acre or Sencor 75 DF at 0.077 kg/acre plus 0.19 L/acre MCPA sodium salt (300 g/L).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Sencor label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Allow 5 days between application of Sencor and application of other pesticides.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 6 hours may reduce control.

Grazing: Do not graze treated cereal crops within 30 days of application, or peas and lentils within 70 days of application.

Preharvest Interval: Do not harvest barley, wheat or potatoes within 60 days of application. Do not harvest lentils, chickpeas, or field peas within 70 days of application. Do not harvest processing peas within 40 days of application.

Recropping: Preplant incorporated treatments may leave a residue in the soil that will affect succeeding crops when using higher rates of product. Do not seed canola, sunflowers, onions, celery, peppers, cole crops, lettuce, spinach, red beets, turnips, pumpkin, squash, cucumbers or melons the year after treatment. Fall seeded crops may be injured when seeded the same year as preplant or postemergence applications of these products.

Aerial Application: No restrictions on label. Reduced weed control and increased crop injury can occur with aerial applications.

Storage: May be frozen.

Environment: Avoid drift onto sensitive plants and contamination of water or wetland areas.

Tank Cleaning:

Spray equipment must be thoroughly cleaned to remove traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution from the spray tank. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of 0.25 L per 100 L of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of the spray tank, nozzles or spray rig.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 14 for additional information.

Hazard Rating:

Keep out of reach of children.

Shotgun

Herbicide Group – 4,5
(Refer to page 27)

Company:

United Agri Products (PCP#24608)

Formulation:

280g/L atrazine and 120 g/L 2,4-D ester formulated as suspension concentrate.

Container size - 2 x 10 L.

Crops and Staging:

Field Corn up to the 4 leaf stage.

NOTE: The 2,4-D component may cause corn to become brittle and prone to breaking from strong wind or cultivation for about two weeks following application.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 1 to 6 leaf stage, unless specified:

Lamb's-quarters
Ragweed (Common)
Redroot pigweed
Smartweed (annual)
Wild buckwheat (1 - 4 leaf)
Wild mustard

Cost (2004 suggested retail price):

$10.59 to 15.57 per acre.

Rate:

Broadcast applications:
Apply at 0.97 L per acre on light textured soils (sandy) and up to 1.46 L per acre on medium to heavy textured soils (silt to clay). One 10 L jug treats 6.8 to 10 acres.

Band applications:
Shotgun may also be applied in a band over the crop row if another method will be used to control weeds between rows. For band treatment, use proportionately less; for example, for 10 inch band on 30 inch row, use 1/3 of the broadcast rate over the total area covered by the sprayer.

Make only one application per year.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 to 81 L per acre.

Pressure: 35 to 40 psi (240 to 275 kPa)

Nozzles: Use flat fan nozzles that deliver a uniform coverage while minimizing fine droplets prone to drift. Banding applications require special even flat fan nozzles, designed to deliver an even spray pattern across the band. Use 50 mesh or larger screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

This product should be applied when the temperature is about 10° to 28°C for optimum results. Cooler temperatures delay or reduce its effectiveness and higher temperatures increase the risk of vapour movement to susceptible crops. Do not apply when there is a risk of severe fall in night temperature. Weed control and crop tolerance will be best when plants are not subjected to excessive moisture or drought stress before and after application.

If conditions are very dry, do not spray until one week after rain.

Tank Mixes:

Herbicides:
Accent (13.5 g/acre) plus recommended adjuvant
Ultim (13.5 g/acre) plus recommended adjuvant – Manitoba only.

Fertilizer: None registered.

Insecticides: None registered.

Fungicides: None registered.

Do not add any other adjuvants, surfactants, oils, fertilizers or other pesticides to the spray mixture.

Note: The above mixes are those listed on the Shotgun label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Rainfall within 6 hours may reduce control. A 24 hour rainfree period is preferred.

Grazing: Do not harvest forage or graze fields until ear emergence of the corn.

Preharvest Interval: No information.

Recropping: Under good moisture conditions most rotational crops may be planted the season following application. When extended periods of dry weather occur during the year of treatment, injury may occur to dry beans, peas or other legumes.

Aerial Application: Do not apply by air.

Storage: Do not store below -10°C. If frozen, warm to 5°C or more and reconstitute by shaking or rolling container.

Environment: Avoid drift onto non-target areas. To avoid contamination, leave a buffer of 10 meters from the sprayed area to water bodies, wetland areas and plants that may be injured by Shotgun. Do not mix within 30 m of these areas. Do not spray near sensitive crops when winds are gusty or in excess of 8 km/h and moving towards sensitive crops.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison.

Danger Eye Irritant.

Caution Skin Irritant.

Keep out of reach of children.

Simazine

Herbicide Group – 5

(Refer to page 27)

Company:

Syngenta (Princep Nine-T - PCP#16370)
United Agri Products (Simazine 480 - PCP#23181)

Formulations:

Princep Nine-T: 90% simazine formulated as a water dispersible granular. Container size - 5 kg.

Simazine 480: 500 g/L simazine formulated as a flowable liquid. Container size – 2 x10L.

Crops and Staging:

Established alfalfa or bird's-foot trefoil (Princep Nine-T only).

Do not use in year of seeding. Apply after final cut in fall until freeze-up. Do NOT apply to the same field more than three consecutive years. Residues may build up with yearly applications.

Established shelterbelts (elm, caragana, green ash, Manitoba maple). - fall or early spring before weeds begin growth.

Do NOT apply to frozen ground.

Weeds and Staging:

Simazine is applied prior to the emergence of the weeds and kills them when they are exposed to the treated layer of soil.

Barnyard grass
Lamb's-quarters
Lady's-thumb
Perennial species
  starting from seed
Purslane
Ragweed
Smartweed
Volunteer clovers
Wild Buckwheat
Wild oats
Yellow foxtail

Cost (2004 suggested retail prices):

Princep Nine-T – $16.60 per kg

Simazine liquids - $11.13 per L

Rates:

Forage crops:
Princep Nine-T - 0.45 kg/acre.

Shelterbelts:
Princep Nine-T - 2 to 3 kg/acre,
Simazine 480 – 3.64 to 5.46 L/acre

Application Information:

Water Volume: Minimum 121 L per acre. In shelterbelts, use a minimum of 202 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan nozzles with 50 mesh or coarser nozzle screens.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

When applying to forage stands, dry soil conditions at the time of weed emergence may result in reduced weed control.

Tank Mixes:

None registered.

Note: The above mixes are those listed on the simazine labels only. For additional mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Moderate rainfall after application enhances activity.

Grazing: In forage stands, allow 30 days between application and grazing, 60 days between application and cutting for feed.

Recropping: Simazine is persistent and residues may persist for several years depending on soil pH, available soil moisture, number of yearly applications, and the sensitivity of the following crop. Corn will tolerate soil residues of simazine and may be planted the year of application. White beans, onions, peas may be injured 12 month after application.

Aerial Application: Do not apply by air.

Storage: May be frozen. Store in a cool, dry place.

Environment: Do not use within 10 m of wells, wetlands, or other open water. Do not mix within 30 m of these areas.

Tank Cleaning:

Refer to page 14.

Spectrum

Herbicide Group – 2,4

(Refer to page 27)

Company:

Dow AgroSciences

Formulation:

Each case of Spectrum contains 2 components:

Spectrum A (PCP#27032): 50 g/L florasulam formulated as a suspension concentrate.

Spectrum B (PCP#27032): 50 g/L clopyralid and 280 g/L of MCPA ester formulated as an emulsifiable concentrate.

Container size- Spectrum A - 0.8 L, Spectrum B - 12 L.

Crops and Staging:

Spring wheat (including durum), barley and oats in the 2 to 6 leaf stage.

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Canada thistle
Chickweed
Cleavers
Dandelion**
Flixweed*
Hemp-nettle
Lamb's-quarters
Redroot pigweed
Shepherd's-purse
Smartweed
Sow-thistle (annual)
Sow-thistle (perennial)****
Stinkweed
Stork's-bill
Volunteer canola
(all varieties)
Wild mustard
Wild buckwheat

Broadleaf weeds suppressed:

Dandelion***

* spring seedlings only
** seedlings and overwintered rosettes < 15 cm
*** overwintered rosettes > 15 cm; mature plants
**** top growth control only

Cost (2004 suggested retail price):

$14.00 per acre.

Rate:

Spectrum A: 40 mL/acre

Spectrum B: 600 mL/acre

One case treats 20 acres.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13 of the Guide to Crop Protection.

Application Information:

Water Volume: 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of Spectrum. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application, may increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Assert (0.65 L/acre) plus Acidulate

Puma^{120} Super at 0.155 L/acre* or 0.31 L/acre.**

In spring wheat (including durum) only:

Everest (17.4 g/acre) plus non-ionic surfactant

Horizon at 95 mL/acre or 115 mL/acre plus Score Adjuvant.**

* For control of green foxtail only

**This tank-mix may cause a reduction in wild oat control.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Spectrum label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze treated crop or cut for feed within 7 days of application.

Pre-harvest: Do not apply within 60 days of harvest.

Recropping: Barley, canola, field peas*, oats and wheat may be grown the year following an application or the field can be summerfallowed.

** Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field peas following drought conditions in the previous year.*

Aerial Application: Do not apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Environment: Spray when winds are under 15 km/hr, but not dead calm. Leave at least 30 m between the downwind edge of the spray boom and the closest edge of non-target land areas such as shelterbelts, hedgerows and shrublands. Leave at least 5 m between the downwind edge of the boom and the closest edge of water and wetland areas such as ponds, streams, rivers, prairie potholes and sloughs.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Caution Poison

May cause eye irritation

Sundance

Herbicide Group – 2
(Refer to page 27)

Company:

Monsanto Canada Inc.

Formulation:

Sundance (PCP#25826): 75% sulfosulfuron formulated as a water dispersible granule.
Container size – 432g per case.

Merge Adjuvant (PCP#24702): – 8 L.

pH adjuster: 7% ammonium hydroxide solution.
Container size: 4L jug.

Crop and Staging:

Spring wheat (including Avonlea, Kyle, Morse and Plenty durum) prior to emergence of the 4th tiller. (Do not underseed forage legumes).

Weeds and Staging:

Wild oat: From emergence to the 6 leaf stage with no more than 3 tillers. For optimum weed control and crop yield, apply prior to tillering.

Green foxtail*: emerged but prior to heading.

Annual Broadleaf weeds: cleavers, common chickweed, redroot pigweed, stinkweed, volunteer canola (NOT including CLEARFIELD canola), wild mustard - Apply to the seedling stage for optimum weed control.

Perennial Grass and Broadleaf weeds: dandelion**, foxtail barley, perennial sow-thistle**, quack grass* - Prior to heading/flowering when plants are actively growing

* Suppression only
** Spring seedlings are controlled and established plants are suppressed.

Cost (2004 suggested retail price):

$15.50 per acre including adjuvant.

Rates:

pH adjuster: 1 L per 400 L of spray solution.

Sundance: 11g per acre (40 acres per case)

Merge: 0.2 L per acre.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 20 to 40 L per acre.

Pressure: 40 psi (275 kPa)

Nozzles: Flat fan. Choose nozzles that provide optimum spray distribution and coverage at the appropriate spray pressure. Do not use flood jet or controlled droplet application equipment.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

For optimum results, do not apply to weeds growing under stress.

Tank Mixes:

Herbicides:

2,4-D ester (500 g/L formulation) at 0.340 L/acre for additional control of lamb's-quarters, wild buckwheat, volunteer CLEARFIELD canola, stork's-bill, dandelion**, narrow-leaf hawk's-beard**, kochia, annual smartweed spp., and perennial sow-thistle**.

Pardner at 0.4 L / acre for additional control of weeds controlled by Pardner plus narrow-leaf hawk's-beard**.

MCPA ester (500 g/L formulation) at 0.385 L/acre for additional control of weeds listed on MCPA ester label plus narrow-leaf hawk's-beard***, wild radish, volunteer sunflower, common plantain, and wild buckwheat*.

Buctril M at 0.405 L/acre for additional control of those weeds listed on the Buctril M label plus dandelion**, hemp-nettle, velvetleaf, Canada thistle* and perennial sow-thistle*.

* Suppression only
** Control of spring seedlings and season-long control of established plants.
*** Spring Seedlings only

Fertilizers: None registered.

Insecticides: None registered.

Note: The above mixes are those listed on the Sundance label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Do not apply if rain is forecast during, or soon after, application.

Grazing: Do not graze in the year of treatment.

Pre-harvest interval: No information available on label. Contact manufacturer for information.

Recropping:

Soils with GREATER than 4% organic matter:

Normal Moisture conditions	Wheat (including durum), canola (all varieties), barley, peas and flax may be grown the season after application.
Under Drought conditions*	Seed only wheat or Clearfield canola the season following application. The crops listed above can be seeded the second season following application.

Soils with LESS than 4% organic matter:

Normal Moisture conditions	Wheat (including durum), and Clearfield canola may be seeded one season after application. In the second season after application, barley, canola (all varieties), peas and flax may be grown in addition to the crops above.
Under Drought conditions*	Seed only wheat (including durum) or Clearfield canola for two seasons following application. Barley, canola (all varieties), peas and flax as well as the crops listed above can be seeded the third season following application.

* Drought conditions (less than 80% of normal June to Sept. rainfall) for high pH soils (greater than 7.5) and severe drought (less than 65% of normal June to Sept. rainfall) for all soils.

Re-seed only to wheat in the year of application. For all other crops, allow the most severe restriction for the soil and moisture conditions and conduct a field bioassay prior to planting.

Aerial Application: Do not apply by air.

Storage: Store under cool, dry conditions (below 50° C) away from foodstuffs, feed or seed.

Environment: Do not apply to water or contaminate water when draining sprayer. Maintain a buffer zone of 30m between the last spray swath and the edge of sensitive areas (shelterbelts, wood lots, or gardens). Leave a buffer zone of 6m between the last spray pass and wetlands or ponds. Use precautions to AVOID DRIFT. Drift may cause soil residues that will injure subsequent crops.

Tank Cleaning:

Do not drain or clean sprayer in an area where wastewater may drain into wells, other water bodies, or surrounding desirable vegetation. Flush tank, boom, and hoses with clean water for a minimum of ten minutes to remove all visible residues of the product. Fill the tank with water and add 1 L of household ammonia (minimum 3% ammonia content) per 100 L of rinse water and agitate. Flush solution through booms and hoses then top up tank with clean water. Agitate for 15 minutes, flush the hoses and booms again and then drain the tank. Remove nozzles and screens and clean separately with the ammonia solution as above. Repeat the ammonia rinse procedure as above. Thoroughly rinse the tank with clean water for at least 5 minutes, flush water through hoses and booms, and drain. Dispose of wash solution by spraying in a waste area or on the treated field.

For additional information, refer to page 14.

Hazard Rating:

Sundance: Slight skin irritant

Merge adjuvant: Warning Poison; skin irritant

pH adjuster: Caution corrosive

Target/Sword

See dicamba+mecoprop+MCPA section on page 90

Thumper

Herbicide Group – 4, 6

(Refer to page 27)

Company:

Bayer CropScience (PCP#22659)

Formulation:

280 g/L bromoxynil and 280 g/L 2,4 D ester formulated as an emulsifiable concentrate.

Container size - 8 L.

Crops and Staging:

Spring wheat (including durum) and barley at the 4 leaf to early flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 1 to 4 leaf stage:

American nightshade
Ball mustard
Bluebur
Cocklebur
Common ragweed
Cow cockle
Flixweed
Green smartweed
Lady's-thumb
Night-flowering catchfly
Pale smartweed
Redroot pigweed
Shepherd's-purse
Volunteer canola
Volunteer sunflower

Weeds controlled at the 1 to 8 leaf stage:

Common groundsel
Lamb's-quarters
Stinkweed
Tame buckwheat
Tartary buckwheat
Wild buckwheat
Wild mustard

Weeds Controlled from 1 to 12 leaf (max. 2 in. tall):

Kochia
Russian thistle

Cost (2004 suggested retail price):

$6.60 per acre.

Rates:

0.4 L/acre (one 8 L container treats 20 acres).

Application Information:

Water Volume: 20 to 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° flat fan nozzles. All strainer and nozzle screens must be 50 mesh or coarser.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Less than acceptable weed control may be expected if weeds are under stress because of excessive moisture, drought, or cool weather.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Achieve Liquid (0.2 L/acre) plus Turbocharge Aduvant
Avenge (1.4 to 1.7 L per acre)
Puma^{120} Super (0.15 to 0.31 L/acre)

In spring wheat (including durum) only:

Horizon (95 mL per acre) plus Score adjuvant

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Thumper label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze or cut for livestock feed within 30 days of application. Withdraw meat animals 3 days before slaughter.

Preharvest Interval: Do not harvest within 30 days of application.

Recropping: No recropping restrictions the year after application.

Aerial Application: May be applied by air in 12 to 16 L/acre of water. Use the higher volume when there is a heavy crop canopy, or when the majority of weeds are cow cockle, smartweed, or pigweed.

Storage: May be frozen.

Environment: Apply when wind is blowing lightly away from non-target areas and water bodies or wetland areas.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

◈ Warning Poison

Eye Irritant

Tordon 22K

Herbicide Group – 4

(Refer to page 27)

Company:

Dow AgroSciences (PCP#9005)

Formulation:

240 g/L picloram acid present as a potassium salt, formulated as a solution.

Container size - 2 X 10 L jugs

Crop and Staging:

Apply at any stage of permanent grass pastures, rangeland and non-cropland.

Weeds, Rates and Staging:

For the control of biennial and deep-rooted perennial weeds listed below:

Weed	Rate L/Acre	Backpack (mL of Tordon 22K per 100 square metres)*
Scentless chamomile	0.445	11
Knapweed (diffuse, spotted)	0.910	22
Canada thistle, pasture sage, poverty weed, Russian knapweed, perennial sow-thistle	1.8	45
Leafy spurge, field bindweed, toadflax	3.6	90

* mix with 18 litres of water and the spray solution over 100 square metres.

For best results, applications should be made when perennial weeds have fully developed, green leaves. Application in late summer (or periods of dry weather) when plants are not actively growing may result in unsatisfactory control.

Cost (2004 suggested retail price):

$41.00 per L.

Only available through Dow AgroSciences trained agents. Contact Dow AgroSciences for further information.

Application Information:

Water Volume: 160 to 325 L per acre.

Pressure: 150 to 350 kPa (20 to 50 psi).

Nozzles: Flat fan recommended. Non-target broadleaf plants are very sensitive to picloram drift.

Avoid conditions that are conducive to drift. (See page 12 for drift control suggestions)

How it Works:

Picloram interferes with cell division, causing leaf cupping, stem distortion and eventual death. Picloram is absorbed through the leaves and roots.

IMPORTANT: Picloram is a very persistent and water-soluble herbicide. Treated soil should not be moved from the treated area. Do not apply to soils that are permeable, have sinkholes, or lie over limestone bedrock. Do not apply to soils whose surfaces are composed of fractured rock or unconsolidated gravel. Application to these sites may allow the movement of herbicide to underlying water sources or aquifers. When applying Tordon 22K over sandy soils ensure that aquifers are not within 1.8 m of the soil surface. If shallow aquifers are present, do not apply Tordon 22K.

Tordon 22K must not be applied on range and pasture acres that are irrigated. Do no compost or mulch clippings from grass treated with Tordon 22K.

Effects of Growing Conditions:

Avoid spraying if temperatures exceed 28°C.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Rain within 6 hours of application may cause poor results. Heavy rainfall may dissolve and carry picloram away from the target area, or it may leach dissolved picloram out of the root zone or to undesirable locations.

Grazing: Do not graze lactating dairy animals within 6 weeks after treatment. There are no grazing restrictions for other livestock. Do not use manure from animals grazing treated forage to fertilize susceptible plants or crops.

Recropping: Tordon 22K may persist in the soil for up to 5 years. For this reason Tordon 22K may only be applied on permanent grass pastures and rangeland unless applied by an authorized pesticide applicator. Avoid the root zone of desirable trees or shrubs.

Aerial Application: Do not apply by air.

Storage: Do not freeze.

Environment: This product is moderately toxic to fish. Apply this product well back from any water bodies to avoid contamination. Heavy rains can move this product from its application site down slope toward sensitive areas. Do not load or mix near wells, dugouts or other water bodies.

Tank Cleaning:

Tordon 22K can cause severe injury to sensitive crops (especially pulses and other broadleaf crops) at very low concentrations. Spray equipment should be flushed out immediately after spraying Tordon 22K. The manufacturer recommends that equipment should be flushed twice with a water/household ammonia rinse (1 L of 3% ammonia per 100 L of water). All nozzles screens and filters should be removed and cleaned after use. Do not clean equipment upslope of water bodies or ditches near crop-land or shelterbelts.

Refer to page 14 for additional information.

Hazard Rating:

☠ Caution Poison.

Trifluralin

(Summerfallow use in the brown soil zone of Saskatchewan)

Herbicide Group – 3
(Refer to page 27)

Company:

Dow AgroSciences (Advance 10G, Treflan QR5)
NuFarm (Rival 10G)
United Agri Products (Bonanza 10G)

Formulation:

Advance 10G (PCP#21742): 10% trifluralin formulated as a granular in 22.7 kg and 454 kg bags.

Bonanza 10G (PCP# 22744): 10% trifluralin formulated as a granular in 22.7 kg, 500 kg bags.

Rival 10G (PCP#18926): 10% trifluralin formulated as a granular in 22.7 kg, 567 kg bags.

Treflan QR5 (PCP#14545): 5% trifluralin formulated as a granular in 25 kg, 725 kg bags.

Crops and Staging:

Spring wheat (including durum) in the brown soils zone of Saskatchewan. **Not for use in Manitoba.**

Apply to summerfallow in May, June, or July for weed control during both years of a fallow-wheat rotation or in the fall (September or October) or spring prior to wheat seeding. Do not apply to stubble when the previous crop was treated with another trifluralin product (Treflan, Advance, Rival or Bonanza). This includes application the previous summer or fall. Do not apply trifluralin to stubble or fallow when the previous year's crop was an oilseed, barley or pulse crop treated with a deep incorporated, spring or fall applied trifluralin product.

Weeds and Staging:

The followed weeds will be controlled prior to emergence:

Fallow Year:

Green foxtail	Wild buckwheat
Wild oat	Cow cockle
Persian darnel	Lamb's-quarters
Barnyard grass	Redroot pigweed
Russian thistle*	

Crop Year:

Green foxtail	Lamb's-quarters
Wild oat *	Wild buckwheat *

* Suppression only

Cost (2004 suggested retail price):

Advance 10G: $3.46 per kg.
Bonanza 10G: $3.21 per kg.
Rival 10G: $3.13 per kg.
Treflan QR5: $1.88 per kg.

Rates:

	RATE (KG/ACRE)			
	5G Formulations		10G Formulations	
Soil Organic Matter (%)	**1 to 3**	**4 to 8**	**1 to 3**	**4 to 8**
May	7.7	9	3.85	4.5
June	6.5	7.7	3.25	3.85
July	5.3	6.5	2.65	3.25
	2-8% Organic Matter			
September to October	4.5*		2.75*	
Spring (April or May, in the year of seeding)	4.5*		2.75*	

* Control of green foxtail only.

During the fallow year, susceptible weeds may not be fully controlled until after the second fallow operation has established a uniform layer of treated soil. Control of wild oats in the crop year may be variable depending on wild oat population as well as soil and climatic conditions. Some wild buckwheat may escape but its growth will be retarded and result in limited competition to the wheat crop.

Application:

Do not apply to sandy soils with less than 1% organic matter. Application to severely eroded knolls is not recommended. Do not apply to wet soils, soils in poor working condition, soils which contain more than 8 percent organic matter, or soils subject to prolonged periods of flooding.

Granules may be applied to standing or pre-worked stubble, provided trash or green growth does not interfere with cultivation (prevent soil mixing).

Over-application caused by overlapping, improper calibration or non-uniform application may result in reduced crop stand, delayed development or reduced yields.

Close applicator lid after filling to avoid prolonged exposure to direct sunlight.

Incorporation:

May - July: Apply granules to the soil surface and incorporate immediately, in the same operation if possible. Do not delay incorporation more than 24 hours after application. Use a deep tillage cultivator, field cultivator or disc implement set to work 2 to 3 inches (5 to 8 cm) deep, and operating at 8 to 10 km/hr. A second incorporation at the same depth and at an angle to the first should be done when weed growth requires it. Wait at least one week before making the second incorporation. After completing two fallow incorporations, additional operations with a rod weeder, shallow tillage or fall 2,4-D application may be required to control remaining weed growth.

September - October: Apply granules after harvest until freeze-up and incorporate immediately, in the same operation if possible with a deep tillage cultivator, field cultivator or disc implement set to work 2 to 3 inches (5 to 8 cm) deep. Do not delay incorporation more than 24 hours after application. A second incorporation may be done in the fall a minimum of 3 days later. Alternatively, in order to conserve trash cover throughout the winter, the second incorporation can be completed in the spring at the same depth and at an angle to the first incorporation. When both incorporations take place in the fall, shallow spring tillage should be completed in the spring. If a discer or air seeder is used for seeding, separate spring tillage may not be necessary.

Granules should not be incorporated when soil is crusted, lumpy or too wet for good mixing action. Fall application is not recommended on soils where a lack of trash cover combined with the required incorporation would leave the soil vulnerable to erosion.

Spring (April - May): Apply granules and incorporate immediately, in the same operation if possible. Do not delay the first incorporation more than 24 hours after application. The second incorporation must be delayed a minimum of 3 days following the first incorporation.

When applied to cold soils, wait 14 days before making second incorporation. The second incorporation should be done at an angle to the first incorporation, and at the same depth. If a discer or air seeder is used for seeding, the seeding operation can be used as the second incorporation.

Seeding:

Allow soil to warm before seeding to reduce risk of injury to crop. Place seed 1.25 to 2.5 inches (3 to 6 cm) deep. If spring seedbed preparation is required, set cultivator 2 inches (5 cm) deep. To reduce the risk of wheat injury, use good quality seed and agronomic practices that will promote good growing conditions. Avoid deep seeding, loose seedbeds and seeding into cold soils. If extended dry periods were present after a fallow application, a 10 percent increase in seeding rate is recommended.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Prolonged drought conditions after a May-July application to fallow may result in higher levels of trifluralin in the soil at the time of seeding. If conditions are not conducive to rapid emergence and growth of the wheat crop (for example cold or dry soil conditions) injury can occur.

Restrictions:

Rainfall: Once the product is incorporated into the soil, rainfall has no effect.

Grazing: Do not graze or cut for hay; insufficient data to support such use.

Recropping: Oats, canaryseed and small seeded forage grasses such as timothy and creeping red fescue should not be grown in rotation following a trifluralin treated crop.

Aerial application: Do not apply by air.

Storage: Store in a cool, dry location, out of direct sunlight.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Keep out of reach of children.

Trifluralin (Green and Yellow Foxtail Control in Cereals)

Herbicide Group – 3
(Refer to page 27)

Company:

Dow AgroSciences (Advance, Treflan)
NuFarm (Rival)
United Agri Products (Bonanza)

Formulations:

Advance 10G (PCP#21742): 10% trifluralin formulated as a granular in 22.7 kg and 454 kg bags.

Bonanza 10G (PCP#22744): 10% trifluralin formulated as a granular in 22.7 kg and 500 kg bags.

Bonanza 400 (PCP#21967): 400 g/L trifluralin formulated as an emulsifiable concentrate in 10 L, 115 L, 205 L containers.

Rival 10G (PCP#18926): 10% trifluralin formulated as a granular in 22.7 kg, 567 kg bags.

Rival EC (PCP#18612): 500 g/L trifluralin formulated as an emulsifiable concentrate in 9 L, 110 L, 200 L, 900 L containers.

Treflan EC (PCP#23933): 480 g/L trifluralin formulated as an emulsifiable concentrate in 9.45 L, 115 L containers.

Treflan QR5 (PCP#14545): 5% trifluralin formulated as a granular in 25 kg, 725 kg bags.

Crops and Staging:

Liquids applied in spring only (after seeding but prior to crop emergence) - spring wheat (including durum), barley.

Granulars applied in fall only (after September 1 but before freeze-up) - spring wheat (including semi-dwarf and durum).

Weeds and Staging:

Pre-emergent control of green and yellow foxtail.

Rates:

LIQUID OR DRY FLOWABLES

PRODUCT	RATES (L OR KG/ACRE)	
	LIGHT AND MEDIUM SOIL TEXTURE	HEAVY SOIL TEXTURE
Rival EC	0.49 to 0.57 L	0.65 L
Treflan EC	0.49 L	0.69 L
Bonanza 400	0.57 L	0.85 L

GRANULARS (WHEAT ONLY)

PRODUCT	RATES (KG/ACRE)
	All soils (2 - 8% organic matter)
Advance 10G/Rival 10G/Bonanza 10G	2.23
Treflan QR5*	4.45

*Spring applications of Treflan QR5 may also be made at the same rate during the months of April and May only in soils between 2 to 6% organic matter.

Cost (2004 suggested retail prices):

Advance 10G: $3.46 per kg.
Bonanza 400: $11.52 per L.
Bonanza 10G: $3.21 per kg.
Rival EC: $13.83 per L.
Rival 10G: $3.13 per kg.
Treflan EC: $14.97 per L.
Treflan QR5: $1.88 per kg.

Application Information:

Water Volume: Minimum 40 L per acre - liquid formulations only.

Pressure: 40 psi (275 kPa) - liquid and dry flowable formulations only.

Nozzles: Flat fan - liquid and dry flowable formulations only..

Incorporation:
Liquid or dry flowable formulations: Apply and incorporate in spring just after seeding. Incorporate to a depth of 1 to 1.5 inches (2 to 4 cm) into a trash free soil (80 percent black when viewed from above) using diamond or tyne type harrows operated at a speed of 6 mph (9 km/h). Incorporate twice, with the second incorporation at right angles to the first. The first incorporation should be performed immediately in the same direction of application. Both incorporations should be done within 24 hours of application. When tank mixing liquid formulations with Avadex BW, follow the same incorporation procedure.
Granular formulations: May be applied to standing or pre-worked stubble. Very heavy trash fields should be worked prior to application to allow product penetration to the soil surface. Incorporate with cultivators or disc implements only. Perform the first tillage operation within 24 hours of application. Incorporate at a working speed of 5 to 8 mph (8 to 13 km/hr) and to a depth of 2 to 3 inches (5 to 8 cm). Wait a minimum of 5 days, then incorporate a second time at right angles to the first. This second incorporation may be delayed until the following spring. Subsequent working should be no deeper than 2 to 3 inches (5 to 8 cm).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

When using fall applications of the granular formulations, crop injury will occur if soil conditions are not conducive to rapid wheat emergence (for example, cold or dry soil conditions). Weed control is best when soil moisture conditions are good. Rainfall has no direct effect on product activity. Flooding (3 to 5 days) will cause the rapid breakdown of the product, resulting in reduced weed control. Flooding for 3 weeks or more will result in total product breakdown and loss of weed control.

Tank Mixes:

Herbicides: None registered.

Fertilizer: Liquid trifluralin may be applied with liquid fertilizer (for example 28-0-0) as a carrier. Conduct a compatibility test before mixing.

Insecticides: None registered.

Note: The above mixes are those listed on the trifluralin labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: None. Flooding can reduce weed control.

Grazing: Do not graze the treated crop or cut for feed prior to crop maturity.

Aerial Application: Do not apply by air.

Recropping: Do not sow tame oats, canaryseed or small seeded forage grasses one year after spring applications or 21 months after fall applications.

Storage: Granular formulations must be stored in a cool, dry location, out of sunlight.

Rival EC - Do not store below 5°C.
Treflan E.C. - Do not freeze.
Bonanza 400 - Do not freeze.

Recommendations for liquid formulations: Crystalization of the active ingredient may occur at less than 5°C. To reconstitute, bring temperature to 15°C and shake well until no crystals are visible. This should be done before adding to the spray tank.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Keep out of reach of children.

Trifluralin (Broadleaf and Grassy Weed Control)

Herbicide Group – 3
(Refer to page 27)

Company:

Dow AgroSciences (Advance, Treflan)
NuFarm Agriculture Inc. (Rival)
United Agri Products (Bonanza)

Formulations:

Advance 10G (PCP#21742): 10% trifluralin formulated as a granular in 22.7 kg and 454 kg bags.

Bonanza 400 (PCP#16929): 400 g/L trifluralin formulated as an emulsifiable concentrate in 10 L, 115 L, 205 L containers.

Bonanza 10G (PCP#22744): 10% trifluralin formulated as a granular in 22.7 kg, 500 kg bags.

Rival EC (PCP#18612): 500 g/L trifluralin formulated as an emulsifiable concentrate in 9 L, 110 L, 200 L, and 900 L containers.

Rival 10G (PCP#18926): 10% trifluralin formulated as a granular in 22.7 kg, 567 kg bags.

Treflan EC (PCP#23933): 480 g/L trifluralin formulated as an emulsifiable concentrate in 9.45 L, 115 L containers.

Treflan QR5 (PCP#14545): 5% trifluralin formulated as a granular in 25 kg, 725 kg bags.

Crops and Staging:

Certain formulations are not registered for all the crops listed here. Refer to the specific product label for details. All products are for pre-plant incorporated use only.

Spring applied liquid, dry flowable or granular formulations - canola, peas, sunflowers, safflower (liquid formulations), dry beans, mustard, fababeans, alfalfa, sainfoin, sweet clover, soybeans, forage legumes (cicer milk-vetch, seedling alsike clover, red clover, bird's-foot trefoil).

Fall applied granular formulations - canola, peas, sunflowers, dry beans, mustard, fababeans, soybeans, barley, lentils and flax.

Trifluralin liquids - prior to planting shelterbelt transplants (elm, caragana, green ash, Scots pine).

Weeds and Staging:

For pre-emergent control of the following species:

Annual brome species
  (downy, Japanese)
Barnyard grass
Chickweed
Cow cockle
Green foxtail
Knotweed
Lamb's-quarters
Persian darnel
Pigweed
Purslane
Wild buckwheat*
Wild oats*
Yellow foxtail

*Some plants may escape herbicide treatment but are not competitive with the crop.

Costs (2004 suggested retail prices):

Advance 10G: $3.46 per kg.
Bonanza 400: $11.52 per L.
Bonanza 10G: $3.21 per kg.
Rival EC: $13.83 per L.
Rival 10G: $3.13 per kg.
Treflan EC: $14.97 per L.
Treflan QR5: $1.88 per kg.

Rates:

For use in canola, peas, sunflowers, dry beans, mustard, fababeans, seedling alfalfa (spring only), seedling sweet clover (spring only), soybeans.

PRODUCT	SOIL TYPE			
	Light soils with less than 6% organic matter		Medium to heavy soils with 6 to 15% organic matter	
	Spring	Fall	Spring	Fall
Advance 10G	Not registered	4.45 kg/acre	Not registered	5.67 - 6.88 kg/acre
Rival EC	0.65 L/acre	0.89 L/acre*	0.89 - 1.13 L/acre	1.13 - 1.37 L/acre*
Rival 10G	3.43 kg/acre**	4.45 kg/acre	4.45 - 5.67 kg/acre**	5.67 - 6.88 kg/acre
Treflan EC	0.69 L/acre	0.93 L/acre*	0.93 - 1.21 L/acre	1.21 - 1.37 L/acre*
Treflan QR5	6.9 kg/acre**	8.9 kg/acre	8.9 - 11.3 kg/acre**	11.3 - 13.8 kg/acre
Bonanza 10G	Not registered	4.45 kg/acre	Not registered	5.67 - 6.88 kg/acre
Bonanza 400	0.81 L/acre	1.11 L/acre*	1.11 L/acre	1.42 L/acre*

* Although liquid and dry flowable formulations are registered for fall application, this use is not recommended as tillage requirements before and after application will predispose fields to erosion.

**Spring applications of granular formulations are recommended for Manitoba only.

For use in barley (fall only), apply:

	SOIL TYPE					
	2 to 4% organic matter		4 to 6% organic matter		6 to 10% organic matter	
PRODUCT	Light Soil Texture*	Medium to Heavy Soil Texture**	Light Soil Texture*	Medium to Heavy Soil Texture**	Light Soil Texture*	Medium to Heavy Soil Texture**
Advance 10G	3.44 kg/acre	4.45 kg/acre	4.45 kg/acre	5.67 kg/acre	Not recommended	Not recommended
Rival 10G, Bonanza 10G	3.44 kg/acre	3.44 kg/acre	4.45 kg/acre	4.45 kg/acre	4.45 kg/acre	5.67 kg/acre
Treflan QR5	6.9 kg/acre	8.9 kg/acre	8.9 kg/acre	11.3 kg/acre	Not recommended	Not recommended

* Light textured soils can be defined as Sandy to Sandy loam.

** Medium to Heavy textured soils can be defined as loam to clay.

Rates: continued

For use in flax or lentils (fall only), apply:

PRODUCT	SOIL TYPE			
	Soils with 2 to 6% organic matter		Soils with 6 to 15% organic matter	
	Light Soil Texture*	Medium-Heavy Soil Texture**	Light Soil Texture*	Medium-Heavy Soil Texture**
Advance 10G, Rival 10G, Bonanza 10G	4.45 kg/acre	4.45-5.6 kg/acre***	5.67 kg/acre	5.67-6.88 kg/acre
Treflan EC	0.93 L/acre	1.21 L/acre	1.21 L/acre	1.21-1.38 L/acre
Treflan QR5	8.9 kg/acre	11.3 kg/acre	11.3 kg/acre	11.3-13.8 kg/acre

* Light textured soils can be defined as Sandy to Sandy-loam.
** Medium to Heavy textured soils can be defined as loam to clay.
*** Rates vary among products. Refer to product label for specific information.

Application Information:

Water Volume: Minimum 40 L per acre (liquid products only).

Pressure: 40 psi (275 kPa) (liquid products only).

Nozzles: Flat fan (liquid products only).

Incorporation: Granular formulations are recommended for use in fall or spring as a pre-plant incorporated treatment on broadleaf crops listed on the product label. The liquid formulations should be used only on soils free of lumps and relatively trash-free (75% black) and are recommended only for spring use. Granular formulations may be applied to standing or pre-worked stubble. Very heavy trash fields should be worked prior to application to allow product penetration to the soil surface. Do not use liquid or formulations of trifluralin as a pre-plant incorporated treatment in barley, as severe injury will result. Only the fall applications of Advance 10G, Rival 10G, Bonanza 10G or Treflan QR5 are registered for use as pre-plant incorporated treatments in barley. For fall application of granular formulations, work the chemical into the soil between September 1 and freeze-up. Use a discer or field cultivator (vibrating shank-type). Disc implements are preferred on stubble. Set equipment to cut at 3 to 4 inches (8 to 10 cm) depth. The initial incorporation should be done within 24 hours of application.

The second incorporation should be done at right angles to the first. The second incorporation may be delayed until spring, except when planting barley, flax or lentils; for these crops both incorporations must be done in fall. Delay the second incorporation 5 days for better weed control. This will allow greater release of the chemical onto soil particles and assure more even distribution. Fall application of granular trifluralin on flax, lentils or barley is not recommended on soils prone to erosion, as the 2 fall incorporations necessary in these crops may leave soils vulnerable to wind or water erosion.

For spring application of liquid and granular formulations, work the chemical into the soil prior to seeding by setting the implement at 3 to 4 inches (8 to 10 cm) cutting depth. The first incorporation must be done within 24 hours of application. The second incorporation must be done at right angles to the first. If incorporating granular trifluralin, delay the second incorporation for 3 days after the first to achieve better weed control.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Injury to flax, barley or lentils may occur if soil and weather conditions are not conducive to rapid crop emergence (cold or dry soils at the time of seeding and crop emergence).

To minimize crop injury, seed into a firm, moist seed bed. Use a press drill or hoe-drill. Sow barley no deeper than 2 inches (5 cm). Sow lentils and flax no deeper than 1.5 inches (4 cm).

Less than acceptable weed control will result if dry conditions prevail at the time of weed emergence.

Rainfall has no direct effect on products' activity. Flooding (3 to 5 days) will cause rapid breakdown of the product resulting in reduced weed control. Flooding for 3 weeks or more will result in total breakdown of the product resulting in loss of weed control.

Tank Mixes:

Herbicides:

Fababeans:
Sencor 75 DF or Sencor 500 (Treflan EC only).

Peas:
Sencor (Treflan EC only).

Soybeans:
Sencor 500 (use only on KG 20 or McCall) (Treflan EC only). This mixture may cause injury to soybeans if seedlings are weak or slow to emerge.

Dry beans (white and red kidney only):
Eptam 8-E (Treflan and Rival liquids only)

Fertilizers: Liquid product may be applied with liquid fertilizer as a carrier. Before the herbicide is added to the tank, compatibility of the herbicide to liquid fertilizer should be tested following instructions on the herbicide container. Trifluralin liquids may be blended with dry bulk fertilizers (do not mix with nitrate fertilizers). Check label for blending instructions.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Trifluralin labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: None. Flooding may reduce weed control.

Grazing: Do not graze the treated crops or cut for feed prior to crop maturity.

Recropping: Oats, canaryseed, and small-seeded grasses may be affected the year after treatment. Corn is sensitive at higher rates of application. Damage to wheat can occur if the crop is seeded into land that has been treated during the previous 21 months with trifluralin products and has received abnormally low amounts of precipitation. Damage is worse if conditions are not conducive to rapid emergence of the wheat (for example, if the crop is seeded deep or if soil conditions remain cool during emergence). Damage tends to be greater on fields treated with granular formulations.

Aerial Application: Do not apply by air.

Storage: Granular formulations must be stored in a cool, dry location, out of sunlight.
Rival EC - Do not store below 5°C.
Treflan EC, Bonanza 400 - Do not freeze.
Bonanza 400 - Do not freeze.

Recommendations for liquid formulations: Crystalization of the active ingredient may occur at less than 5°C. To reconstitute, bring temperature to 15°C and shake well until no crystals are visible. This should be done before adding to the spray tank.

Tank Cleaning:

Refer to page 14

Hazard Rating:

Keep out of reach of children.

Triumph Plus

Herbicide Group – 1, 2, 4
(Refer to page 27)

Company:

Bayer CropScience

Formulation:

The Triumph Plus package contains 2 components:

Triumph FM (PCP#22348): 56 g/L fenoxaprop-p-ethyl and 256 g/L MCPA ester formulated as an emulsifiable concentrate.
Container size – 2 x 13.3 L.

Plus (PCP#21645): 75% thifensulfuron methyl formulated as a dry flowable.
Container size – 2 x 162 g (imbedded in Triumph FM jug).

Crops and Staging:

Spring wheat (not including durum)- 2 to 6 leaf stage, maximum 3 tillers

Seedling forage grasses for seed and forage, treat at the 2 to 4 leaf stage:

Bromegrass (meadow and smooth)

Fescue (creeping red, tall)

Wheatgrass (intermediate, northern, slender, tall, western)

Crop injury (yellowing, stunting and in some instances pinching of tillers) can occur with applications of Triumph Plus made beyond the recommended leaf stage. Roblin wheat is particularly sensitive to applications of Triumph Plus made at leaf stages beyond the recommended range. To minimize the risk of injury, apply the product within the recommended range.

Weeds and Staging:

Grass weeds:

Apply at the 1 to 6 leaf stage with a maximum of 2 tillers. Target the 2 to 3 leaf stage for optimum control:

Foxtail (green and yellow)
Wild oats

Broadleaf weeds:

Apply at the 2 to 4 leaf stage or less than 4 inches (10 cm) across, unless otherwise stated:

Annual sunflower
Ball mustard
Burdock
Canada thistle*
Chickweed (1 to 6 leaf)
Cocklebur
Corn spurry
Cow cockle
Field horsetail**
Flixweed***
Green smartweed
Hemp-nettle
Hoary cress**
Kochia
Lady's-thumb
Lamb's-quarters
Mustards (except dog and green tansy)
Plantain**
Prickly lettuce
Ragweeds
Redroot pigweed
Russian pigweed
Russian thistle
Stinkweed***
Shepherd's purse***
Vetch
Volunteer canola^
Wild buckwheat (1 to 3 leaf)
Wild mustard
Wild radish

* Suppression only
** Top growth control only
*** Spring seedlings only
^ May not give satisfactory control of CLEARFIELD canola varieties

Cost (2004 suggested retail price):

$20.96 per acre.

Rates:

The Triumph Plus package will treat 40 acres.

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 23 to 45 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: 80° or 110° stainless steel flat fan nozzles tilted forward at a 45° angle. Screens 50 mesh or larger.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Do not apply to crop that is stressed by severe heat, frost, low fertility, drought, water-saturated soil, disease or insect damage as crop injury and poor weed control may result.

Tank Mixes:

Herbicides: In spring wheat (not durum): Lontrel (85 mL/acre).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Do not tank mix with any other chemical additive, pesticide or fertilizer. Do not use a surfactant with this tank mix. An interval of 7 days prior to application, or 4 days after application with Triumph Plus, is required before any other pesticide can be applied.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: Within 2 hours may reduce weed control.

Grazing: Do not graze treated crops or cut for feed prior to crop maturity. Treated forages are not to be harvested for feed or grazed the year of application.

Preharvest Interval: Do not apply within 80 days of harvest.

Recropping: No restrictions the year after treatment.

Aerial Application: Do not apply by air.

Storage: Do not freeze. If stored for 1 year or longer, shake well before using.

Environment: Do not apply within 15 m of a body of water or wetland area.

Tank Cleaning:

The "Plus" component of Triumph Plus can cause severe injury to sensitive crops at very low concentrations. Sprayers used to apply Triumph Plus should be flushed out immediately after use. The manufacturer recommends that sprayers used to apply this product be flushed twice with a water/household ammonia (1L of 3% household ammonia per 100 L water) rinse prior to using the sprayer on sensitive crops. All nozzles, screens and filters should be removed and cleaned after applying this product.

Refer to page 14 for additional information.

Hazard Rating:

◈ Warning Poison

Skin and Eye Irritant

Trophy

Herbicide Group – 4,2
(Refer to page 27)

Company:

Nufarm

Formulation:

The Trophy package has 2 components:

Trophy A (PCP#27246): 180 g/L fluroxypyr formulated as an emulsifiable concentrate.

Trophy B (PCP#27245): 500 g/L MCPA ester formulated as an emulsifiable concentrate.

Container size: Trophy A – 4.8L, Trophy B – 9L.

Crops and Staging:

Spring wheat (NOT including durum) & barley - 3 leaf up to flag leaf fully emerged stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 2 to 4 leaf stage, unless specified include:

Burdock
Cleavers (1-4 whorls)
Clover (sweet)
Cocklebur
Flixweed
Hemp Nettle (2-6 leaf)
Kochia
Lamb's-quarters
Mustards (except dog and tansy)
Prickly lettuce
Ragweed
Shepherd's-purse
Stinkweed
Sunflower (annual)
Vetch
Volunteer canola
Volunteer flax
Wild radish
Wild mustard

Weeds suppressed include:

Smartweed (green)
Stork's-bill
Wild buckwheat (1-4 leaf)

Cost (2004 suggested retail price):

$10.05 per acre.

Rate:

Trophy A: 0.24 L per acre

Trophy B: 0.45 L per acre.

One case of Trophy treats 20 acres. Make only one application per year.

Application Information:

Water Volume: Minimum 40 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Trophy activity is influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought, heat or cold stress) or if extremely heavy infestations exist.

Tank Mixes:

Herbicides:

In barley only:
Achieve Liquid (0.2 L/acre) plus Turbocharge adjuvant.*

In spring wheat only (NOT including durum):
Assert (0.53 to 0.65 L/acre) plus pH adjuster
Horizon (95 to 115 mL/acre) plus Score adjuvant.*
Puma^{120} Super (0.31 L/acre)

*Temporary crop injury (Achieve only) or reduced wild oat control may occur with these tank mixes.

Insecticides: None registered.

Fertilizers: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 13.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: Do not graze, harvest forage or cut hay within 7 days of application.

Preharvest: Leave at least 60 days from application to harvest.

Recropping: Wheat, barley, oats, rye, forage grasses, flax, canola, mustard, lentils and peas may be grown the year after a Trophy application. There are no recropping restrictions the second year after application.

Aerial Application: Do not apply by air.

Storage: May be frozen. If frozen, bring to room temperatures and agitate before use.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

DangerPoison.

Warning Eye Irritant.

Caution Skin Irritant.

Ultim*

Herbicide Group – 2

(Refer to page 27)

***(For use only in Manitoba)**

Company:

E. I. duPont Canada (PCP#24736)

Formulation:

37.5% rimsulfuron and 37.5% nicosulfuron formulated as a water dispersible granule. Container size – 134.8 g (4 x 33.7 g water soluble bags).

Crops and Staging*:

The following field corn hybrids at the 1 to 4 leaf stage:

Pioneer Brands 39K72, 39K73, 39W54, 39M27. For use in Manitoba only.

** NOTE - Since applications to corn in Manitoba has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance.* ***Application to corn is at the risk of the user.***

Weeds and Staging:

No information is provided on the label for leaf staging. The manufacturer recommends the following staging:

wild oats – 3 to 6 leaf stage

green foxtail, yellow foxtail, barnyard grass, volunteer cereals – 1 to 6 leaf stage (up to 2 tillers)

quackgrass – 3 to 6 leaf stage (with extended leaf 4 to 8 inches long)

redroot pigweed – 2 to 6 leaf stage

Contact the manufacturer for additional weeds not listed on the label.

Cost (2004 suggested retail price):

$23.71 per acre

Rates:

Apply Ultim 75 DF at 13.5 g/acre.

One water soluble bag of Ultim will treat 2.5 acres (10 acres per container).

Add a non-ionic surfactant (AgSurf, Agral 90, Citowett Plus) at 0.2 L per 100 L of spray solution. Apply Ultim with 24 hours of mixing, as product degradation may occur resulting in reduced weed control. Refer to the product label for complete mixing instructions.

Application Information:

Water Volume: Minimum 40 L per acre; for best results apply 56 to 77 L per acre.

Nozzles and Screens: Flat fan nozzles with 50 mesh or larger screens.

Effects of Growing Conditions:

Rapid fluctuations in temperature (greater than 20°C difference within 24 to 36 hours) will stress the corn crop. For maximum crop safety, allow 48 to 72 hours for the corn to acclimatize before applying Ultim.

Apply ONLY when the temperature in the 24 hours before AND after application is between 5°C and 28°C. Temperatures beyond this range increase the potential for crop injury. Separate applications of Ultim herbicide followed by a broadleaf herbicide (minimum 12 hours later) will reduce the potential for injury.

WARNING: Crop injury may result if application is made to corn that has been stressed by abnormally hot, humid or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If corn has been injured by frost, wait 48 to 72 hours before applying Ultim.

Tank Mixes:

Herbicides: None registered.

Insecticides: None registered. Ultim should NOT be applied to corn that has been treated with Counter, Cygard, Thimet, Lorsban, or Di-Syston. Leave seven days between the application of Ultim and that of a foliar organophosphate insecticide.

Fungicides: None registered.

Note: The above mixes are those listed on the Ultim label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: Within 2 to 4 hours of application may reduce weed control.

Grazing: Do not graze treated crops or cut for hay.

Preharvest: Leave at least 30 days from application to harvest.

Recropping: Field corn, winter wheat and spring barley may be planted the year following Ultim application. Perform a field bioassay before planting any other crops, or where Ultim is more persistent (sandy soils, with low organic matter and pH greater than 7),

Aerial Application: Do not apply by air.

Storage: Store product in original containers in a secure, dry area, away from other pesticides, food, or feed.

Environment: Leave a 5 m buffer between the last spray path and woodlots or shelterbelts. Leave a 10 m buffer before wetland areas or water bodies.

Tank Cleaning:

Drain tank and hose down interior surfaces. Flush tank, hoses, boom, and nozzles with clean water for a minimum of 5 minutes. Fill spray tank with a water-ammonia cleaning solution (1 litre of minimum 3% household ammonia for every 100 L of water). Flush hoses, boom and nozzles with the cleaning solution, then add more water to completely fill the tank. Circulate for 15 minutes, then flush hoses, boom and nozzles with the cleaning solution, and drain the tank. Remove and clean the nozzles and screens separately in a bucket containing a cleaning solution as above. Repeat the above process and then thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom. Prior to using the sprayer for the next application, flush the tank, boom and hoses for 5 minutes with fresh water. Do not clean equipment where cleaning solution could flow towards water bodies, ditches, cropland, shelterbelts, or areas where people are likely to frequent or walk.

For additional information, refer to page 14.

Hazard Rating:

Warning – Eye Irritant

Caution – Skin Irritant

Unity*

***(For use only in Saskatchewan)**

Herbicide Group – 2,6

(Refer to page 27)

Company:

Bayer CropScience

Formulation:

The Unity package contains two components:

Unity 280 EC (PCP#24497): 280 g/L bromoxynil formulated as an emulsifiable concentrate.

Unity 75 WG (PCP#24596): 75% triasulfuron formulated as a dispersible granule.

Container size - one case of Unity contains 2 x 4 L jugs of Unity 280 EC and 4 x 43 g water soluble bags of Unity 75 WG.

Crops and Staging:

Spring wheat (including durum), and barley - 2 leaf to just prior to flag leaf emergence. For use in the brown and dark brown soil zones of Saskatchewan only.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 2 to 4 leaf stage:

Chickweed
Cleavers
Cow cockle
Flixweed
Green smartweed
Hemp-nettle
Kochia
Lady's-thumb
Lamb's-quarters
Redroot pigweed
Shepherd's-purse
Stinkweed
Wild buckwheat
Wild mustard
Volunteer canola

Caution:

Several crops are sensitive to low concentrations of Unity 75 WG in the soil. Therefore, careful consideration should be given to crop rotation plans prior to use. The Unity tank mix is registered for use in Saskatchewan in the brown and dark brown soil zones only. The degradation of Unity 75 WG in the soil is affected by rainfall, soil temperature and soil pH. If you are considering using this product, be sure to read the recropping restrictions outlined in the product label.

Cost (2004 suggested retail price):

$7.25 per acre.

Rates:

Unity 75 WG: 4.3 g per acre

Unity EC: 0.2 L per acre

One case treats 40 acres. Add a recommended non-ionic surfactant such as Citowett Plus, Agral 90, Agsurf, Super Spreader or Companion at 0.25 L per 100 L spray solution.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Pressure: 40 psi (275 kPa).

Nozzles: Flat fan. 50 mesh screens or coarser.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Adequate control may not be achieved under unfavourable conditions such as drought, flooding or prolonged temperature extremes.

Tank Mixes:

Herbicides:

In Spring wheat (NOT including durum) and Barley:
2,4-D ester (0.34 l/acre – 500 g ai/L formulation)
Lontrel (0.11 L/acre)
MCPA ester (0.34 L/acre)

In Spring wheat (NOT including durum):
Puma® Super (0.4 L/acre)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the Unity label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions.

Restrictions:

Rainfall: Do not apply if rainfall is expected within 2 hours.

Grazing: Do not graze treated crops or cut for hay until 30 days after application.

Recropping: Some crops have shown a high sensitivity to low residues of Unity 75 WG in the soil. The risk of injury to rotational crops is affected by a number of factors including soil factors (soil type and soil pH), environmental factors (soil temperature, precipitation), crop species, and recropping interval. Breakdown of Unity 75 WG in soil is more rapid in high moisture, high temperature, and low pH soils.

To avoid injury to subsequent crops after an application of recommended rates of the Unity tank mix, the following recropping intervals outlined in the following table should be observed:

Recropping Restrictions:

CROP	MINIMAL INTERVAL (MONTHS)		
	Soil pH 6.4 or less	**Soil pH 6.5 to 7.4**	**Soil pH 7.5 or greater**
Spring wheat (hard red, Canada prairie, extra strong)	No Restrictions	No Restrictions	No Restrictions
Barley, oats, durum wheat	10	10	10
Flax	10	10	22
Peas	10	22	22
Canola	10	22	34
Canaryseed	22	22	22
Mustard	22	22	34
Lentils, sunflowers, all other crops	Bioassay	Bioassay	Bioassay

A field bioassay involves growing test strips of the crop(s) intended for production the following year in fields previously treated with the Unity tank mix. Crop response will dictate whether or not to rotate to the crop(s) used in the test strip. See Bioassay Guidelines section of product label. Apply the Unity tank mix only once per year to the same field.

Restrictions continued:

Aerial application: Do not apply by air.

Storage: Do not freeze.

Environment: Leave at least a 15 m buffer zone around wildlife habitat. Do not apply to irrigated land.

Tank Cleaning:

Unity can cause injury to sensitive crops at low concentrations. Clean sprayer immediately after use. Do not clean near desirable vegetation, wells or other water sources. Drain and flush tank and boom with water and household detergent. Add 1 L of 3% household ammonia for every 100 L of water to spray tank and agitate for 15 min. before flushing through nozzles. Wash nozzles and screens in the water/ammonia mix in pail. Rinse system with clean water for 5 minutes.

For additional information, refer to page 14.

Hazard Rating:

Warning Poison

Velpar DF

Herbicide Group – 5
(Refer to page 27)

Company:

E. I. duPont Canada (PCP#25225)

Formulation:

75% hexazinone formulated as a water dispersible granule
Container size: 2 kg.

Crops and Staging:

Established alfalfa for seed. Apply in late fall prior to freeze-up when alfalfa is dormant or in early spring before alfalfa growth resumes. Apply only on alfalfa that has been established for 18 months or longer. If burning or irrigation is to be carried out, do not apply until these operations have been completed. Do not apply to frozen ground.

Weeds and Staging:

Application stage is dictated by the crop above

Dandelion
Narrow-leaf hawk's beard
Sow-thistle
Quackgrass
Scentless chamomile

Cost (2004 suggested retail price):

$22.71 to $46.30 per acre.

Rates:

0.272 to 0.544 kg/acre. Use the lower rate on medium-textured soils with low organic matter. Do not apply to soils with less than 1% organic matter.

Application Information:

Water Volume: 81 L per acre.

Pressure: 30 to 40 psi (200 to 275 kPa).

Nozzles: Flat fan.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Adequate soil moisture is required for activation of the product.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Rainfall is beneficial for activation of the product.

Grazing: Do not graze the treated crop or cut for feed.

Recropping: Insufficient information is available. Velpar residues will persist for 2 or more years after application.

Aerial Application: Do not apply by air.

Storage: May be frozen.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

Danger Corrosive

Caution Flammable

Venture L

Herbicide Group – 1
(Refer to page 27)

Company:

Syngenta (PCP#21209)

Formulation:

125g/L fluazifop-p-butyl formulated as an emulsifiable concentrate. Container size 2 x 8 L.

Crops and Staging:

Preharvest intervals must be followed to avoid unacceptable residues of fluazifop-p-butyl in harvested crops.

Crop	Crop Stage	Harvest Interval
Alfalfa	No restrictions	41
Bird's-foot trefoil and red clover	No restrictions	Do not graze or feed in year of treatment
Canola	Do not apply past 5th leaf	80
Creeping red fescue	2-5 leaf	For seed only
Flax, Solin	No restrictions	80
Lentil	Do not apply past 6th node	70
Lupins	No restrictions	80
Mustard	Do not apply past 5th leaf	75
Peas	No restrictions	66
Potatoes	No restrictions	90
Soybeans	Up to 3rd trifoliate leaf	90
Sunflowers	No restrictions	120

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Cost (2004 suggested retail price):

$26.40 per L.

Application Information:

Water Volume: 20 to 81 L per acre.

Pressure: 30 to 60 psi (200 to 400 kPa).

Nozzles: 80° or 110° flat fan nozzles.

How it Works:

Refer to Table 2 on page 29.

Effects of Growing Conditions:

Less than acceptable weed control may be expected if weeds are under stress because of excessive moisture, drought, or cool weather.

Weeds, Rates and Staging:

Weed	Weed Stage	Rate (L/acre)	Acres per 8 L
Volunteer corn	2 to 5 leaf stage	0.24	33
Volunteer wheat and barley, Persian darnel, barnyard grass	2 to 5 leaf stage	0.32	25
Wild oats	2 to 5 leaf stage	0.40	20
Quackgrass suppression in canola and soybeans	3 to 5 leaf stage	0.40	20
Green and yellow foxtail	2 to 4 leaf stage	0.57	14
Quackgrass suppression (all crops)	3 to 5 leaf stage	0.80	10

Tank Mixes:

Herbicides:

In canola:
Muster (Argentine varieties only)
Lontrel

In alfalfa, bird's-foot trefoil & red clover:
2,4-DB

In creeping red fescue:
Ally

In soybean:
Basagran + Assist

In potato:
Sencor (reduced grass control from Venture L may result)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

When applying other broadleaf herbicides in the same field, apply the product no sooner than 3 days after application of Venture.

Note: The above mixes are those listed on the Venture L label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Restrictions:

Rainfall: Activity is not affected by rainfall if it occurs at least two hours after application.

Grazing: Do not graze crops or cut for hay in the year of treatment.

Preharvest Interval: See crop stage section.

Recropping: Seed only broadleaf crops listed on this label if seeding must occur within 60 days of application.

Aerial Application: Do not apply by air.

Storage: Product is not affected by temperatures below 0°C.

Environment: Do not apply within 15 m of fish-bearing waters and wildlife habitats.

Tank Cleaning:

Refer to page 14.

Hazard Rating:

☠ Caution Poison

Experimental studies have shown that the active ingredients in this product may cause birth defects in laboratory animals.

Women capable of bearing children should avoid contact with this product.

Special Weed Problems

This section identifies specific weeds and some herbicides recommended for control. Refer to the particular crop section or the product label for information on specific products that may be used on the crops and for application instructions.

Absinthe

2,4-D LV Ester (500 g/L) - In grass pastures with no legumes, spray 1.82 L/acre in late June, prior to flowering. Re-treat regrowth in late summer when plants have 6 to 10 inches (15 to 25 cm) of new growth. More than 1 season of treatment may be required.

Banvel II - In grass pasture and rangeland only, apply 0.5 L/acre in 20 to 30 gallons (90 to 135 L) per acre for topgrowth control when leaves are fully expanded.

Alders

2,4-D LV Ester (500 g/L) - In grass pastures and non-crop land, apply 1.78 L/acre to the foliage of actively growing brush.

Banvel II + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply Banvel II at 2.1 L per 1,000 L of water with 2,4-D LV ester or amine at 4.0 L per 1,000 L of water to the foliage of actively growing brush in the spring or early summer and wet the foliage until the point of runoff.

Aspen Poplar (Trembling Aspen)

Banvel II + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply Banvel II at 1.32 L/acre with 2,4-D LV ester or amine at 1.78 L/acre in 20 gallons/acre (90 L/acre) water to the foliage of actively growing brush in spring or early summer.

Remedy EC - In grass pasture and rangeland apply 1.62 to 3.24 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Single stem foliar and backpack methods are also available - See Remedy section for details.

Baby's Breath (Perennial)

Banvel II - In grass pastures with no legumes, apply 3.72 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water when actively growing.

Biennial Wormwood

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing plants.

Black Medic

Bromoxynil+MCPA ester, /Dichlorprop+2,4-D,/Mecoprop, /dicamba+mecoprop+MCPA, /2,4-D amine or LV ester - Apply in registered crops at registered rates to black medic in the 1 to 4 leaf stage for suppression only.

Chokecherry

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Remedy EC - In grass pasture and rangeland apply 1.62 to 3.24 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Single stem foliar and backpack methods are also available - See Remedy section for details. May need the higher rates for control and retreatment the following year.

Common Tansy

Glyphosate - Apply at 1.9-2.8 L/acre in 10 gallons of water/acre (40 L/acre) to actively growing plants that are 8-10 inches (20-25 cm) tall (summerfallow, stubble and noncropland).

Escort - In pastures, rangeland and rough turf, apply 8 g/acre in 10 to 20 gallons/acre (45 to 90 L/ acre) of water to actively growing plants of less than 4 inches (10 cm) tall. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Curled Dock

Banvel II - As a patch treatment or in pasture and rangeland, apply 0.92 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds for top growth control.

Glyphosate - As a spot treatment, apply 2.83 to 4.86 L/acre in 10 gallons/acre (45 L/acre) water when most plants have reached the early bud stage. Do not disturb treated plants for at least 10 days following treatment.

MCPA amine, 2,4-D amine - Apply 0.445 to 0.69 L/acre of formulations containing 500 g/L MCPA or 2,4-D amine to give top growth control.

Dichlorprop + 2,4-D - 0.71 L/acre for suppression before plants are 2 inches (5 cm) tall.

Remedy EC - In grass pasture and rangeland apply 0.4 to 1.62 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Backpack method is also available - See Remedy section for details.

Diffuse Knapweed

Banvel II - In grass pastures, rangeland and non-crop land, apply Banvel II at 1.86 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds.

Tordon 22K - In rangeland and grass pasture, apply 0.91 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Downy Brome and Japanese Brome

Glyphosate - Prior to crop emergence, apply 0.51 to 0.77 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water before downy brome is 6 inches (15 cm) in height.

Rustler - Prior to crop emergence, apply 1.0 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water between emergence and heading of downy brome.

Trifluralin - Apply at recommended rates for weed control in broadleaf crops prior to emergence.

Field Bindweed

Banvel II - As a patch treatment or in rangeland, apply 1.0 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water. Apply when field bindweed is in the flowering stage and allow 3 weeks after treatment before resuming normal summerfallow tillage.

Basagran - In labelled crops, apply 0.71 L/acre followed by 0.71 L/acre 7 to 10 days later. Apply in 20 to 35 gallons/acre (90 to 160 L/acre) water before field bindweed has developed a dark green colour and before it has begun trailing. Use a recommended surfactant (see recommendations under the appropriate crop).

2,4-DB - As a spot treatment in labelled crops apply 2.83 to 4.86 L/acre in 10 gallons/acre (45 L/acre) water at the bud stage. Do not disturb plants for at least 10 days following treatment. Heavy rainfall within 2 hours of application may wash chemical off the foliage and a repeat treatment may be required. Rainfall occurring within 6 hours after application may reduce control.

2,4-D amine - In grass pastures containing no legumes or as a spot treatment, apply 1.82 L/acre of formulations containing 500 g/L 2,4-D amine at early flowering stage.

Glyphosate - As a spot treatment, apply 2.8 to 4.9 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) at the full bloom stage or beyond. Allow 7 or more days after application before tillage.

Remedy EC - In grass pasture and rangeland apply 0.4 to 1.62 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Backpack method is also available - See Remedy section for details.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Field Horsetail

Amitrol 240 - Apply 5.0 to 6.7 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is young and actively growing.

MCPA amine, potassium and sodium salt mixtures - Apply 0.57 L/acre of formulations containing 500 g/L MCPA after the weeds have fully emerged for top growth control. May be used in wheat, oats, barley, flax and rye.

Foxtail Barley

Glyphosate - Prior to crop emergence, apply 1 to 2 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water to foxtail barley at the seedling to heading stage. Late fall applications may provide better control of established plants than spring applications.

Glyphosate - In Roundup Ready canola, apply 2 applications, each at 0.5 L/acre, for season long control.

Gramoxone - Apply 2.23 L/acre in 98 gallons/acre (445 L/acre) water or 75 mL in 2.2 gallons (10 L) water/1076 square feet (100 sq. m) for top growth control only.

Kerb 50-W - Apply 0.364 to 0.445 kg/acre product in 20 gallons/acre (90 L/acre) water between October 1 and freeze-up. Use the lower rate on grey-wooded soils or where perennial bluegrass or fescues are the predominant pasture species. Do not use Kerb for foxtail barley removal in seed grass stands or desired foliage stands of timothy or fescue grass species. At recommended rates, pasture stands of perennial bluegrass and fescue may be reduced by 10 to 15 percent. Where perennial bluegrass and fescues are the dominant pasture species, use the lower rate of Kerb. Spray overlaps may seriously harm desirable pasture grass species. Where the grass stand comprises mostly foxtail barley and reseeding to a desirable grass species is required, delay seeding into the Kerb-treated soil until the end of June. Do not harvest or graze within 60 days of application with Kerb. Avoid using Kerb on soils having more than 6 percent organic matter.

Rustler - Prior to crop emergence, apply 1.26 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water to foxtail barley before initiation of the seed head for suppression only.

Sundance - Apply at 11 g/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water prior to flowering.

Goat's-Beard

2,4-D amine - Apply 0.91 L/acre of formulations containing 500 g/L in early fall or early spring.

Dichlorprop + 2,4-D - Apply 1.62 L/acre in early spring or fall.

Banvel II - In grass pasture and rangeland only, apply 1.86 L/acre in 20 to 30 gallons (90 to 135 L) per acre when leaves are fully expanded.

Gumweed

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 0.89 L/acre to the foliage of actively growing plants.

Hemp Dogbane

2,4-D amine or LV ester - Apply 1.38 to 1.82 L/acre of formulations containing 500 g/L 2,4-D in fall before frost and while plant leaves are green.

Glyphosate - Apply 2.83 to 4.86 L/acre when hemp dogbane is in the early bud stage. Apply in 10 gallons/acre (45 L/acre) water. Do not disturb treated plants for at least 7 days after application.

Hoary Cress

Amitrol 240 - For non-selective patch treatment in pastures and non-crop land, apply 8.9 to 13.8 L/acre.

Glyphosate - As a spot treatment in labelled crops, apply 2.83 to 4.86 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water when most plants have reached the early bud stage. Do not disturb treated plants for at least 10 days following treatment.

Leafy Spurge

Amitrol 240 - Apply 15.2 to 18.5 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is between the late stages of flowering and early seed development.

Banvel II - Apply 0.84 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water for top growth control when the weed is actively growing. Patch treatment or pasture.

2,4-D amine - Apply 1.82 L/acre of formulations containing 500 g/L 2,4-D at early flowering stage. Repeat at least once to new growth later in the season. Control of established plants and new seedlings will require continued applications for a period of at least 4 to 5 years.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Locoweeds, Lupines, and Milk-vetches

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 to 2.75 L/acre at the full bloom stage.

Remedy EC - To control vetches in grass pasture and rangeland apply 0.4 to 1.62 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Backpack method is also available - See Remedy section for details.

Milkweed

Amitrol 240 - Apply 7.6 to 11.3 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures in the early summer when all the shoots have emerged.

Glyphosate - When making preharvest applications, use 1.0 L/acre. For patch treatments, apply 4.86 L/acre in 10 gallons/acre (45 L/acre) water. Apply when most plants have reached the bud to bloom stage. Reduced results may occur on plants treated after full bloom as not all milkweed plants reach the required stage of growth at the same time. Repeat treatments may be required. Do not disturb plants for 10 days following treatment. Do not apply to plants covered with dust.

Narrow-leaf Hawk's-beard

2,4-DB - Apply to forage legume crops at recommended rates at the 2 to 4 leaf stage of narrow-leaf hawk's-beard, after legume growth in the fall has stopped.

2,4-D LV ester (500 g/L) - In fall stubble, apply 0.57 to 0.90 L/acre to fall rosettes.

Express Pack - In registered crops and summerfallow, apply at registered rates to spring seedlings less than 4 inches (10 cm) across or to fall rosettes.

Glyphosate - Prior to crop emergence, apply 0.51 to 0.77 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water. Use the high rate if narrow-leaf hawk's-beard is between 3 and 6 inches (8 to 15 cm) in height.

Glyphosate - In Roundup Ready canola, apply 0.5 L/acre at the 0 to 6 leaf stage. Not all products are registered. Check glyphosate pages.

Pasture Sage

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.2 L/acre to the foliage of actively growing plants.

Banvel II - In grass pastures, rangeland and non-crop land, apply Banvel II at 1.86 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds.

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Perennial Smartweed

Glyphosate - Apply 2.0 L/acre in 10 gallons/acre water. Apply when vines are a minimum of 8 inches (20 cm) tall, but before flowering.

Poplar

Glyphosate - As a non-selective spot treatment, apply 1.21 to 2.43 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in the summer through early fall when brush is actively growing.

Remedy EC - In grass pasture and rangeland apply 1.62 to 3.24 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Single stem foliar and backpack methods are also available - See Remedy section for details.

Banvel II + 2,4-D - In grass pasture and rangeland only, apply Banvel II at 2.1 L plus 2,4-D 500 amine at 4 L or 2,4-D 600 ester at 3.3 L per 220 gallons (1000 L) of water and apply by wand to the point of runoff when leaves are fully expanded.

Poverty Weed

Banvel II - As a spot treatment or in grass pasture or rangeland apply 1.86 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water when weed is actively growing. Banvel II at 0.61 L/acre will provide only top growth control.

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Prairie Everlasting, Prairie Sage

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing plants in the early fall, and repeat in the spring.

Purple Loosestrife

(dryland situations only)

Glyphosate - Apply 2.43 L/acre in 30 to 60 gallons/ acre (135 to 270 L/acre) water when purple loosestrife is actively growing and at or beyond the bloom stage. If using hand held equipment, apply a 1 to 2 percent solution until plants are wet. Use a 33 percent product solution if using a wiper applicator. Do not treat plants over open water. If possible, remove and destroy the flower heads before treatment to ensure prevention of seed set. For large monocultures of purple loosestrife, gradually work from the periphery inward over a number of years to allow competing vegetation to invade the treated area. Sprayed areas should be monitored for new seedlings to prevent re-infestation of purple loosestrife.

Red Bartsia

2,4-D amine or LV ester - Apply 0.57 L/acre of formulations containing 500 g/L 2,4-D in 10 gallons/acre (45 L/acre) water. On hayland, treat within 10 days after first cutting. Roadsides and pastures should be sprayed as soon as the red bartsia appears, usually in early July. Repeat treatment if necessary for later germination.

Roses

Banvel II + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply Banvel II at 1.48 L/acre with 2,4-D LV ester or amine at 1.78 L/acre to the foliage of actively growing brush in the spring or early summer.

Escort - In pasture and rangeland, apply Escort at 0.012 kg/acre with non - ionic surfactant at 0.2 L per 100 L spray solution in 10 to 20 gallons/acre (45 to 90 L/acre) water. Apply between mid-June and mid-August after the brush has leafed out, but before the leaves begin to turn their fall colours.

Remedy EC - In grass pasture and rangeland, apply 1.62 to 3.24 L/acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Single stem foliar and backpack methods are also available - See Remedy section for details.

Russian Knapweed

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Banvel II - In grass pasture and rangeland only, apply 3.72 L/acre in 20 to 30 gallons (90 to 135 L) per acre when leaves are fully expanded.

Saskatoon

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Scentless Chamomile

Ally plus 2,4-D - Apply 3 g/acre Ally plus 0.34 to 0.45 L/acre 2,4-D LV ester or amine (500 g/L formulations) in 10 gallons/acre (45 L/acre) water for control of scentless chamomile up to the 4 leaf stage in wheat, barley, and creeping red fescue. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Banvel II - Apply 0.51 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds for top growth control.

Bromoxynil+MCPA ester - Apply in registered crops at label rates when scentless chamomile is in the 2 to 4 leaf stage.

Curtail M - In registered crops, apply 0.81 L/acre in 10 gallons/acre (45 L/acre) water when scentless chamomile is actively growing and in the 2 to 4 leaf stage.

Escort - In pastures, rangeland and rough turf, apply 8 g/acre in 10 to 20 gallons/acre (45 to 90 L / acre) of water to actively growing plants of less than 4 inches (10 cm) tall. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Liberty - In registered crops, apply 1.1 L/acre to plants up to 4 inches (10 cm) in height.

Lontrel - In registered crops, apply 0.23 L/acre in 10 gallons/acre (45 L/acre) water when scentless chamomile is actively growing and in the 2 to 4 leaf stage.

Prevail - In registered crops, apply at a rate of 20 acres per case in 10 gallons/acre (45 L/acre) water when scentless chamomile is actively growing and in the 2 to 4 leaf stage.

Refine Extra - Apply 8 g/acre in 10 gallons/acre (45 L/acre) water to actively growing seedlings for suppression. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Tordon 22K - In rangeland and grass pasture, apply 0.445 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Stinging Nettle

2,4-D amine - Apply 0.91 to 1.82 L/acre of formulations containing 500 g/L 2,4-D amine.

Stork's Bill

Ally Toss-N-Go - Apply with 2,4-D or MCPA amine or LV ester in registered crops at registered rates to stork's-bill in the 2 to 4 leaf stage.

Attain - Apply at a rate of 40 acres per case to registered crops when stork's-bill is in the 1 to 8 leaf stage.

Basagran – In registered crops apply 0.91 L/acre at the 2 to 6 leaf stage.

Dichlorprop + 2,4-D - Apply at 0.71 L/acre to registered crops when stork's-bill is in the 2 to 4 leaf stage.

Glyphosate - in Roundup Ready canola, apply 0.5 L/acre in the 0 to 6 leaf stage.

Liberty - in registered crops apply 1.35 L/acre to plants in 1 to 3 leaf stage.

Linuron - Apply with MCPA amine in registered crops at registered rates to stork's-bill in the 2 to 4 leaf stage.

Odyssey/Absolute – in registered crops, apply 17 g/acre of the Odyssey component plus adjuvant.

Prestige - Apply at the rate of one case per 20 acres to registered crops when stork's-bill is in the 1 to 8 leaf stage.

Roundup Transorb - in Roundup Ready canola, apply 0.5 L/acre in the 0 to 6 leaf stage.

Spectrum – In registered crops apply at 20 acres per case from the 2 to 4 leaf stage.

Toadflax (Yellow)

Ally plus 2,4-D - Apply 2 to 3 g/acre Ally plus 0.34 to 0.45 L/acre 2,4-D LV ester or amine (500 g/L formulations) in 10 gallons/acre (45 L/acre) water for toadflax suppression in wheat, barley, and creeping red fescue. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Amitrol 240 - Apply 7.6 to 11.3 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is in the advanced rosette to prebud stage.

Dichlorprop + 2,4-D - Apply 0.71 L/acre in 10 to 18 gallons/acre (45 to 80 L/acre) water in wheat or barley for toadflax suppression. Apply when majority of toadflax is no taller than 6 inches (15 cm). The use of Dichlorprop + 2,4-D for suppression of toadflax in wheat or barley should be part of a long-term planned approach for toadflax control, which includes spring and fall tillage, fall patch spraying, summerfallow or chemical fallow.

Glyphosate - Apply 2.83 to 4.86 L/acre when most plants have reached the early bud stage of growth. Allow 7 more days after application before tillage. A rate of 1.0 L/acre may be used with preharvest applications or when controlling in summerfallow situations.

Refine Extra - In registered crops, apply 8 g/acre in 10 gallons/acre (45 L/acre) water for suppression of toadflax. Apply when toadflax is less than 15 cm (6 inches) in height. Add non-ionic surfactant at 0.2 L per 100 L spray solution.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. WARNING - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Western Snowberry (Buckbrush)

2,4-D amine or LV ester (500 g/L) - Apply 1.82 L/acre 2,4-D amine or LV ester in a minimum of 20 gallons/acre (90 L/acre) water in spring or early summer. Retreatment may be necessary the following year.

Banvel II plus 2,4-D LV ester (500 g/L) - Apply 1.48 L/acre Banvel II tank mixed with 1.82 L/acre 2,4-D LV Ester in 20 gallons/acre (90 L/acre) water in spring or early summer after the leaves are fully expanded.

Escort - Apply 10 g/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water between mid-June and mid-August after the brush has leafed out, but before the leaves turn their fall colours.

White Cockle

2,4-DB - Apply Embutox 625 at 1.1 L/acre or Caliber 400 at 1.7 L/acre or Cobutox 600 at 1.1 L/acre for top growth control to registered crops only.

Mecoprop - Apply 2.2 L/acre in 18 gallons water/acre (*80 L/acre) for top growth control of established plants. Will also control seedlings. Apply to registered crops only.

Wolf Willow (Silverwillow)

Banvel II + 2,4-D amine or LV ester (500 g/L) - In grass pastures with no legumes, apply Banvel II at 2.1 L per 1000 L of water with 2,4-D LV ester or amine at 4.0 L per 1000 L of water to the foliage of actively growing brush in the spring or early summer and wet the foliage until the point of runoff.

Wild Tomato

2,4-D or MCPA amine or ester (500 g/L) - Apply 0.34 to 0.45 L/acre to registered crops up to the 8 leaf stage of wild tomato.

Bromoxynil+MCPA ester - Apply 0.40 L/acre to registered crops from the 1 to 6 leaf stage of wild tomato.

Willow

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Banvel II+ 2,4-D - In grass pasture and rangeland only, apply Banvel II at 1.7 L plus 2,4-D 500 amine at 3.24 L per acre in 20 to 30 gallons (90 to 135 L) of water per acre when leaves are fully expanded.

Glyphosate - As a non-selective spot treatment, apply 1.21 to 2.43 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in the summer through early fall when brush is actively growing.

Remedy EC - In grass pasture and rangeland apply 1.62 to 3.24 L /acre in 45 gallons/acre (200 L/acre) of water in broadcast treatment to fully expanded, actively growing foliage. Single stem foliar and backpack methods are also available - See Remedy section for details.

Soil Residual Herbicides

When applied at recommended rates in a crop, most herbicide residues will disappear within a few weeks after application and impose no restriction on cropping options the next year. However, some herbicide residues do not degrade quickly, and can persist in the soil for months or years following application, thereby restricting the crops that can be grown in rotation. Herbicide residues in the soil are deactivated in various ways including:

- Break down by chemical reactions,
- Break down by soil microbes,
- Escape to the atmosphere as a gas (volatilization),
- Break down by light (photodegradation),
- Leaching,
- Binding to soil particles.

Herbicides often disappear from the environment by more than one of these mechanisms. Many herbicides considered to be non-residual are bound temporarily to soil particles while they are broken down gradually by either soil microbes or chemical reactions. The binding action insures that the herbicide is not available to the crop in quantities that will cause damage.

As a general rule, breakdown processes are favoured by warm, moist soil conditions. During the winter, when the ground is frozen, and in the summer when the soil is dry, herbicide degradation is reduced. The residual activity of certain herbicides is also affected by soil organic matter and soil pH. These soil factors are seldom uniform across a field. Herbicide carryover is aggravated by low levels of organic matter and is more likely to occur on eroded hilltops than in other parts of a field. The risk of herbicide carryover will also be greater in sprayer overlaps which are most common around headlands and slough margins.

Growers should be aware of the residual properties before applying any herbicide if they are to avoid cropping restrictions in following years. Knowledge of the limitations associated with herbicides that leave a soil residue, along with an accurate record of application (i.e. rates, locations) will serve to minimize rotational problems. Each herbicide used in mixes should be considered separately.

Soil tests using chemical extraction cannot always give a good indication of the potential injury risk from herbicide residue because of the influence of organic matter, clay and pH. Because of this, a field bioassay or laboratory bioassay, where plants are grown directly in the treated soil are best for detecting the potential for injury. These tests are not intended to be used to shortcut restrictions on the label, but provide information on rotational crops where none is available.

Injury symptoms from other causes can resemble herbicide carryover injury (i.e. cold weather, flooding, drought, insects, diseases, etc.). Consult with your local agronomist on potential causes before spending money on testing.

Herbicides that leave a soil residue and are of particular concern in Western Canada are found in the following chart.

Recropping Restrictions for Residual Herbicides:

Figures listed are the number of cropping seasons before each crop can be grown ("1" means that the crop can be grown the year following application). Products that have preseeding restriction (in days) are listed in italics. A blank space means that there are no recommendations given on the product label and a field bioassay is recommended by many product manufacturers to determine if these crops are safe to plant. A field bioassay is a strip of a test crop that covers an area of the fieldthat is representative of fieldthe variation and should include an untreated area. Laboratory soil residue bioassay services are also available from Alberta Research Council – Contact Harold Feddema at (780) 632-8238 or Sandi Checkel at (780) 632-8217 for more information.

PRODUCT	Alfalfa	Barley	Canaryseed	Clearfield canola	Non-Clearfield canola	Fababeans	Field corn	Dry beans	Field peas	Flax	Forage grasses	Lentils	Mustard (yellow)	Oats	Potatoes	Rye	Soybeans	Sunflowers	Wheat (durum)	Wheat (spring)	Wheat (winter)
2,4-D*	1	*0d*	1	1	1		1	1	1	1	1	1		1		0d			1	0d	0d
Absolute*		1	1	1						2				1					1	1	
Accent		1					1							1					1	1	
Adrenalin		1		1					1					1					1	1	
Ally (pH less than 7, Brown and Dark Brown soils)		1	4		2					2		3	4	1					1	1	
Ally (pH less than 7, other soils)		1	4		1					1		3	4	1					1	1	
Ally (pH 7 to 7.9, Brown and Dark Brown soils)		1	4		3					3		4	4	2					1	1	
Ally (pH 7 to 7.9, other soils)		1	4		2					3		4	4	1					1	1	
Amitrol		*1d*	1	*1d*	*1d*		*10d**	*10d**	*10d**	1		1	1	1			*6d*	1	*1d*	*1d*	*1d*
Assert (Black and Grey Wooded soils)		1	2	1	1				1	1				2				1	1	1	
Assert (Brown and Dark Brown soils)		1	2	1	2				2	2				2				1	1	1	
Atrazine, Laddock*, Primextra II Magnum, Shotgun						1*	1		1*	1*											
Attain, Trophy	2	1	2	1	1	2	2	2	1	1	1	1	1	1	2	1	2	2	1	1	1
Avadex	0	0	0	0	0	1	1	1	0	0		1	0	2		1	1	1	0	0	0
Banvel II*		1		1*	1*		1	1*						1			1		0*	0*	1
Curtail M, Eclipse, Prevail, FlaxMax, Prestige	2	1	2	1	1	2	1	2	1*	1	1	2	1	1		1	2	2	1	1	1
Edge	0		2	0	0	0		0	0		2	0	0	2			0	0	1*	1*	
Everest, K2 (Brown soils)																			1	1	
Everest, K2 (Dark Brown soils)		1*		1*	1*				1*	1*									1	1	
Everest, K2 (Black soils)		1*		1*	1*			1*	1*	1*									1	1	
Everest, K2 (Grey-Wooded soils)		1*		1*	1*				1*										1	1	
Frontier							0*	0*													1
Frontline, Spectrum (F=Frontline only)		1		1	1				1(F)										1	1	1
Kerb	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2
Lontrel		1		1	1					1	1		1	1		1			1	1	
Muster, Muster Gold	2	1	2		2	2			2	1	2*	2	2	1					1	1	1
Odyssey*	1	1	1	1					1	2		1		1					1	1	
PrePass		0d		1	1				1	1		1	1	0d					0d	0d	0d
Prism		1				1*	1								1		1				0
Pursuit, Pursuit Ultra	1	1		0					0			1							2	1	
Reflex*							1	0									0			1	0
Sencor				2	2	0			0						0*		0*	2			
Sundance - organic matter less than 4%		2		1	2				2	2									1	0	
Sundance - organic matter less than 4% under drought**		3		1	3				3	3									1	0	
Sundance - organic matter greater than 4%		1		1	1				1	1									1	0	
Sundance - organic matter greater than 4% under drought**		2		1	2				2	2									1	0	
Trifluralin	0	1*	2	0	0	0	1*	0	0	1*	2	0	0	2		0	0	0	1*	0*	1*
Ultim		1					1														1
Unity - soil pH less than 6.5*		1	2		1				1	1			2	1					1	0	
Unity - soil pH 6.5 to 7.4*		1	2		2				2	1			2	1					1	0	
Unity - soil pH greater than 7.4*		1	2		2				2	2			3	1					1	0	

* The minimum recropping intervals are listed. These intervals may be longer than those listed depending on the use rates, region, soil types, environment, time of application and crop varietiy. Refer to product page for more information.

** Drought restrictions apply to drought conditions (80% of normal June to September rainfall) for high pH soils (greater than pH 7.5) and severe drought (less than 65% of normal June to Sept. rainfall) for all soils.

0 - May be seeded or reseeded the year of application. No recropping restrictions. 1 - Next cropping season after application. 2 - Two cropping seasons after application. NR - Not recommended. **Note: The recropping intervals listed may not be sufficient to prevent crop injury during periods of below average rainfall.**

Effect of Rainfall on Herbicide Efficacy

Required Interval	Product
15 minutes	Reglone
30 minutes	Horizon
1 hour	Achieve Liquid, Achieve Liquid Gold, Assure II, Bromoxynil, Bromoxynil+MCPA ester, Clethodim, Elite, Everest, Gramoxone, Hoe-Grass, Horizon BTM, Poast Ultra, Puma^{120} Super
2 hours	2,4-D LV Ester, Ally+2,4-D LV Ester, Atrazine (post-emergent applications), Fusion, MCPA Ester, Triumph Plus, Trophy, Unity, Venture
4 hours	Accent, Ally + 2,4-D Amine, 2,4-D Amine, Harmony Total, Liberty, MCPA Amine, Muster, Muster Gold, Prism, Refine Extra, Ultim
6 hours	Assert, Avenge, Curtail M, Everest, Express Pack, FlaxMax, K2, MCPA-K, MCPA Sodium Salt, Prestige, Prevail, Rustler, Sencor, Shotgun, Tordon 22K
8 hours	Basagran, Laddock
No specific recommendation*	Absolute, Adrenalin, Amitrol 240, Attain, Caliber, Cobutox, Compitox, Dicamba, Dicamba+ Mecoprop+MCPA, Dichlorprop-D, DyVel, DyVel DSp, Eclipse, Embutox, Escort, Estaprop, Frontline, Frontline+2,4-D, Glyphosate, Grazon, Linuron, Lontrel, Mecoprop, Odyssey, PrePass, Pursuit, Remedy, Spectrum, Thumper, Tropotox Plus, Turboprop

* The products listed make no specific time recommendation on the label. The required rainfree period could be up to 8 hours. See the product page in the guide or consult the product label.

Note: The term "Rainfastness" refers to the time needed between application and rainfall to avoid significant reduction in efficacy. Rainfall shortly after application of most post-emergent herbicides may reduce weed control. Effect will vary with product, the interval between spraying and rainfall and the intensity and duration of the rainfall. These guidelines are based on label information. Use the longest time interval on the component products when considering tank mixes.

Plant Disease Control

Integrated Plant Disease Management

Cereals

Cereal Leaf Diseases

Cereal leaf diseases continue to hurt growers on two fronts: yield and quality. The following management practices are recommended for effective control of leaf diseases in all cereal crops, especially high-value barley.

Scouting: Scout fields prior to, during, and following flag leaf emergence to check for disease levels. Healthy flag and upper leaves are critical, since more than 50% of grain filling is contributed by upper leaves.

Crop Rotation: Rotate crops (e.g. wheat / oilseed / barley / pulse) to reduce the build-up of disease inoculum in crop residue. If at all possible, do not follow wheat with wheat or barley with barley. When a short rotation is absolutely necessary, seed in the second year a variety that is more resistant to an anticipated disease problem.

Clean Seed: Use certified seed and/or seed that has been tested at an accredited laboratory and is known to have low or no seed-borne disease. Treating with an appropriate fungicide will reduce the possibility of introducing seed-borne inoculum into a field.

Seed Treatments: Seed treatments together with quality seed help emerging crops by controlling most seed-borne disease and allowing the crop to get a good start (quicker, more uniform plant emergence and better seedling vigour). Seed treatments protect young plants against seedling diseases but do not prevent later infection by cereal leaf diseases.

Resistant Varieties: Provincial crop/seed guides provide a comprehensive listing of the performance of adapted varieties, including their resistance status to specific diseases.

Foliar Fungicides: Foliar fungicides applied at the proper time in accordance with manufacturer's recommendations can control cereal leaf diseases and help to attain target yields. The greatest benefit occurs when disease pressure is high or with varieties that have poor resistance.

Spraying Practices: For maximum effectiveness, foliar fungicides should be applied preventively (when infection levels are low). Good spray coverage with minimal drift is essential. Ideally, the best time to spray is when the wind is light; humidity is above 60%; and air temperature is between 10 and 25°C.

Fusarium Head Blight

Fusarium head blight (FHB) of cereals causes a reduction in yield as a result of floret sterility and the loss of light weight kernels during combining. More important is the effect on grain quality and food safety. In Manitoba, FHB currently occurs throughout all crop regions and will damage wheat crops whenever environmental conditions favour the disease. In Saskatchewan, FHB is established in eastern regions but has occurred in other areas throughout the province.

For FHB to occur, three requirements must be met: the disease causing fungus must be present, a susceptible host grown, and environmental conditions favourable for infection and disease development. If any one of these factors is missing disease cannot occur; the severity of disease in the field is a result of the interaction of the above factors.

Field Management to Prevent Recurrence

Weather patterns are by far the greatest factor in the recurrence of FHB. The disease is most likely to develop when the plants are flowering, temperatures range from 15 to 30°C and moisture is continuous for 48 to 60 hours. If conditions remain warm and moist, the pathogen can continue to sporulate and spread to other kernels or heads. Under these optimum conditions, crop management has little impact on FHB outbreaks.

Crop Rotation and Selection

A break of at least one year – preferably two years – is advised between cereal, grass and corn production. Regardless of the rotation, producers should consider planting cereals that are less susceptible to FHB. Results from previous years show that durum wheat is more susceptible than hard red spring wheat varieties. Barley is more tolerant than wheat, and oats are more tolerant than either wheat or barley.

Avoid planting the more susceptible types of wheat in high risk areas such as the southern part of the Red River Valley and southeastern Saskatchewan. Planting two or more varieties of wheat with differing flowering times or varying planting dates will help reduce the risk of infection. Susceptible crops should definitely not be planted on infected corn stubble. Corn trash is slower to decompose than cereal trash, and acts as a source of inoculum for a much longer time period. In fields of wheat on wheat stubble, the incidence was about one and a half times higher than in fields of wheat planted into pulse crop residue.

A forecast risk of FHB occurrence in wheat during the growing season was conducted by Manitoba Agriculture, Food and Rural Initiatives (MAFRI) and the Agrometeorological Centre of Excellence (ACE) with support from Bayer CropScience in 2004. This information appears on the ACE website: www.aceweather.ca under Maps and Interpretations.

Potatoes

One of the major threats to Manitoba and Saskatchewan's potato industry is the fungal disease late blight. At present, there are no fungicides registered for use on potatoes that are capable of eradicating the fungus once it is present. As a result, producers are forced to adopt preventive management to control this disease. One of the main components of this strategy is the application of fungicides at specified spray intervals. This interval varies with the type of fungicide used. Shortening or lengthening of this interval should be based on current weather conditions and the status of the disease in the crop.

In Manitoba and Saskatchewan, potato producers can make use of a weather-based late blight forecasting program. The purpose of this program is to predict when environmental conditions are most conducive to disease development and issue warnings based on those predictions.

Accurate weather monitoring and scouting techniques are very important for achieving the most effective use of fungicides. Combining precise weather monitoring with spray interval scheduling may lower input costs for the farmer and lead to a more productive, higher quality crop. These weather monitoring systems monitor key environmental variables, such as relative humidity, temperature, leaf wetness and precipitation. Risk forecast maps and interpretations appear on the ACE website at: www.aceweather.ca

The following management practices are recommended for effective control of late blight in potato crops.

Scouting: To effectively schedule preventative fungicide applications and eliminate unnecessary fungicide use, local weather forecasts should be used to identify conditions conducive to disease development. Scout fields to identify hot spots and other sources of disease when conditions are favourable.

Crop Varieties and Disease Resistance: Select varieties with resistance to late blight wherever possible. Where practical, the use of short season varieties may help reduce the period of use for fungicides.

Healthy Seed: Obtain seed from sources with effective disease management practices. The use of certified seed is recommended in Manitoba and legislated in Saskatchewan. Grade seed carefully while cutting and discard suspicious looking tubers and seed pieces.

Sanitation and Cull Clean-up: Follow a program of sanitation for storage facilities and equipment to eliminate sources of the disease. Avoid leaving tubers, including debris or slivers from seed cutting, in cull piles for any length of time. Dispose of cull piles in an approved manner so they do not serve as a source of late blight inoculum for future infections. Dispose by burying, using a cover, spreading out on the field over winter, or feeding to livestock.

Cultural Practices and Rotation: Use proper cultural practices including: rotate potato crops with non-late blight hosts; use proper hilling to reduce infection in tubers; increase spacings of plants to reduce canopy density; manage irrigation use to avoid increasing disease risk during prolonged periods of wetness; identify and destroy hot spots of infected fields; control weed hosts and remove and destroy volunteer potatoes.

Canola

For the past 25 years, sclerotinia stem rot has been one of the two prominent diseases affecting canola in Manitoba and Saskatchewan. An important factor of disease development is environmental conditions. The disease is much more widespread and severe during wet years.

Fungicide applications are an important element in controlling the development and spread of sclerotinia. Fungicide spray decisions are based on soil moisture, weather conditions, crop stage and density, and the history of disease in your area.

The sclerotinia resting bodies require moist soil conditions for up to 10 days for germination to occur and the spore-bearing structures (apothecia) to form. Usually these conditions do not occur until the crop canopy closes. The spores released from the apothecia utilize the canola petals as a food source and fall into the canola canopy where they infect individual plants. Lesions form up and down the stem, wilting leaves and eventually killing the plant. Fungicide should be applied between the 20 to 50% flower stages to protect the petals from being colonized by the spores.

Pulse Crops

There are a variety of pulse crops produced in Manitoba and Saskatchewan including field pea, field bean, lentil, chickpea and soybean. Pulse crops are adapted to different regions and will require unique agronomic and disease management practices. Some diseases will attack all pulse crops, e.g. sclerotinia (white mould) and seedling/root rots caused by *Pythium, Rhizoctonia, Fusarium, Botrytis* species. Some diseases may occur on more than one type of pulse crop, but the pathogen species infecting each is specific to that crop. This is the case for the ascochyta blights, powdery mildews and anthracnose. It is important to source a variety of information on pulse disease control from grower organizations such as the Saskatchewan Pulse Growers (www.saskpulse.com), provincial specialists, and field diagnostic guides. Most foliar diseases are favoured by warm, moist conditions and lush crop canopies, but root rots and powdery mildew can be present in dry years as well. In general, pulse disease management will need to include the following practices;

Use of clean seed and seed treatments: Plant certified seed or seed that has been tested at an accredited lab and known to have zero or acceptable levels of seed-borne disease and high germination. Seed treatments will help protect seed and seedlings from low levels of seedborne and soilborne pathogens.

Crop Rotation: It is important to keep at least three years between the same type of pulse crop to allow for the breakdown of crop residue on which disease pathogens survive. Since there are diseases that affect more than one type of pulse crop, it is still important to maintain at least two years even between different pulse crops.

Crop Varieties with Disease Resistance: Refer to provincial seed guides for varieties adapted to your region. Choose varieties with disease resistance, when available.

Scouting and Foliar Fungicide Application: Begin crop scouting at the vegetative stages for aggressive diseases such as ascochyta blight in chickpea. Scout for other foliar diseases at early bloom, e.g. ascochyta blight and anthracnose in lentil. It is too late to apply fungicide to control sclerotinia (white mould) once symptoms are observed, so forecasting to determine risk is necessary.

There are now a number of foliar fungicides registered for pulse crops. Use foliar fungicides only when disease risk and potential loss are significant. Rotate fungicides from different fungicide groups to prevent the development of resistant pathogen populations.

Resistance Management

Any fungal pathogen population may contain strains naturally resistant to a fungicide and other fungicides within the same Group. A gradual or total loss of disease control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of a fungicide, and others within the same Group, with different Groups that control the same pathogens.
- Avoid application of more than the maximum number listed in the label. Also avoid consecutive sprays of a fungicide, or other fungicides in the same Group, in a season.
- Fungicide use should be based on an integrated pest management (IPM) program that includes scouting and accurate recording related to pesticide use and crop rotation. An IPM program also considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for signs of resistance development. If disease continues to progress after treatment with a product, do not increase the use rate. Discontinue use of the product and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

Table 1: Fungicide Groups Based on Mode of Action

Group	Chemical Group/ Mode of Action	Foliar Fungicide Common Name	Seed Treatment Common Name
1	MBC/ inhibition of tubulin formation	Senator 70WP	Crown*, DCT*, Mertect SC, Senator PSPT
2	Dicarboximides/ affect cell division, DNA/RNA synthesis and metabolism	Ronilan EG, Rovral Flo	Foundation Lite *
3	DMI/ inhibition of sterol biosynthesis	Folicur, Stratego 250EC* Tilt 250E	Baytan 30, Charter, Dividend XL RTA*, Helix*, Helix XTra*, Raxil FL, Raxil 250 FL
4	PA/ affect RNA synthesis	Ridomil Gold/Bravo*	Allegiance FL, Apron FL, Apron Maxx RTA*, Dividend XL RTA*, Helix*, Helix XTra*, Poncho 600 FS*, Prosper FL*
7	Carboximides/ affect mitochondrial	Lance	Crown*, Gaucho CS FL*, Gaucho 480FL*, Poncho 600 FS*, Prosper FL*, Vitaflo 280*, Vitavax rs Fungicide*, Vitavax Single Solution
11	QoI/ inhibition of mitochondrial respiration	Headline EC, Quadris, Reason 500SC, Stratego 250EC*, Tanos 50 DF*	None
12	PP/ affect proteins involved in membrane permeability	None	Apron Maxx RTA*, Helix*, Helix XTra*, Maxim 480FS, Maxim PSP
15	Cinnamic acids/ affect cell wall synthesis	Acrobat MZ*	None
22	Benzamide/ inhibition of tubulin formation	Gavel 75 DF*	None
28	Carbamates/ affect cell membrane permeability	Tattoo C*	None
29	2,6-Dinitroanilines/ uncouples oxidative phosphorylation	Allegro 500F	None
M	Multi-site activity	Acrobat MZ*, Bravo 500, Copper-based Fungicides, Gavel 75 DF*, Kumulus DF, Mancozeb-based Fungicides, Polyram DF, Ridomil Gold/ Bravo*, Tattoo C*	Agrox B-2, Agrox CD, Captan Flowable, DB Red L, DCT*, Dithane F-45, Foundation Lite*, Gaucho CS FL*, Gaucho 480FL*, Mancozeb-based Fungicides, Polyram 16D, Poncho 600 FS*, Prosper FL*, Thiram 75WP, Vitaflo 280*, Vitavax rs Fungicide*
U	Cyanoacetamideoximes/ unknown	Curzate 60 DF, Tanos 50 DF*	None

* Products contain more than one active ingredient and appear in more than one group.

MBC = Methyl Benzimidazole Carbamates
DMI = Demethylation Inhibitors (SBI Class I)
PA = Phenyl Amides
QoI = Quinone outside Inhibitors
PP = Phenyl Pyrroles
SBI = Sterol Biosynthesis Inhibitors

Foliar Fungicide Directory

Table 2. Foliar Fungicides for Disease Control in Potatoes

FUNGICIDES	Page	Botrytis Gray Mould	Early Blight, Late Blight
Acrobat MZ	228		•
Allegro 500 DF	229		②
Bravo 500	230	•	•
Copper 53W	231		•
Copper Spray	231		•
Curzate 60 DF	233		②
Dithane DG Rainshield NT	241		•
Gavel 75 DF	234		•
Headline EC	235		•
Kocide 101	231		•
Kocide DF	231		•
Lance	239		①
Manzate DF	241		•
Parasol Flowable	231		•
Parasol WP	231		•
Penncozeb 75 DF	241		•
Polyram DF	242		•
Quadris	243		•
Reason 500SC	244		•
Ridomil Gold/Bravo	245	•	•
Tanos 50 DF	250		•
Tattoo C	251		②

① early blight only
② late blight only
Note: Before using any pesticide on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

Table 3. Foliar Fungicides for Disease Control in Cereals

FUNGICIDES	Page	Wheat	Barley	Oats	Rye	Crown Rust (Oats)	Leaf Rust	Stem Rust	Stripe Rust	Septoria (Wheat, Barley, Oats,)	Tan Spot (Wheat)	Net Blotch, Scald (Barley)	Powdery Mildew	Spot Blotch	Fusarium Head Blight
Bravo 500	230	X								①	•				④
Dithane DG Rainshield NT	241	X					•			①	•				
Headline EC	235	X	X		X		③		②	①	•	•	③	②	
Manzate DF	241	X					•			①	•				
Penncozeb 75 DF	241	X					•			①	•				
Stratego 250EC	249	X					①	①	①	①	•		①		
Tilt 250E	252	X	X	X		•	②	②	①	•	•	•	②		

① wheat only
② wheat **and** barley
③ wheat **and** rye
④ suppression only

Table 4. Foliar Fungicides for Oilseed and Pulse Crops

FUNGICIDES	Page	Canola	Beans	Chickpeas	Lentils	Peas	Alternaria Blackspot (Canola)	Anthracnose (Beans, Lentils)	Ascochyta Blight (Chickpeas, Lentils, Peas)	Bacterial Blight (Beans)	Blackleg (Canola)	Botrytis Gray Mould (Beans,Chickpeas, Lentils)	Downy Mildew, Anthracnose (Beans)	Mycosphaerella Blight (Peas)	Powdery Mildew (Peas)	Rust (Beans)	Sclerotinia/White Mold (Canola, Beans)
Bravo 500	230			X	X	X		①	•					•			
Copper 53W	231		X							•			•				
Dithane DG Rainshield NT	241				X			①	①								
Headline EC	235		X	X	X	X		•	•					•	•	•	
Kocide 101	231		X							•							
Kumulus DF	238					X									•		
Lance	239	X	X	X	X		⑥		④			⑤					•
Parasol WP	231		X							•							
Quadris	243	X	X	X	X	X	•	•	•		•			•		•	•
Ronilan EG	246	X	X									②					•
Rovral Flo	247	X					•										③
Senator 70 WP	248		X														②
Tilt 250E	252	X	X								•					•	

① lentils only
② beans only
③ canola only
④ except peas
⑤ except beans
⑥ suppression only

Table 5. Foliar Fungicides for Disease Control in Specialty Crops

FUNGICIDES	Page	Alfalfa grown for seed	Canaryseed	Bluegrass, Fescues and Ryegrass grown for seed	Leaf and Stem Spot Diseases	Leaf Rust, Stem Rust (Grass Seed)	Powdery Mildew (Grass Seed)	Septoria (Canaryseed)
Dithane DG Rainshield NT	241	X			•			
Headline EC	235			X		•	①	
Tilt 250E	252		X					①

① suppression only

Acrobat MZ

Fungicide Group – 15, M
(Refer to page 225)

Company:

BASF Canada PCP# 24546

Formulation:

9% dimethomorph and 60% mancozeb formulated as a wettable powder.

Container size - 10 kg.

Crops:

Potatoes.

Diseases Controlled:

Early and late blight. Reduction of tuber blight.

Crop Stage:

Make the first application when the disease threatens or when the first visible signs of disease occur in the field or nearby. Apply every 5 to 7 days under high disease pressure or every 7 to 10 days under low disease pressure. Do not apply more than 3 times per season. It is recommended to apply this product alternately with a fungicide having a different mode of action. Under high level of late blight infection, apply after top kill to control tuber blight.

How it Works:

Acrobat MZ has protectant, systemic and antisporulant activity. The active ingredient mancozeb is a dithiocarbamate fungicide with contact activity. The active ingredient dimethomorph penetrates into the leaf tissue and moves within the leaf.

Cost (2004 suggested retail price):

$28.50/acre.

Rate:

Apply at 1.0 kg per acre.

Water Volume:

Use sufficient water to obtain adequate spray coverage.

Ground - 18 gallons/acre (80 L/acre).

Aerial - minimum 5 gallons/acre (20 L/acre).

Tank Mixes:

None registered.

Restrictions:

Rainfall: Do not apply when rain is expected within 2 - 3 hours. Apply to dry foliage.

Preharvest Interval: 14 days.

Recropping: Do not replant in treated area within 120 days of last application.

Environment: Do not apply to terrain where there is a potential for surface runoff to enter aquatic systems. This product is highly toxic to aquatic organisms. Do not apply within 100 m of streams, ponds, rivers and lakes when applying by air and within 50 m when applying by ground.

Storage: Store under cool, dry conditions in secure, well ventilated buildings away from food or feed.

Resistance: Plant disease fungi can develop resistance when exposed to one type of product or even products of similar chemistry. Use cultural practices and fungicide rotation as well as early preventive fungicide applications. Consult the government Potato Specialist or Plant Disease Specialist for disease outbreak forecast and recommendations.

Hazard Rating:

Caution: Potential skin sensitizer.

Allegro 500F

Fungicide Group – 29
(Refer to page 225)

Company:

Syngenta – PCP# 27517

Formulation:

40% fluazinam. Container size – 18.9 L.

Crops:

Potatoes

Diseases Controlled:

Late blight *(Phytophthora infestans)*

Crop Stage:

Begin applications when the plants are 15-20 cm tall or when conditions favour disease development. Repeat application at 7-10 day intervals. Do not make more than 10 applications in a season.

How it Works:

Fluazinam is a pyridinamine fungicide with protective (contact) activity.

Cost (2004 suggested retail price):

$6.10/acre ($15.24/ha)

Rate:

0.16 L/acre (0.4 L/ha)

Water Volume:

80-240 L/acre (200-600 L/ha). Spray volumes vary with amount of plant growth; apply in sufficient water to obtain adequate coverage of foliage.

Tank Mixes:

None registered.

Restrictions:

Resistance: This product is a Group 29 fungicide and is considered to have low risk of resistance development. It is recommended to not rely exclusively on the same product for pest control. Prolong the effectiveness of this product by not making more than 3 consecutive applications before alternating to a fungicide with a different mode of action. Use as part of an integrated pest management program.

Preharvest interval: 14 days

Re-cropping: Can be replanted with potatoes as soon as practical after the last application, 30 days for other root crops and leafy vegetates, and 70 days for all other crops. Fluazinam will carry over, do not use in areas treated with this product during the previous season.

Environment: Do not apply during periods of dead calm or when winds are gusty. Do not contaminate aquatic habitats when cleaning and rinsing spray equipment or containers. Do not overspray non-target terrestrial or aquatic habitats.

Precautions:

Keep out of reach of children. Causes skin irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes. Wear coveralls over long sleeved shirt, long pants, chemical resistant gloves, socks and footwear. Avoid breathing spray mist.

Hazard Rating:

Caution Poison

Warning - skin irritant.

Potential skin sensitizer.

Bravo 500

Fungicide Group – M
(Refer to page 225)

Company:

Syngenta PCP# 15723

Formulation:

500 g/L chlorothalonil formulated as a suspension.
Container sizes - 2 x 10 L case, 200 L drum.

Crops:

Wheat (all wheat including durum & winter wheat), potatoes, lentils, chickpeas and field peas.

Diseases Controlled:

Wheat - Tan Spot, Septoria Glume Blotch, Septoria Leaf Spot. Suppression of Fusarium Head Blight (Scab).

Potatoes - Early and late blight, Botrytis vine rot.

Lentils - Anthracnose and Ascochyta blight.

Chickpeas - Ascochyta blight

Peas - Mycosphaerella blight

Crop Stage:

Wheat - For control of Tan Spot and Septoria, begin application at flag leaf emergence and repeat 10 to 14 days later when ears are visible. A third application when ears are fully emerged may be necessary, if conditions favour disease spread. For suppression of Fusarium Head Blight apply Bravo at early flowering. For best results, this application must be made before flowering has started in the majority of tillers and before the beginning of weather favouring disease. Do not make more than 3 applications per season. Rates and number of applications will depend upon disease severity and weather conditions.

Potatoes - Begin application when plants are 6 to 8 inches (15 to 20 cm) high or when disease threatens. Repeat applications at 7 to 10-day intervals or as necessary to maintain disease control. Under severe disease conditions use the higher rates at 7-day intervals.

Lentils - One application, must occur at early flowering. Two applications - first application before flowering when bud formation is evident. Second application must occur at early to mid-flowering, 10 to 14 days after the first application, but before rows close in to form a dense canopy. Rates and number of applications will depend upon disease and weather conditions.

Chickpeas - Make first application at early flowering and remaining applications at 10-day intervals. Do not make more than 3 applications per season. Rate and number of applications will depend upon disease severity and weather conditions.

Peas - Begin application at early flowering and repeat 10 days later at early pod set or mid-flowering if necessary. Make a third application 10 to 14 days after the second application at pod fill or later flowering should conditions remain favourable for disease. Do not make more than 3 applications per season. Rates and the number of applications will depend upon disease severity and weather conditions.

How it Works:

The active ingredient chlorothalonil is a chloronitrile fungicide with contact activity.

Cost (2004 suggested retail prices):

$8.77 to $14.61/acre for wheat.
$7.01 to $14.61/acre for potatoes.
$11.69 to $23.38/acre for lentils.
$11.69 to $23.38/acre for chickpeas.
$11.69 to $17.53/acre for peas.

Rate:

Wheat - 0.6 L to 1 L/acre for the control of Tan Spot and Septoria.

0.8 L to 1 L/acre for the suppression of Fusarium Head Blight.

Potatoes - 0.48 to 1 L/acre for control of late blight. 0.65 to 1 L/acre for control of early blight and botrytis stem rot.

Lentils - 0.8 to 1.6 L/acre.

Chickpeas - 1.2 to 1.6 L/acre for first application. 0.8 to 1.2 L/acre for subsequent applications.

Peas - 0.8 L to 1.2 L/acre for the control of Mycosphaerella blight.

Water Volume:

Volume will vary with amount of plant growth. Use sufficient water to obtain adequate spray coverage. Spray volume will usually range from 20 to 140 gallons/acre (90 to 640 L/acre) for dilute sprays and 4.4 to 8.8 gallons/acre (20 to 40 L/acre) for concentrate sprays.

Chickpeas - 20 gallons/acre (90 L/acre). Ground application only.

Tank Mixes:

Fungicides: On potatoes, Bravo may be applied with Quadris for the control of early blight. Do not apply sequential applications of this tank mix and do not exceed 3 tank mix applications per season. Do not apply to potatoes later than 2 days before harvest.

Do not combine with pesticides, surfactants or fertilizers unless prior use has shown the combination physically compatible and non-injurious under your conditions of use.

Herbicides: On lentils, do not apply in combination with POAST herbicide and MERGE surfactant or within 48 hours of the application of POAST and MERGE.

Fertilizers: None registered.

Insecticides: None registered.

Restrictions:

Grazing: Do not graze treated areas. Do not feed straw from treated crop to livestock.

Preharvest Interval: Potato - 1 day.
Lentils and chickpeas - 48 days.
Wheat - 30 days.
Peas - 32 days.

Recropping: None.

Application: On lentils do not make more than 2 applications in the same season. On chickpeas, wheat and peas, do not make more than 3 applications per season. Do not apply by air for chickpeas.

Storage: Store in a cool, dry, ventilated place. Protect from excessive heat.

Environment: Do not apply if weather conditions favour drift from area being treated. Do not contaminate lakes, streams or ponds. Allow a buffer zone of 15 m between area being treated and aquatic systems for ground applications and a buffer zone of 100 m for aerial application.

Re-entry: Do not re-enter treated area within 48 hours. If required, and at least 4 hours have passed since application, individuals may re-enter treated area for short-term tasks not involving hand labour. Long pants, long-sleeved shirt, and chemical resistant gloves must be worn.

Hazard Rating:

Warning, causes severe eye damage.

Copper-based Fungicides

Copper 53W/Copper Spray/Kocide 101/Kocide DF/ Parasol WP/Parasol Flowable

Fungicide Group – M
(Refer to page 225)

Company:

United Agri Products, (Copper 53W – PCP# 09934, Copper Spray – PCP# 19146)

E.I. du Pont Canada Company (Kocide 101 – PCP# 14417, Kocide DF – PCP# 24538)

Nufarm (Parasol WP – PCP# 24671, Parasol Flowable PCP# 25901)

Formulation:

Copper 53 W - 53% tribasic copper sulphate formulated as a wettable powder.
Container size - 10 kg.

Copper Spray - 50% copper oxychloride formulated as a wettable powder.
Container size - 10 x 2 kg case.

Kocide 101 - 50% copper as copper hydroxide formulated as a wettable powder.
Container size - 5 kg.

Kocide DF - 40% copper as copper hydroxide formulated as a dry flowable.
Container size - 10 kg.

Parasol WP - 50% copper equivalent as copper hydroxide formulated as a wettable powder.
Container size - 10 kg.

Parasol Flowable - 24.4% copper equivalent as copper hydroxide formulated as a flowable.
Container size - 10L.

Crops:

Potatoes, beans.

Diseases Controlled:

PRODUCT	POTATO	BEAN
Copper 53W	Early and late blight	Anthracnose Downy Mildew Bacterial Blight
Copper Spray	Early and late blight	
Kocide 101	Early and late blight	Bacterial blight (bacterial and halo)
Kocide DF	Early and late blight	
Parasol WP	Early and late blight	Bacterial blight (bacterial and halo)
Parasol Flowable	Early and late blight	

Crop Stage:

Potatoes - (Copper 53W, Copper Spray) - Apply when plants are 5 to 7 inches (12 to 18 cm) tall. Repeat at 7 to 10 day intervals.

(Kocide 101, Kocide DF, Parasol WP, Parasol Flowable) - Apply when plants are 6 inches (15 cm) tall. Apply combined with 0.7 to 0.9 kg of a mancozeb (80%) product per acre, at 7 to 10 day intervals. Kocide and Parasol may be applied with a desiccant at vine kill or alone after vine kill, prior to harvest, to reduce the risk of late blight tuber infection. See table for rates.

Beans - First application when plants are 6 inches (15 cm) tall, as a protectant. Repeat every 7 to 14 days depending on local conditions.

How it Works:

The active ingredients tribasic copper sulphate, copper oxychloride and copper hydroxide are inorganic fungicides with contact activity.

Cost (2004 suggested retail prices):

Copper 53W - $12.20/kg
Copper Spray - $12.41/kg
Kocide 101 - $10.86/kg (2003 suggested retail price)
Kocide DF - $10.86/kg (2003 suggested retail price)
Parasol WP - $10.00/kg (2003 suggested retail price)
Parasol Flowable - $14.90/L

Rates:

Product	Potato (kg/acre)	Acres treated by 1 container	Beans (kg/acre)	Acres treated by 1 container
Copper 53W	2.2	4.5	2.2	4.5
Copper Spray	1.6	6.25		
Kocide 101	0.44 - 0.9 1.38 (vine kill)	11.4 - 5.5 3.7 (vine kill)	0.9 - 1.3	5.5 - 3.8
Kocide DF	0.44 - 0.69 1.38 (vine kill)	23 - 14.5 7.3 (vine kill)		
Parasol WP	0.44 - 1.0 1.36 (vine kill)	23 - 10 7.3 (vine kill)	0.9 - 1.3	11 - 7.7
Parasol Flowable	0.32 - 0.72 L/ acre 1 L/ acre (vine kill)	31.2 - 13.9 10 (vine kill)		

Water Volumes:

Kocide 101, DF - Enough to ensure thorough coverage. Check label.

Copper 53W and Copper Spray - 88 gallons/acre (400 L/acre).

Parasol WP, Parasol Flowable - Enough to ensure thorough coverage.

Tank Mixes:

None registered.

Restrictions:

Storage: Store in cool, dry, ventilated area, away from feed or food. Keep away from heat, fire and sparks.

Preharvest Interval: 1 day.

Environment: Do not apply or allow to drift onto streams or any body of water.

Hazard Rating:

◈ Warning Poison (Copper 53W, Copper Spray)

▽ Caution Poison (Kocide 101, Parasol Flowable, Parasol WP)

⯃ Danger Poison (Kocide DF)

Curzate 60 DF

Fungicide Group – U

(Refer to page 225)

Company

E.I du Pont Canada Company PCP# 26284

Formulation:

60% cymoxanil formulated as a dry flowable.

Container size - 1.8 kg

Crops:

Potatoes

Diseases Controlled:

Late blight *(Phytophthora infestans)*

Crop Stage:

Use Curzate 60 DF only in a tankmix with Manzate DF. Do not use Curzate 60 DF alone.

Initial applications should start when local conditions indicate that late blight is imminent. Make additional applications at 5 to 7 day intervals. Apply no more than 7 applications per crop.

How it Works:

The active ingredient cymoxanil is a highly active, locally systemic fungicide. It works at several levels of preventative, curative and inhibitive (against sporulation) activity.

The active ingredient mancozeb in Manzate 200 DF is a dithiocarbamate fungicide with contact activity.

Cost (2004 suggested retail price):

$11.65/acre.

Rate:

Apply Curzate 60 DF at 0.09 kg/acre
Plus
Manzate DF at 0.65 kg/acre

Water Volume:

Utilize sufficient water to obtain thorough coverage: 20 to 100 gallons/acre (80 to 400 L/acre).

Tank Mixes:

None registered.

Restrictions:

Rainfall: Curzate is rainfast within 2 hours after application.

Re-entry: Do not re-enter treated area within 24 hours.

Preharvest Interval: 8 days

Application: Do not apply by air.

Storage: Store product in original container in a secure, dry area away from food or feed. Protect against humid air and water. Not for use or storage in or around the home. Keep container tightly closed.

Environment: A buffer zone of 50m is required between the down-wind edge of the boom and sensitive aquatic habitats such as ponds, lakes, rivers, streams, and wetlands. Do not contaminate these habitats when cleaning and rinsing equipment or containers. Do not clean sprayer near well or water source or near desirable vegetation.

Hazard Rating:

☠ Danger Poison

Gavel 75 DF

Fungicide Group – 22, M
(Refer to page 225)

Company:

Dow AgroSciences PCP# 26842

Formulation:

66.7% mancozeb and 8.3% zoxamide formulated as a dry flowable.

Crops:

Potatoes

Diseases Controlled:

Early and late blight

Crop Stage:

Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. Begin applications at the first sign of disease or when blight is reported in the area. Apply at 0.90 kg/acre every 7 days under high disease pressure when either disease is present and environmental conditions favour continued disease development. Gavel 75 DF can be applied at 0.68 kg/acre every 7 days under low disease pressure and environmental conditions unfavorable for disease development. Do not apply more than 6 applications per season.

How it Works:

The active ingredient zoxamide is a benzamide fungicide with contact activity.

The mancozeb component is a dithiocarbamate fungicide with contact activity.

Cost (2004 suggested retail):

$12.82 to $16.97/acre.

Rate:

Apply at 0.68 to 0.90 kg/acre

Nozzles:

Ground – generally hollow cone, disc – D5 to D7. Consult spray nozzle accessory catalogues for specific information on proper equipment calibration.

Aerial – hollow cone brass nozzles with a D-series orifice disc and core (whirlplate) are recommended. Nozzles should point straight down or slightly backward.

Water Volume:

Thorough, uniform coverage is essential for good disease control.

Ground – 20 gal/acre (90 L/acre).

Aerial – 4 to 8 gal/acre (18 to 36 L/acre). Use 36 liters of water under high disease pressure to provide better crop coverage.

Effects of Weather:

Must be dry on plant leaf prior to rainfall.

Tank Mixes:

None registered.

Restrictions:

Preharvest Interval: 3 days.

Recropping: A 30 day plant back interval (PBI) is required for leafy vegetables and root and tuber vegetables. For all other crops not included on the label, the PBI should be 140 days.

Storage: Do not allow product to freeze. Keep away from fire and sparks. Store in a cool, dry, well ventilated place away from feed or food.

Environment:
Ground application – a buffer zone of 25m for application by ground sprayer should be established between the last spray swath and the edge of aquatic systems. A buffer zone of 5m for application by ground sprayer should be established between the last spray swath and the edge of terrestrial habitats such as hedgerows, windbreaks, woodlots, vegetative strips and other vegetation.

Aerial application – a buffer zone of 20m is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats.

Re-entry: do not re-enter treated areas within 48 hours of application.

Hazard Rating:

Caution – causes moderate eye irritation. This product is a dermal sensitizer. Avoid contact with skin, eyes or clothing.

Headline EC

Fungicide Group – 11

(Refer to page 225)

Company:

BASF Canada PCP# 27322

Formulation:

250 g/L of pyraclostrobin formulated as an emulsifiable concentrate

Container size: 6.5 L jug; 104 L tote

Crops and Staging:

Wheat, barley, rye: For control of leaf diseases apply single application immediately after flag leaf emergence; use higher rate to obtain extended protection; apply second application 10 to 14 days later if disease persists or weather conditions are favourable for disease development.

Chickpea, lentil, dry field pea, dry field bean, faba bean: Apply at the beginning of flowering or at the onset of symptoms for the more aggressive diseases (anthracnose in lentil, ascochyta blight in chickpea); apply a second application 10 to 14 days later if disease persists or weather conditions are favourable for disease development.

Potatoes: Apply prior to row closure when conditions become favourable for the development of disease (whichever comes first); use higher rates and shorter spray intervals under heavy disease pressure; do not make more than 6 applications of Headline per season.

Grasses grown for seed (bluegrass, fescue, ryegrass): Apply prior to disease development; apply second application 14 to 21 days later if disease conditions persist.

How it Works:

The active ingredient pyraclostrobin is a member of the strobilurin class of chemistry used as a broad spectrum preventive and curative fungicide.

Cost (2004 suggested retail price):

$16.57/acre at the 0.16 L/acre rate

Crops, Diseases Controlled, Staging and Rates:

Cereals (Ground and aerial application)

Crop	Diseases controlled:	Rates (L / acre):
Wheat	Tan spot *(Pyrenophora tritici-repentis)* Septoria leaf spot *(Septoria tritici; S. nodorum)* Leaf rust *(Puccinia recondita)*	0.12 – 0.24
	Powdery mildew *(Erysiphe graminis* f. sp. *tritici)* Spot blotch *(Cochliobolus sativus)* Stripe rust *(Puccinia striiformis)*	0.16 – 0.24
Barley	Net blotch *(Pyrenophora teres)*	0.12 – 0.24
	Scald *(Rhynchosporium secalis)* Spot blotch *(Cochliobolus sativus)* Stripe rust *(Puccinia striiformis)*	0.16 – 0.24
Rye	Leaf rust *(Puccinia recondita)*	0.12 – 0.24
	Powdery mildew *(Erysiphe graminis)*	0.16 – 0.24

Pulses (Ground and aerial application)

Crop	Diseases controlled:	Rates (L / acre):
Chickpea	Ascochyta blight *(Ascochyta* spp.*)*	0.16 – 0.24
Lentil	Anthracnose *(Colletotrichum truncatum)* Ascochyta blight *(Ascochyta* spp.*)*	0.16
Field pea	Mycosphaerella blight *(Mycosphaerella* spp.) Powdery mildew *(Erysiphe* spp.)	0.16
Dry beans *(Phaseolus* spp.*)*	Anthracnose *(Colletotrichum* spp.*)* Powdery mildew *(Erysiphe* spp.*)* Rust *(Uromyces* spp.)	0.16
Faba bean	Powdery mildew *(Erysiphe* spp.*)* Rust *(Uromyces* spp.)	0.16

Grasses grown for seed (Ground application only)

Crop	Diseases controlled:	Rates (L / acre):
Bluegrass; fescue; ryegrass	Leaf and stem rust *(Puccinia recondita; P. graminis)* Powdery mildew – suppression *(Erysiphe graminis)*	0.16 – 0.27

Potato (Ground application only)

Crop	Diseases controlled:	Spray interval:	Rates (L / acre):
Potato	Early blight *(Alternaria solani)* Late blight *(Phytophthora infestans)*	7 to 14 days 5 to 7 days	0.18 – 0.27 0.18 – 0.36

Water Volume:

Ground: Use a minimum water volume of 40 L/acre (9 g/acre) on cereals, pulses and grasses and 80 L/acre (18 g/acre) on potatoes. Ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 20 L/acre (4.5 g/acre). Ensure thorough coverage of foliage. Use the highest rate recommended for the disease being controlled.

Tank Mixes:

None registered.

Restrictions:

Resistance: Like any fungicide with a single mode of action, the quinone outside inhibitor fungicides, while highly effective, may result in resistant strains of fungi where they have been overused or misused.

When applying multiple applications in one season, pay close attention to the labelled recommendations for resistance management such as using labelled rates, rotating to fungicides outside group 11 and not exceeding the maximum number of applications per season.

Preharvest interval: Cereals–apply no later than the end of flowering. Pulses–30 days. Grasses–14 days. Potatoes–3 days.

Recropping: Crops listed on label may be planted immediately following last application. Wait 14 days before planting all other crops.

Application: Do not apply during periods of dead calm or when winds are gusty.

Environment: Avoid overspray or drift to sensitive habitats. Maintain specified buffer zones.

Storage: Store in a cool, dry, locked, well-ventilated area. Do not freeze.

Hazard Rating:

☠ Danger Poison

Danger - Skin and eye irritant

Crops, Diseases Controlled, Staging and Rates:

Cereals (Ground and aerial application)

Crop	Diseases controlled:	Rates (L / acre):
Wheat	Tan spot *(Pyrenophora tritici-repentis)* Septoria leaf spot *(Septoria tritici; S. nodorum)* Leaf rust *(Puccinia recondita)*	0.12 – 0.24
	Powdery mildew *(Erysiphe graminis* f. sp. *tritici)* Spot blotch *(Cochliobolus sativus)* Stripe rust *(Puccinia striiformis)*	0.16 – 0.24
Barley	Net blotch *(Pyrenophora teres)*	0.12 – 0.24
	Scald *(Rhynchosporium secalis)* Spot blotch *(Cochliobolus sativus)* Stripe rust *(Puccinia striiformis)*	0.16 – 0.24
Rye	Leaf rust *(Puccinia recondita)*	0.12 – 0.24
	Powdery mildew *(Erysiphe graminis)*	0.16 – 0.24

Pulses (Ground and aerial application)

Crop	Diseases controlled:	Rates (L / acre):
Chickpea	Ascochyta blight *(Ascochyta* spp.*)*	0.16 – 0.24
Lentil	Anthracnose *(Colletotrichum truncatum)* Ascochyta blight *(Ascochyta* spp.*)*	0.16
Field pea	Mycosphaerella blight *(Mycosphaerella* spp.*)* Powdery mildew *(Erysiphe* spp.*)*	0.16
Dry beans *(Phaseolus* spp.*)*	Anthracnose *(Colletotrichum* spp.*)* Powdery mildew *(Erysiphe* spp.*)* Rust *(Uromyces* spp.*)*	0.16
Faba bean	Powdery mildew *(Erysiphe* spp.*)* Rust *(Uromyces* spp.*)*	0.16

Grasses grown for seed (Ground application only)

Crop	Diseases controlled:	Rates (L / acre):
Bluegrass; fescue; ryegrass	Leaf and stem rust *(Puccinia recondita; P. graminis)* Powdery mildew – suppression *(Erysiphe graminis)*	0.16 – 0.27

Potato (Ground application only)

Crop	Diseases controlled:	Spray interval:	Rates (L / acre):
Potato	Early blight *(Alternaria solani)* Late blight *(Phytophthora infestans)*	7 to 14 days 5 to 7 days	0.18 – 0.27 0.18 – 0.36

Water Volume:

Ground: Use a minimum water volume of 40 L/acre (9 g/acre) on cereals, pulses and grasses and 80 L/acre (18 g/acre) on potatoes. Ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 20 L/acre (4.5 g/acre). Ensure thorough coverage of foliage. Use the highest rate recommended for the disease being controlled.

Tank Mixes:

None registered.

Restrictions:

Resistance: Like any fungicide with a single mode of action, the quinone outside inhibitor fungicides, while highly effective, may result in resistant strains of fungi where they have been overused or misused.

When applying multiple applications in one season, pay close attention to the labelled recommendations for resistance management such as using labelled rates, rotating to fungicides outside group 11 and not exceeding the maximum number of applications per season.

Preharvest interval: Cereals–apply no later than the end of flowering. Pulses–30 days. Grasses–14 days. Potatoes–3 days.

Recropping: Crops listed on label may be planted immediately following last application. Wait 14 days before planting all other crops.

Application: Do not apply during periods of dead calm or when winds are gusty.

Environment: Avoid overspray or drift to sensitive habitats. Maintain specified buffer zones.

Storage: Store in a cool, dry, locked, well-ventilated area. Do not freeze.

Hazard Rating:

☠ Danger Poison

Danger - Skin and eye irritant

Kumulus DF

Fungicide Group – M
(Refer to page 225)

Company:

BASF Canada PCP# 18836

Formulation:

80% sulphur formulated as a water dispersible granular. Container size - 25 kg bag.

Crop:

Field peas.

Disease Controlled:

Powdery mildew.

Crop Stage:

Spray at first appearance of disease and repeat at 7 to 10 day intervals as necessary.

How it Works:

The active ingredient sulphur is an inorganic fungicide with contact activity.

Cost (2004 suggested retail price):

$1.68/acre.

Rate:

0.6 kg/acre (one bag treats 41 acres).

Water Volume:

Minimum of 10 gallons/acre (45 L/acre). Higher water volumes may be required later in the growing season. Use sufficient water volume to thoroughly cover all foliage.

Effects of Weather:

Do not apply when rain or night frost is expected. Do not apply if temperature is above 27°C (in shade) and high humidity prevails, or if any of the above conditions are expected within 3 days after treatment. Do not apply under intense sunshine. Do not apply when weather favours drift.

Tank Mixes:

None registered.

Do not mix with dinitro compounds, tetradifon or oils.

Restrictions:

Preharvest Interval: 1 day.

Application: Do not apply by air. Avoid drift onto neighbouring crops.

Storage: Keep away from heat, fire or sparks. Store in cool, dry, locked, well-ventilated area without floor drain.

Caution: Drift to off-target crops may cause leaf burn.

Lance

Fungicide Group – 7
(Refer to page 225)

Company:

BASF Canada PCP# 27495

Crops:

Canola, beans, lentils, chickpeas, potatoes.

Formulation:

70% water dispersible granular (WDG).
Container size - 2 x 2.83 kg per jug.

Crop Stage and Diseases Controlled:

Crop*	Disease	Application Rate (g/acre)	Application Timing
Canola	Sclerotinia stem rot *(Sclerotinia sclerotiorum)*	142	Apply at 20-50% flowering. Apply a second time 7-14 days later up to 50% bloom if disease persists, or weather conditions are favourable for disease development.
	Black spot** (*Alternaria brassicae* and *A. raphani*)	142	Apply at late flowering to early green pod to control black spot.
Dry beans	White mold *(Sclerotinia sclerotiorum)*	227 – 312	Apply at 20-50% flowering. Apply a second time 7-14 days later if disease persists, or weather conditions are favourable for disease development. Use the higher rate to obtain extended protection and maximum yield benefit.
Chickpeas Lentils	Ascochyta blight *(Ascochyta spp.)* White mold *(Sclerotinia sclerotiorum)* Gray mold *(Botrytis cinerea)*	170	Apply at the beginning of flowering. Apply a second time 7-14 days later if disease persists, or weather conditions are favourable for disease development.
Potatoes	Early blight	70 – 126	Apply prior to disease development and at 14 day intervals if conditions continue to favour disease development.

* Refer to the label for crops not detailed above. ** Suppression only.

How it Works:

The active ingredient boscalid is an anilid fungicide. It provides a protective effect because it inhibits spores and spore germination.

Cost:

$22.58 - 49.61/acre.

Water Volume:

Aerial Application: Use a minimum water volume of 40 L/ha. Ensure thorough coverage of foliage

Ground Application: Use a minimum water volume of 100 L/ha and ensure thorough coverage of foliage.

Application:

The product should be used in a preventative spray program and applied as a protectant. When mixing use vigorous agitation. Do not allow the product to sit more then one hour without further vigorous agitation prior to application.

Ground or Aerial Application: Canola, beans, chickpeas and lentils.

Ground Application Only: Potatoes.

Effects of Weather:

Apply under favourable conditions to minimize or avoid the occurrence of drift. Aerial drift is increased under certain meteorological conditions therefore do not apply when wind speed is greater than 16 km/hour at flying height at the site of application

Tank Mixes:

None registered.

Do not tank mix with any other products.

Restrictions:

Crop-Specific Restrictions and Limitations:

Crop*	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre Per Application (grams)	Maximum Number of Applications per season	Maximum Rate Per Acre Per Season (grams)
Beans, Dry	21 days	312	2	624
Canola	21 days	142	2	284
Chickpeas	21 days	170	2	340
Lentils	21 days	170	2	340
Potatoes	30 days	126	4	504

* Refer to the label for crops not detailed above.

Rainfall: If rainfall is imminent, delay application.

Grazing Restrictions: All crops can be grazed or fed to livestock.

Re-Entry Interval: Do not re-enter treated area for 4 hours after application or until dry.

Drift: Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed

Recropping: A plant back restriction of 14 days is required for all crops not on the label.

Hazard Rating:

☠ Caution Poison

Potential Skin Sensitizer

Warning - Eye Irritant

Mancozeb-based Fungicides

Fungicide Group – M
(Refer to page 225)

Dithane DG Rainshield NT/Manzate DF/Penncozeb 75 DF

Company:

Dow AgroSciences (Dithane DG Rainshield NT – PCP# 20553)

E.I du Pont Canada Company (Manzate DF – PCP# 21057)

Cerexagri: distributed by United Agri Products (Penncozeb 75 DF – PCP# 25397)

Formulations:

Dithane DG Rainshield NT - 75 % mancozeb formulated as a dispersible granule. Container size - 20 kg.

Manzate DF - 75 % mancozeb formulated as a dry flowable granule. Container sizes - 20 kg, 2.5 kg.

Penncozeb 75 DF - 75 % mancozeb formulated as a dry flowable powder. Container sizes - 20 kg.

Crops:

Potatoes
Wheat
Lentils (Dithane DG Rainshield NT only)
Seed alfalfa (Dithane DG Rainshield NT only)

Diseases Controlled:

	DITHANE DG RAINSHIELD NT	MANZATE DF PENNCOZEB 75 DF
Wheat	Tanspot, leaf rust, Septoria leaf blotch	Tanspot, leaf rust, Septoria leaf blotch
Lentils	Anthracnose and ascochyta blight	
Potatoes	Early and late blight	Early and late blight
Seed Alfalfa	Leaf spot and stem spot diseases	

Crop Stage:

Potatoes - Apply when plants are 4 to 6 inches (10 to 15 cm) high; repeat at 7 to 10-day intervals. Start with low rate and increase to maximum as foliage develops. The spray interval may be reduced to 5 to 6 days during periods of wet weather favoring late blight and/or vigorous crop growth.

Wheat - May be applied early (when crop is in the 3 leaf to tillering stage) and/or late (when head is fully emerged, but prior to flowering). Do not make more than 2 applications per season.

Lentils – Apply the first application before flower when bud formation is evident. A second application should be applied 10 to 14 days after the initial application, but before rows close in to form a dense canopy. If conditions for disease development persist, a third application may be applied 10 to 14 days later. Do not apply more than 3 applications of Dithane DG per season.

Seed Alfalfa - Apply first application prior to 50% bloom. Repeat 7 to 10 days after the first application and 10 days after the second application. Do not make more than 3 applications per season.

How it Works:

The active ingredient mancozeb is a dithiocarbamate fungicide with contact activity.

Cost (2004 suggested retail prices):

$3.78 to $7.65/acre.

Rates:

Potatoes - 0.45 to 0.90 kg/acre.

Wheat (early) - 0.45 kg/acre; **wheat (late)** - 0.9 kg/acre.

Lentils - 0.9 kg/acre.

Alfalfa - 0.6 kg/acre.

Nozzles:

Dithane DG Rainshield NT - flat fan.

Water Volume:

Thorough, uniform coverage is essential for good disease control.

Manzate DF, Penncozeb 75 DF - 17 to 100 gallons/acre (80 to 400 L/acre) by ground sprayer; 5 to 7 gallons/acre (20 to 32 L/acre) by air.

Dithane DG Rainshield NT - 10 to 18 gallons/acre (45 to 80 L/acre) for ground application. Use 4 gallons/acre (18 L/acre) for aerial application.

Effects of Weather:

Dithane DG Rainshield NT - must be dry on plant leaf prior to rainfall; approximately 1 hour is required without rainfall at moderate temperatures and humidity.

Tank Mixes:

None registered.

For Dithane DG Rainshield NT, add other co-applied fungicides, insecticides, growth regulators, micronutrients and spray adjuvants to the tank last. Do not let Dithane DG Rainshield NT settle out in unagitated spray tank.

Restrictions:

Grazing: Do not graze or feed treated straw or plant tops to livestock.

Preharvest Interval: Potatoes - 1 day.
Lentils – 35 days.
Wheat - 40 days.

Storage: Store in cool, dry, well-ventilated place. Keep away from fire and sparks.

Environment: Do not apply if weather favours drift from areas treated. Do not apply to water or wetlands.

Hazard Rating:

◈ Warning Poison - Eye irritant

Polyram DF

Fungicide Group – M
(Refer to page 225)

Company:

BASF Canada PCP# 20087

Formulation:

80% metiram formulated as dry flowable. Container size - 20 kg.

Crops:

Potatoes.

Diseases Controlled:

Early blight, late blight

Crop Stage:

Apply at 7 to 10 day intervals using 1.10 to 1.75 kg per hectare until plants cover the row. Then increase the rate to 2.25 kg per hectare until tops are killed, or use 1.10 to 1.75 kg per hectare at 5 to 7 day intervals starting when plants are 15 cm high and continuing until top killing. When conditions (rain or dew) favour infections, use the shorter intervals in each case.

How it Works:

The active ingredient metiram is a dithiocarbamate fungicide with contact activity.

Cost (2004 suggested retail price):

$3.71 - 7.51/acre.

Tank Mixes:

Herbicides: None registered.

Fertilizers: Polyram DF can be mixed with fertilizers after a physical compatibility test has been conducted.

Insecticides: Mixtures with diazinon or malathion should be prepared immediately prior to use and not allowed to stand in the tank.

Fungicides: None registered.

Rates:

0.45 to 0.91 kg per acre

Water Volumes:

Ground - 8.8 to 17.6 gallons/acre (40 to 80 L/acre);

Aircraft - 4.9 gallons/acre (22 L/acre).

Nozzles:

Hollow cones or flat fans recommended.

Effects of Weather:

When weather conditions favor disease development the shorter spray intervals may be required.

Restrictions:

Grazing: Do not use treated crop parts for feed or food.

Preharvest Interval: May be applied up to the day before harvest.

Recropping: None.

Storage: Store in a cool, dry place away from flame or sparks. If product becomes wet or overheated, effectiveness is reduced and flammable vapors may be produced. Do not freeze.

Environment: Do not apply when environmental conditions favor drift from treated area. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or otherwise.

Quadris

Fungicide Group – 11

(Refer to page 225)

Company:

Syngenta PCP# 26153

Formulation:

250 g/L azoxystrobin formulated as a flowable suspension concentrate

Container size – 4 x 3.78 L jugs

Crops:

Canola, chickpeas, lentils, peas, beans and potatoes

Diseases Controlled:

Canola - Blackleg, Sclerotinia stem rot and Alternaria black spot.

Chickpeas - Ascochyta blight.

Lentils – Anthracnose and Ascochyta blight.

Field Peas – Mycosphaerella blight

Beans – Anthracnose

Potatoes – Early blight and Late blight; Rhizoctonia stem rot, stolon canker and black scurf

Crop Stage:

Canola

Blackleg: Apply at the 2 to 6 leaf stage of canola.

Sclerotinia stem rot: Apply at early bloom (prior to 30% bloom). This will also suppress alternaria black spot.

Alternaria black spot: Apply at early pod stage (90% petal fall).

Chickpeas, Lentils, Field Peas, Beans

Make first application no later than onset of flowering. Second application 10 to 14 days later.

How it Works:

The active ingredient azoxystrobin is a methoxyacrylate compound (strobilurin) used as a broad spectrum preventative and curative fungicide.

Cost (2004 suggested retail price):

$20.16/acre for control of blackleg and alternaria black spot.
$28.23 to $40.33/acre for control of sclerotinia stem rot.
$20.16/acre per application on legumes.
$20.16 to $32.26/acre per application on potatoes.

Rate:

Canola

Blackleg and alternaria black spot: Use 0.2 L/acre.

Sclerotinia stem rot: Use 0.28 to 0.4 L/acre. Use the higher rate if there is a history of sclerotinia stem rot infection in the area and when conditions favour development (heavy crop canopy, high humidity and/or excessive rain).

Chickpeas, Lentils, Field Peas, Beans

For each application use 0.2L/acre.

Potatoes

Early blight: 0.2 to 0.32 L/acre starting prior to disease development

Late blight: 0.32 L/acre starting prior to disease development

Rhizoctonia stem rot, stolon canker and black scurf: (in-furrow treatment) 4 to 6 mL/ 100m of row, applied at planting.

Water Volume:

Ground: Use sufficient water volume to obtain adequate coverage. Use minimum 9 gallons/acre (40 L/acre). In-furrow treatment in 5 to 15 gallons/acre (20 to 57 L/acre).

Aerial: Use minimum of 4 gallons/acre (18 L/acre). Ensure uniform application.

Tank Mixes

Herbicides: None registered.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: On potatoes, Quadris may be applied with Bravo for the control of early blight. Do not apply sequential applications of this tank mix and do not exceed 3 tank mix applications per season. Do not apply to potatoes later than 2 days before harvest.

Restrictions:

Resistance: Like any fungicide with a single mode of action, the quinone outside inhibitor fungicides, while highly effective, may result in resistant strains of fungi where they have been overused or misused.

When applying multiple applications in one season, pay close attention to the labelled recommendations for resistance management such as using labelled rates, rotating to fungicides outside group 11 and not exceeding the maximum number of applications per season.

Grazing: Do not feed dried pea vines to livestock.

Preharvest Interval: Canola - 30 days. Legumes – 15 days. Potatoes – 1 day; 90 days for in-furrow application

Recropping: Do not plant broadleaf or root crops within 30 days of application. Do not plant cereals within 45 days of application.

Application: In-furrow application: Mount the spray nozzle so that spray is directed into the furrow as a 15-20 cm band just before the seed is covered.

Do not exceed 2 applications or 0.56 L/acre per season on legume crops. Do not exceed 3 applications or 0.96 L/acre per season on potatoes. Do not apply sequential treatments of Quadris to potatoes but apply in alternation with fungicides that have a different mode of action and to which disease resistance has not developed.

Do not apply if rainfall is imminent. Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 15 km/hr at 2 m above ground at the site of application.

Environment: For ground application maintain a 1 m buffer zone between areas sprayed and aquatic systems.

For aerial application allow a 6 m buffer zone. Quadris can be extremely phytotoxic to certain apple varieties. Do not apply where there is the possibility of spray drift reaching apple or crabapple trees.

Storage: Store in a cool, dry, well-ventilated area. Do not store below 0°C

Hazard Rating:

None.

Reason 500SC

Fungicide Group – 11

(Refer to page 225)

Company:

Bayer CropScience PCP# 27462

Formulation:

500 g/L fenamidone. Container sizes – 2, 4 or 10 L.

Crops:

Potatoes.

Diseases Controlled:

Early and late blight.

Crop Stage:

Begin application when plants are 15 to 20 cm high or when disease threatens. Apply a fungicide with a different mode of action within 7 to 10 days after each application using the shorter interval when conditions favor disease development. Ensure even application.

How it Works:

The active ingredient fenamidone is a member of the strobilurin class of chemistry and has preventative and protectant activity, inhibiting spore germination and sporulation.

Cost:

TBA.

Rate:

Apply at 80 mL per acre as a tank mix with either Dithane DG* at 500 g/ac or Bravo 500 at 500 mL/ac. *When using other formulations of mancozeb, adjust application rates to apply 375 g active ingredient per acre.

Water Volume:

21 gallons/acre (80 L/ac).

Tank Mixes:

Herbicides: None registered.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: To be applied ONLY as a tank mix with mancozeb-based fungicides or Bravo 500. Follow mixing instructions provided on the label.

Restrictions:

Resistance: The strobilurin fungicides, while highly effective, have quickly led to resistant strains of fungi where they have been overused or misused. Especially where multiple applications in one season are considered, pay close attention to the labeled recommendations for resistance management.

Rainfall: Reason is rainfast within 2 hours of application.

Preharvest Interval: 14 days.

Recropping: A 30 day plantback interval is required for potatoes and all other crops.

Application: Do not apply by air or through any type of irrigation system. Do not exceed 6 applications or 0.48 L/ac in a year. Alternate with fungicides having a different mode of action.

Environment: Maintain an 8 m buffer zone between areas sprayed and aquatic systems. Toxic to fish and other aquatic organisms; do not apply where runoff is likely to occur.

Storage: Do not allow product to freeze. If stored more than 1 year, shake well before using. Keep away from fire, open flame or other sources of heat. Store in tightly closed container away from fertilizer, seeds, feed or food.

Hazard Rating:

Caution: Eye Irritant.

Ridomil Gold/Bravo

Fungicide Group – 4, M

(Refer to page 225)

Company:

Syngenta PCP# 26443

Formulation:

500 g/L chlorothalonil and 480 g/L metalaxyl-M in a Twin-Pak Jug. Container size - 8.83 L.

Crops:

Potatoes.

Diseases Controlled:

Early blight, late blight, late blight tuber rot, Botrytis vine rot.

Suppression of Pythium leak and Pink rot.

Crop Stage:

Begin preventive applications, early in the season when conditions are favorable for disease, (before infection), but no later than when the plant foliage meets within the row uniformly across the field. Apply a second and third application at 14-day intervals. The labelled rate of a registered contact fungicide should be applied 7 days after each application. Do not make more than three applications per season.

How it Works:

The metalaxyl component is an acylalanine fungicide with systemic activity.

The chlorothalonil component is a chloronitrile fungicide with contact activity.

Cost (2004 suggested retail price):

$29.66/acre.

Rate:

One 8.83 L jug treats 10 acres. The entire contents of the jug must be added to the spray tank or an improper mixture will result.

Water Volume:

Ground: use sufficient water to ensure thorough coverage of foliage. Use a water volume of 20 to 140 gallons/acre (90 to 640 L/acre).

Aerial: use a minimum water volume of 5 gallons water/acre (23 L/acre).

Tank Mixes:

None registered.

Restrictions:

Resistance: Strains of late blight resistant to metalaxyl may develop especially when applied after disease is present. If late blight develops in the field, contact government Potato Specialist or Plant Disease Specialist.

Application: Do not make more than 3 applications per season.

Storage: Protect from excessive heat.

Environment: Do not apply where runoff is likely to occur. Do not use on coarse textured gravelly soils, soils with less than 2% organic matter or in areas where the water table may be high. Avoid application when weather favors drift. Avoid application by ground or air near or around bodies of water. Do not contaminate streams or ponds by spray drift, by cleaning equipment, or disposal of wastes. A buffer zone of 100 m for aerial application and 15 m for ground application should be observed to protect water bodies.

Hazard Rating:

◈ Warning Poison

Warning, causes severe eye damage.

Ronilan EG

Fungicide Group – 2

(Refer to page 225)

Company:

BASF Canada PCP# 24894

Formulation:

50% vinclozolin formulated as an extruded granular fungicide.

Container size - 12 kg (1.2 kg/PVC bag, 2 x 1.2 kg/pouch, 5 x 2.4 kg/pouch per box).

Crops:

Canola, beans.

Disease Controlled:

Canola - Sclerotinia stem rot.

Beans - Sclerotinia (white mold) and Botrytis (grey mold).

Crop Stage:

Canola - Apply once at 20 to 50% bloom, usually 4 to 8 days after first blooms appear in field. For two applications, apply first at early bloom (20 to 30%) with a second application 7 days later at late bloom (70%) if the disease persists or weather conditions are favourable for disease development.

Beans - Apply at early to mid bloom (30 to 50%) with or without a second application 7 to 14 days later at full bloom if disease persists, or weather conditions are favourable for disease development.

How it Works:

The active ingredient vinclozolin is an dicarboximide fungicide with contact activity.

Cost (2004 suggested retail prices):

Canola - $13.42 to $26.83/acre

Beans - $26.83 to $40.25/acre

Rate:

Canola - 0.30 to 0.40 kg/acre. The high rate is used when conditions favor sclerotinia development (heavy crop canopy, high humidity, and/or excessive rain).

For split application, 0.2 kg/acre per treatment at early bloom (20 to 30%) and again 7 days later.

Beans - 0.4 kg/acre if using 2 applications/year, 0.6 kg/acre if using 1 application/year. Do not apply more than 0.8 kg/acre per season.

Water Volume:

Canola (by ground) - 10 to 20 gallons/acre (45 to 90 L/acre).

Canola (by air) - Use minimum 3.6 gallons/acre (16 L/acre).

Beans - Use sufficient spray volume to obtain thorough coverage of foliage. Use 10 to 20 gallons/acre (45 to 90 L/acre). Ground application only.

Pressure:

Minimum 275 kPa (40 psi).

Effects of Weather:

Do not apply if rain or frost is expected. If rainfall is imminent, delay spraying. Do not treat plants at very low temperatures. Do not apply to any crops that have been stressed due to conditions such as flooding, drought, etc.

Tank Mixes:

Herbicides: None registered.

Fertilizers: Do not use a product that contains boron.

Insecticides: Lorsban and Pyrinex. See label for details on rates. Ronilan EG must be added first to mix.

Fungicides: None registered.

Restrictions:

Grazing: Do not allow livestock to graze on treated crop.

Preharvest Interval: 45 days for beans, 40 days for canola.

Storage: Store in original, tightly closed container in cool, dry, locked, well ventilated area without floor drain. Ronilan may be frozen.

Environment: For ground application, maintain 15 m buffer zone between area sprayed and aquatic systems, for aerial application allow a 100 m buffer zone.

Hazard Rating:

Caution Poison

Rovral Flo

Fungicide Group – 2

(Refer to page 225)

Company:

Bayer CropScience PCP# 24378

Formulation:

240 g/L iprodione formulated as a liquid flowable. Container size - 8.4 L.

Crop:

Canola

Diseases Controlled:

Sclerotinia stem rot and Alternaria black spot.

Crop Stage:

Sclerotinia stem rot: Apply when the canola crop is in the 20 to 50% bloom stage. This will be approximately 4 to 8 days after the canola crop begins to flower. Best protection will be achieved when the fungicide is applied at the 20 to 30% bloom stage - prior to petals beginning to fall. Rovral Flo fungicide can be applied until the 50% bloom stage - when the canola crop is at its maximum yellow color, and prior to significant petal fall.

Alternaria black spot: Apply as a single spray at early green pod, or as split application: the first at full bloom, followed by a second application at early green pod stage. Early green pod stage occurs when almost all canola pods are fully formed and still green with only a few flowers or undeveloped pods remaining at the top of the plant. Good coverage of plants is essential.

How it Works:

The active ingredient iprodione is a dicarboximide fungicide with protective and eradicant activity.

Cost (2004 suggested retail price):

$16.46 to $32.66/acre.

Rate:

Sclerotinia stem rot (white mold): 0.63 to 1.25 L/acre. Use of the higher rate is recommended for fields with a history of heavy disease pressure. Split applications: 0.42 to 0.63 L/ac at 20 percent bloom stage followed by 0.42 L/ac at 50 percent bloom stage.

Alternaria black spot: A single application of 0.85 L/acre at early green pod. Split applications: 0.42 L/acre at full bloom followed by 0.42 L/acre at early green pod stage of canola.

Water Volume:

Ground: 10 gallons/acre (45 L/acre)

Air: Not less than 4.5 gallons/acre (20 L/acre)

Nozzles:

Flat fan or hollow cone for ground and properly calibrated aerial equipment.

Tank Mixes:

None registered.

Rovral should not be mixed with other pesticides, adjuvants or fertilizers except where stated.

Effects of Weather:

Do not spray in heavy dew or when rain is imminent within one hour. Spraying should be carried out in crosswind where possible. Avoid spraying in a dead calm and when wind speeds exceed 20 km/hr.

Restrictions:

Application: Do not make more than 2 applications per season

Preharvest Interval: 38 days

Storage: Protect from frost.

Hazard Rating:

▽ Caution Poison

Senator 70WP

Fungicide Group – 1
(Refer to page 225)

Company:

Engage Agro Corporation PCP# 25343

Formulation:

70% thiophanate-methyl formulated as wettable powder. Container size - 2 kg.

Crop:

White beans.

Diseases Controlled:

Sclerotinia (white mold).

Crop Stage:

Apply when conditions favour disease development (i.e. , warm, humid weather and heavy, dense foliage). This usually occurs during early bloom stage, prior to rows closing in. If conditions favouring disease persist, repeat applications may be warranted.

How it Works:

The active ingredient thiophanate-methyl is a benzimidazole fungicide with systemic activity.

Cost: (2004 suggested retail price)

$55.88 to $71.85/acre.

Rate:

0.7 to 0.9 kg/acre (one container treats 2.14 to 1.67 acres).

Water Volume:

10 to 20 gallons/acre (40 to 90 L/acre).

Nozzles:

Ground - Flat fan.

Air - Hollow cone.

Tank Mixes:

None registered.

Senator 70WP is compatible with most pesticides.

Do not mix with lime or other alkaline materials.

Restrictions:

Resistance: Experience has shown that strains of fungi resistant to thiophanate-methyl may develop in treated crops.

Grazing: Do not feed or allow livestock to graze on treated crops.

Storage: Store in a dry place.

Stratego 250EC

Fungicide Group – 3, 11

(Refer to page 225)

Company:

Bayer CropScience PCP# 27528

Formulation:

125 g/L propiconazole and 125 g/L trifloxystrobin formulated as an emulsifiable concentrate. Container size - 2 x 8 L case

Crops:

Spring Wheat (including Hard Red, Durum, Canada Prairie, Soft White) and Winter Wheat

Diseases Controlled:

Septoria leaf blotch, Tan spot, Powdery mildew, Leaf and Stem rust, Stripe rust

Crop Stage:

Single application – 4-leaf stage up to early heading (GS 14 to 55).

Two applications – First at 4-leaf to flag leaf stage. Second before early heading but not within 14 days of first application.

Apply at the very early stages of disease development. Typically, one application from tillering up to flag leaf emergence is required.

How it Works:

The active ingredient propiconazole is a triazole fungicide with broad spectrum systemic activity.

The active ingredient trifloxystrobin is a strobilurin fungicide with broad spectrum preventative activity.

Cost (2004 suggested retail price):

$10.95/acre

Rate:

0.2 L/acre.

Water Volume:

Ground: 9 to 18 gallons/acre (40 to 80 L/acre)

Air: Not less than 4.5 gallons/acre (20 L/acre)

Effects of Weather:

If rain occurs within one hour of application, reapplication may be necessary.

Avoid spraying in a dead calm or when winds are gusty.

Tank Mixes:

None registered.

Restrictions:

Application: Do not make more than 2 applications per season.

Grazing: Do not graze or harvest as forage if 2 applications made in one season. If a single application is made, do not allow livestock to graze or harvest as forage within 30 days after application.

Preharvest Interval: 45 days

Recropping: A plant back restriction of 14 days is required for all crops not on the label

Environment:. Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Hazard Rating:

Danger – Eye irritant, potential skin sensitizer

Tanos 50 DF

Fungicide Group – 11, U
(Refer to page 225)

Company:

E.I. duPont Canada Company PCP# 27435

Formulation:

25% famoxadone and 25% cymoxanil as a dry flowable.

Crops:

Potatoes.

Diseases Controlled:

Early and late blight.

Crop Stage:

Make the first application following one or two applications of a preventative broad spectrum fungicide such as chlorothalonil or mancozeb. Tanos may be applied on a 7-day interval, but it is recommended to alternate with a fungicide having a different mode of action to prevent resistance.

How it Works:

The active ingredient cymoxanil is a highly active, locally systemic fungicide. It works at several levels of preventative, curative and inhibitive (against sporulation) activity. The active ingredient famoxadone is a member of the strobilurin class of chemistry used as a broad spectrum preventative and curative fungicide.

Cost:

TBA.

Rate:

Apply at 225 to 340 g/ acre.

Water Volume:

Use sufficient water to obtain thorough coverage. With a conventional sprayer use no less than 100 to 120 L/ acre. With an air-assisted sprayer use no less than 44 L/ acre. Ground application only.

Effects of Weather:

Do not apply during periods of dead calm or when winds are gusty. TANOS Fungicide must not be applied to any crop suffering from stress as a result of drought, water logging, low temperatures, insect attacks, nutrient or lime deficiency or other factors reducing crop growth.

Tank Mixes:

None registered.

Restrictions:

Resistance: Plant diseases can develop resistance when exposed to one type of product or even products of similar chemistry. Use cultural practices and fungicide rotation as well as early preventive fungicide applications. Consult the government Potato Specialist or Plant Disease Specialist for disease outbreak forecast and recommendations.

Rainfall: Rainfast within 12 hours of application.

Preharvest Interval: 14 days.

Recropping: Crops that are on the product label (potatoes and field tomatoes) may be planted back at any time. A 30-day plantback interval is required for cereal grains. All other crops may be planted following a 1 year interval.

Application: Do not apply Tanos to more than 500 ac (200 ha) per day. Do not apply by air. Tank mix solutions containing boron may affect product solubility. When using boron containing solutions, add the correct amount of Tanos first and boron containing solution last. Do not make more than 6 applications per year.

Re-entry: Do not re-enter treated areas within 24 hours of application.

Environment: Toxic to fish and aquatic organisms. Observe prescribed buffer zones. Toxic to birds, mammals and harmful to beneficial arthropods. Minimize off-target drift to reduce the effects on wildlife at the field boundary. Do not apply to areas prone to run-off and delay spraying if rainfall is imminent.

Hazard Rating:

◈ Warning Poison

Eye Irritant.

Tattoo C

Fungicide Group – 28, M

(Refer to page 225)

Company:

Bayer CropScience PCP# 24544

Formulation:

375 g/L propamocarb HCl and 375 g/L chlorothalonil formulated as a suspension. Container size - 10 L.

Crops:

Potatoes.

Disease Controlled:

Late blight.

Crop Stage:

Begin applications when conditions are favorable for disease, but before infection, and continue on 7 to 14-day intervals until threat of disease is over. Use the 7-day interval when the risk and conditions for disease are high. To avoid resistance, rotating and alternating applications with fungicides having different modes of action is recommended if multiple fungicide applications are required. Make no more than 3 applications per season.

How it Works:

The active ingredient propamocarb HCl is a carbamate fungicide with systemic activity. Chlorothalonil is a chloronitrile fungicide with contact activity.

Cost (2004 suggested retail price):

$27.14/acre

Rate:

1.09 L per acre.

Water Volume:

17.8 to 26.7 gallons/acre (80 to 120 L/acre).

Tank Mixes:

None registered.

Restrictions:

Resistance: Plant disease fungi can develop resistance when exposed to one type of product or even products of similar chemistry. Use cultural practices and fungicide rotation as well as early preventive fungicide applications. Consult the government Potato Specialist or Plant Disease Specialist for disease outbreak forecasts and recommendations.

Grazing: Do not feed treated crops to livestock.

Preharvest Interval: 7 days.

Recropping: Do not plant a new crop in the treated area within 120 days of the last application.

Application: Do not apply by air. Treatment with any product containing chlorothalonil must be separated by a minimum of 7 days.

Re-entry: Do not re-enter treated areas within 48 hours after treatment. If required, individuals may re-enter treated areas within 48 hours for short tasks not involving hand labour, provided that 4 hours have passed since application and that long pants and a long-sleeve shirt are worn.

Storage: Keep away from fire, open flame or other sources of heat. Do not store below freezing. Store the tightly closed container away from seeds, fertilizers, plants and foodstuffs.

Environment: Do not apply directly to water or areas where surface water is present. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment wash waters. Allow a buffer zone of 15 m around bodies of water when applying.

Hazard Rating:

Danger – Corrosive to eyes

Potential skin sensitizer

Tilt 250E

Fungicide Group – 3
(Refer to page 225)

Company:

Syngenta PCP# 19346

Formulation:

250 g/L propiconazole formulated as an emulsifiable concentrate. Container sizes - 5 L, 4 x 5 L.

Crops:

Winter and spring wheat (including hard red, durum, Canada Prairie, soft white)
Spring Barley
Oats
Canaryseed
Canola
Beans

Diseases Controlled:

WHEAT	BARLEY	OATS	CANOLA	BEANS
Septoria leaf spot Tan spot Septoria glume blotch Stripe rust Powdery mildew Leaf and stem rust	Spot blotch Net blotch Scald Leaf rust Stem rust Septoria leaf spot Powdery mildew	Septoria leaf blotch Crown rust	Blackleg	Rust

Diseases Suppressed:

Canaryseed – Septoria leaf mottle

Crop Stage:

Wheat, barley, oats: Apply at the very early stage of disease development, anytime from the beginning of stem elongation to before the head is half emerged. Best results have been achieved when applied just when the flag leaf emerges. Tilt 250E last about 3 weeks in the plant. If conditions favourable to disease continue, another application may be necessary. The second application is usually applied at the time of head emergence.

Canaryseed: Apply at emergence of flag leaf.

Canola: Apply during the rosette stage (after second true leaf and prior to bolting).

Beans: Apply at the first detection of disease in the field. A second application may be made 14 to 21 days later.

How it Works:

The active ingredient propiconazole is a triazole fungicide with broad-spectrum systemic activity.

Cost (2004 suggested retail price):

$13.97/acre

Rate:

0.2 L/acre. One 5 L container treats 25 acres.

Water Volume:

Ground: Minimum 18 gallons/acre (80 L/acre).

Air: 4 to 5 gallons/acre (16-20 L/acre).

Nozzles:

Ground: Flat fan.

Air: Flat fan or hollow cone.

Tank Mixes:

Herbicides: In wheat and barley only, Tilt 250E may be applied with: 2,4-D amine, estimine 2,4-D, MCPA amine, estimine MCPA, Buctril-M or Pardner. In wheat only, Tilt 250E may be applied with Horizon.

To ensure weed and disease control:

1. Weeds and crops must be at correct growth stage.
2. Tank mixes of Tilt 250E and Buctril-M, Horizon or Pardner can only be applied by ground.
3. 2,4-D and MCPA formulations can be applied by ground or air.

Fertilizers: Tilt 250E may be applied with small amounts of nitrogen. Rate of actual N not to exceed 4 kg/acre (9lb/acre). Add the nitrogen to the spray tank before adding Tilt 250E. Excessive nitrogen or application during hot weather may result in crop injury. Do not add nitrogen when tank-mixing Tilt 250E with a herbicide.

Insecticides: None registered.

Fungicides: None registered.

Effects of Weather:

If rainfall occurs within 1 hour of application, reapplication is necessary. For aerial application, wind speed should be less than 18 km/hr to avoid drift. High humidity and low temperature (10 to 20°C) allow for better deposition of spray droplets.

Restrictions:

Grazing: Do not graze animals on treated green crops within 3 days of application. Do not feed straw treated with herbicide tankmixes to livestock.

Preharvest Interval: Wheat, oats, barley - 45 days. Canola - 60 days. Beans – 28 days

Recropping: None.

Storage: Do not freeze.

Environment: Do not contaminate food or feed. This pesticide is toxic to fish. Do not spray any body of water by direct application, drift or by cleaning equipment.

Hazard Rating:

Caution Poison

Seed Treatment Directory

Table 3. Seed Treatments in Field Crops

								CROP DISEASES/INSECTS CONTROLLED																													
	POTATO SEED PIECE							SEED AND SEEDLING ROTS/BLIGHTS														SMUTS CARRIED ON SEED							SMUTS IN SEED		Blackleg	Alternaria Black Spot	INSECTS				
PRODUCTS	Page	Blackleg	Black Scurf/Silver Scurf	Common Scab	Seed Piece Decay	Storage Rot	Verticillium Wilt	Canola (Rapeseed)	Cereals	Chickpeas	Corn	Field Beans	Flax	Forage Legumes	Grasses	Lentils	Mustard	Peas	Soybeans	Sunflowers	Triticale	Barley - Covered	Barley - False Loose	Corn - Head Smut	Oats - Smut	Rye - Bunt	Rye - Stem Smut	Wheat - Bunt/Common Smut	Barley - Loose Smut	Wheat - Loose Smut	Canola (Rapeseed)	Canola (Rapeseed)	Wireworms	Seed Corn Maggot	Flea Beetles	Root Maggots	Colorado Potato Beetle
Agrox B-2, Agrox CD	255										●	●						●	●															●			
Apron FL, Allegiance FL	256							●	●	●	●	●		●	●	●		●	●	●[4]																	
Apron Maxx RTA	258									●		●[3]				●		●	●																		
Baytan 30	259								●[1]													●	●					●	●	●							
Captan Flowable	260										●	●						●	●																		
Charter	261								●													●	●		●			●	●	●							
Crown	262									●[2]						●[2]																					
Cruiser 5FS	263																																●[5]				
DB-Red L	264								●													●	●		●	●		●									
DCT	265																																			●	
Dithane F-45*	266				●	●																															
Dividend XL RTA	266								●[1,7]													●	●					●		●							
Foundation Lite	268							●									●														●	●					
Gaucho CS FL, Gaucho 480 FL	269							●[8]									●[8]														●[8]	●[8]			●		
Helix, Helix XTra	271							●									●														●	●			●		
Imidacloprid-based Insecticides* (includes Genesis 240 and Admire SPT)	272																																				●[5]
Mancozeb-based Fungicides* (includes Potato ST16, MancoPlus, Tuberseal)	273				●																																
Maxim 480FS	274							●	●	●	●	●	●		●	●	●	●	●	●	●																
Maxim PSP*	275		●		●																																
Mertect SC*	276		●		●	●																															
Polyram 16D*	277			●	●																																
Prosper FL/Poncho 600	278							●[6]			●[6]																						●	●	●		
Raxil FL, Raxil 250 FL	279								●													●	●		●			●	●	●							
Senator PSPT*	281	●	●		●		●																														
Thiram 75WP	282										●	●		●	●		●	●	●																		
Vitaflo 280	283								●[7]		●	●[3]	●			●		●[2]	●		●	●	●	●	●		●	●	●	●							
Vitavax rs Fungicide	285							●									●														●	●					
Vitavax Single Solution	286								●				●									●	●		●		●	●	●	●							

[1] wheat and barley only.
[2] also seed-borne ascochyta.
[3] also seed-borne anthracnose.
[4] also controls downy mildew.
[5] also controls flea beetle, potato leafhopper, and aphids.
[6] controlled only with Poncho products.
[7] also controls seed-borne septoria.
[8] only controlled with Gaucho CS FL.

* Before using any pesticide on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

Agrox B-2/Agrox CD

Fungicide Group – M
(Refer to page 225)

Company:

Norac Concepts Inc.
(Agrox B-2 – PCP# 26956; Agrox CD – PCP# 26957)

Formulation:

Agrox B-2: 33.5% captan, 11% diazinon.

Agrox CD: 15% captan, 15% diazinon.

Products formulated as wettable powders.

Container sizes: Agrox B-2 - 2.0 kg; Agrox CD - 0.2 kg, 0.6 kg, 1 kg

Crops and Staging:

Corn, beans, soybeans, peas. Seed treatment.

Diseases Controlled:

Agrox B-2: Seedling blight and seed rot.

Agrox CD: To supplement previous fungicide treatment for seedling blight and seed rot.

Insects Controlled:

Seed corn maggot.

Cost (2004 suggested retail prices):

Agrox B-2: $3.46 per 25 kg seed ($3.68 for corn)

Agrox CD: $2.81 per 25 kg seed

Rate:

Agrox B-2: Dry application: 0.08 kg (0.085 kg for corn) per 25 kg of seed. **Slurry machines:** adjust to apply 0.31 L (0.315 for corn) of slurry per 25 kg of seed and use 0.25 kg of seed treatment per L of water. **Mixing in bucket:** 0.084 kg seed treatment per 0.5 L of water per 25 kg of seed.

Agrox CD: 0.05 kg per 25 kg of seed (one 0.6 kg container treats 300 kg, one 0.2 kg container treats 100 kg).

How it Works:

The active ingredient captan is a phthalimide fungicide with contact activity. The diazinon component is an organophosphate insecticide with contact and stomach activity.

Registered Mixes:

Use Agrox CD only on seed previously treated with captan or thiram; do not use on seed already treated with an insecticide (other than methoxychlor or malathion). Do not use Agrox B-2 on seed already treated with an insecticide.

Restrictions:

Use: Seed within one month of treatment.

Storage: Store in a cool, dry place. Treated seed must be labeled. Label unplanted treated seed as follows: **"Poisonous to man and animals. This seed has been treated with diazinon for the control of insects, and captan for disease control. Do not use for food or feed purposes. Do not sell to oil mills."**

Resistance: Any insect or fungal population may contain individuals naturally resistance to Agrox B-2 and Agrox CD. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Rotate these products with insecticides/fungicides with a different mode of action.

Precautions:

Do not mix with unprotected hands; use rubber gloves. Wash hands, face and arms thoroughly after handling. Do not breathe in mist of dust. Work in a well-ventilated area and wear a suitable respirator, goggles and gloves. Do not contaminate food, feed or any body of water.

Hazard Rating:

☠ Danger Poison

Apron FL/Allegiance FL

Fungicide Group – 4
(Refer to page 225)

Company:

Bayer CropScience
(Apron FL – PCP# 24262; Allegiance FL – PCP# 26674)

Formulation:

317 g/L metalaxyl formulated as a liquid seed treatment. Container size: 3.79 L.

Crops and Staging:

Alfalfa, beans, chickpeas, canola, clover, corn, grasses, low tannin lentils, peas, sainfoin, soybeans, sunflowers, and vetch. Crops for export only - wheat, barley, oats, rye, bird's-foot trefoil and sorghum.

Seed treatment.

Diseases Controlled:

Seed rots and seedling blights caused by *Pythium* spp., early season *Phytophthora* of soybeans, and downy mildew of sunflowers. Also (for export purposes only) downy mildew of peas, corn, and sorghum.

Cost (2004 suggested retail price):

$195.78 per L.

Rates and Water Volumes:

Use the high rate if planting into cold, wet soils.

Rates for crops processed in Canada:

CROP	APRON FL or ALLEGIANCE FL (L per 100 kg seed treated)	WATER (L required to make up a total volume of 0.5 L)	KG SEED TREATED PER 3.79 L JUG
Chickpeas*, dry peas*	0.016 - 0.11	0.484 - 0.39	23,625 - 3,436
Canola (rapeseed)*, processing peas	0.032 - 0.11	0.468 - 0.39	11,812 - 3,436
Sainfoin, vetch, alfalfa, beans, clover, corn	0.046 - 0.11	0.454 - 0.39	8,217 - 3,436
Grasses (forage), soybeans	0.046 - 0.093	0.454 - 0.407	8,217 - 4,064
Grasses (turf)	0.093	0.407	4,064
Sunflowers (high rate is for downy mildew control)	0.11 - 0.189	0.39 - 0.311	3,436 - 2,000
Low tannin lentils	0.016	0.484	23,625

*low rate provides good *Pythium* protection for 2 to 3 weeks.

Rates for crops to be exported:

CROP	APRON FL or ALLEGIANCE FL (L per 100 kg seed treated)	WATER (L required to make up a total volume of 0.5 - 0.62 L)	KG SEED TREATED PER JUG
Corn (downy mildew)	0.189 - 0.62	0.311 - 0	2,000 - 609
Peas (downy mildew)	0.146	0.354	2,589
Sunflowers (downy mildew)	0.62	0	609
Cereals (*Pythium*)	0.046 - 0.11	0.454 - 0.39	8,217 - 3,436
Sorghum (*Pythium*)	0.093 - 0.11	0.407 - 0.39	4,064 - 3,436
Sorghum (downy mildew)	0.189	0.311	2,000
Bird's-foot trefoil	0.046 - 0.11	0.454 - 0.39	8,217 - 3,436

Registered Mixes:

Compatible on the seed with the following products:

Alfalfa, Peas	Thiram 75WP
Beans	Vitaflo 280
Low tannin lentil	Crown

Application Information:

Apron FL and Allegiance FL should be mixed with water to form a slurry seed treatment. Apron FL and Allegiance FL contain no colourant. An appropriate colourant such as Prolzed Colourant must be added to slurry before treating seed.

When preparing the slurry the following procedure should be used:

1. Partially fill the mixing tank with water.
2. Add the required quantity of Apron FL or Allegiance FL onto the water surface.
3. Allow products to disperse and switch on agitation.
4. Top up with extra water to required volume and maintain agitation during use.
5. Add colourant last.

Treatment of highly mechanically damaged, poor quality or low vigour seed may result in reduced germination and/or reduced seed and seedling vigour. If seed lot quality is unknown conduct a germination test prior to treating.

How it Works:

The active ingredient metalaxyl is an acylalanine fungicide with systemic activity.

Restrictions:

Use:

1. Treated seed must not be used for food, feed or oil processing.
2. Treated seed may be toxic to birds and other wildlife. Clean up any spilled seed. Ensure that treated seed is properly incorporated at planting.
3. Treated seed must be labelled as follows; **"This seed has been treated with Apron FL or Allegiance FL seed protectant which contains metalaxyl. Do not use for feed, food or oil processing."**

Storage: Do not store above 35°C or below 0°C. Store in original container, away from pesticides, food or feed. All bags containing seed for export must be labelled "FOR EXPORT ONLY."

Grazing: Do not graze or feed livestock on treated areas for 4 weeks after planting.

Resistance: Experience has shown that strains of fungus resistant to metalaxyl may develop. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Rotate Apron FL and Allegiance FL with fungicides with a different mode of action in your disease control program.

Hazard Rating:

Warning – Skin and eye irritant.

Apron Maxx RTA

Fungicide Group – 4, 12
(Refer to page 225)

Company:

Syngenta – PCP# 27577

Formulation:

0.73% fludioxonil plus 1.10% metalaxyl-M formulated as a liquid seed treatment.

Crops and Staging:

Chickpea, lentil, dry beans, dry peas, and soybeans. Seed treatment.

Diseases Controlled:

Seedling blights (damping-off) and seed rots caused by *Pythium*, *Fusarium* and *Rhizoctonia* species. Dry beans only: anthracnose caused by *Colletotrichum* sp.

Cost (2004 suggested retail price):

$38.75 per L; $3.15 per 25 kg of seed

Rate:

81.3 mL per 25 kg of seed (325 mL per 100 kg of seed)

Compatibility with Rhizobia-based Inoculants:

Apron Maxx RTA is compatible with Rhizobia-based inoculant. Check with inoculant manufacturer for details prior to use. Mixing with inoculants may increase drying time while treating. Recalibrate the seed drill before planting treated seed.

Application Information:

Apron Maxx RTA is a ready-to-apply seed treatment formulation for use in commercial seed treatment plants and for on-farm treatment using standard gravity flow or mist type seed treatment equipment. Also used in treat-on-the-go air seeders. Ensure uniform coverage of the seed, as uneven seed coverage may not give the desired level of disease control. Treatment of highly damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Allow the seed to dry before bagging, storing or seeding.

How it Works:

Fludioxonil is a phenylpyrrole fungicide with contact activity. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseased caused by the Oomycetes, including *Pythium* damping-off.

Restrictions:

Use: All treated seed must be labelled **"This seed has been treated with fludioxonil and metalaxyl-M fungicides".** Do not use for food, feed or oil purposes.

Storage: Store away from feeds and feedstuffs. Store between 0 and 30°C.

Environment: This product is toxic to fish and other aquatic organisms. Do no apply directly to aquatic habitats and estuarine-marine habitats. Do not contaminate water by cleaning of equipment or disposal of wastes. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up.

Re-planting: Do not plant any crop other than soybeans, dry beans, chickpea, lentils or dry peas within 30 days to fields in which treated seeds were planted.

Resistance: Experience has shown that strains of fungus resistant to metalaxyl-M may develop. A gradual or total loss of pest control may occur over time if this active ingredient is used repeatedly in the same fields. Rotate Apron Maxx RTA with fungicides with a different mode of action in your disease control program.

Precautions:

Keep out of reach of children. Harmful if swallowed. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist. When handling product, contaminated equipment or treated seed, wear long pants, a long-sleeved shirt and chemical resistant gloves. Wear a dust mask or respirator to minimize exposure to dust.

Baytan 30

Fungicide Group – 3

(Refer to page 225)

Company:

Bayer CropScience PCP# 24677

Formulation:

317 g/L triadimenol formulated as a liquid seed treatment. Container size: 3.78 L.

Crops and Staging:

Wheat, barley. Seed treatment.

Diseases Controlled:

Wheat and Winter Wheat: Loose smut, common bunt, powdery mildew (early season control), and suppression of take-all.

Barley: True loose smut, covered smut, false loose smut, leaf stripe, suppression of net blotch, scald and common root rot.

Cost (2004 suggested retail price):

$3.10 to $6.20 per 25 kg seed

Rates and Water Volumes:

0.05 L per 100 kg seed will control:

Barley - smuts, leaf stripe and suppression of common root rot and scald

Spring wheat - smuts, powdery mildew

Winter wheat - smuts

0.1 L per 100 kg of seed will control:

Barley - suppression of net blotch

Winter wheat - powdery mildew and suppression of take-all

BAYTAN 30 (L per 100 kg seed)	WATER REQUIRED (L)	SLURRY RATE (L)	KG SEED TREATED PER JUG
0.05	0.2 - 0.45	0.25 - 0.5	7,460
0.1	0.15 - 0.4	0.25 - 0.5	3,780

Registered Mix:

Baytan 30 can be applied on seed treated with Vitaflo 280.

How it Works:

The active ingredient triadimenol is a triazole fungicide with systemic activity.

Restrictions:

Use: Treated seed must be labelled **"This seed has been treated with Baytan 30 containing triadimenol. Do not use for feed, food or oil processing."**

Storage: Do not store treated seed. Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed.

Grazing: Do not graze livestock on treated areas for 40 days after planting.

Resistance: Baytan 30 contains a triazole fungicide. Some loss of disease control may occur over time if Baytan 30 or other fungicides in the triazole group are used repeatedly or consecutively in successive years on same fields, due to the development of resistant strains of pathogens. Alternate use of Baytan 30 with fungicides with a different mode of action in your disease control program.

Comments:

Seed treated with this product will sometimes emerge at a slower rate than untreated seed. Avoid late planting of treated winter wheat. Germination and seed vigour may also be reduced under adverse environmental conditions such as cool, wet soil or seed planted deeper than 4 cm. Baytan 30 contains no colourant. An appropriate colourant such as Prolzed Colourant must be added before treating seed. No colourant is required if applying over seed treated with Vitaflo 280.

Captan Flowable

Fungicide Group – M
(Refer to page 225)

Company:

Norac Concepts Inc. PCP# 12028

Formulation:

30% captan formulated as a flowable seed treatment.

Container sizes: 20 L; 1000 L returnable container

Crops and Staging:

Corn, beans, peas, soybeans. Prior to storage or as a seed treatment.

Diseases Controlled:

Storage rot, seed decay, root rot, damping off, seedling blight.

Cost (2004 suggested retail prices):

$0.79 per 25 kg of bean, pea or soybean seed.

$0.34 to $0.56 per 25 kg corn seed.

Rates:

CROP	CAPTAN FLOWABLE (L per 25 kg seed)
Beans	0.07
Corn (field)	0.03* - 0.05
Peas	0.07
Soybeans	0.07

*At this rate product is to be applied only by a professional applicator using equipment which will assure complete and uniform coverage.

Application Information:

Mix the recommended amount of Captan with the amount of water required for the slurry treater equipment to be used. Seed treated by the slurry method should not be bagged or stacked until it has dried.

How it Works:

Captan is a phthalimide fungicide with protective activity.

Restrictions:

Storage: Do not freeze. Product must be stored at ambient temperatures above 0°C and must not be stored with herbicides, feed, food or fertilizer. All treated seed should be labelled, **"Poisonous to man and animals. Do not use for food or feed. Do not sell to oil mills. This seed has been treated with Captan."**

Precautions:

Never handle material with bare hands. When treating, auguring or handling, work in a well-ventilated area and wear a respirator, goggles, rubber gloves and clothing that completely covers arms and legs.

Charter

Fungicide Group – 3
(Refer to page 225)

Company:

BASF Canada PCP# 26455

Formulation:

25 g/L triticonazole formulated as a liquid flowable seed treatment. Container size: 10 L (containing 3.1 L Charter); 200 L (containing 66.7 L Charter) returnable container

Crops and Staging:

Wheat, barley and oats. Seed treatment.

Diseases Controlled:

CROP	DISEASES CONTROLLED	DISEASES SUPPRESSED
Wheat	Seed rot and Seedling blights caused by seed and soil-borne *Fusarium* spp., Loose smut, Common smut	*Fusarium* crown and root rot
Barley	Seed rot and Seedling blights caused by seed and soil-borne *Fusarium* spp. Loose smut, Covered smut False loose smut	*Fusarium* crown and root rot
Oats	Loose smut, Covered smut	

Cost (2004 suggested retail price):

$1.60 per 25 kg of seed

Rate:

25 mL concentrate per 25 kg of seed (75 mL diluted product per 25 kg seed)

Application Information:

Charter is a concentrate seed treatment formulation for use in commercial seed plants, in on-farm standard gravity flow or mist type treatment machines, and in on-the-go air seeder treatment systems. Concentrate must be diluted with water to provide adequate application volume to ensure good coverage of the seed. For best results, mix 2 parts water with 1 part Charter while adding sufficient dye to ensure optimal seed colouration. Uneven seed coverage may result in poor levels of disease control. Seed should be well conditioned and cleaned before treating. Treated seed should not require drying after treatment.

How it Works:

The active ingredient triticonazole is a triazole fungicide that provides systemic broad spectrum protection against seed and soil-borne diseases.

Restrictions:

Storage: Store product in the original container. Keep container closed. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizer and seed. Protect from frost and freezing. Store treated seed under cool, dry conditions. Do not store treated seed for more than 18 months as it may decrease in germination at a faster rate than untreated seed. Treated seed stored for more than 6 months should be tested for germination before use. Treated seed should be labelled **"This seed has been treated with Charter, do not use for food, feed or oil processing"**

Environmental: Do not feed or otherwise expose wildlife or birds to treated seed. Ensure proper disposal of any surplus treated seed not intended for later planting. Do not contaminate domestic, irrigation or natural bodies of water.

Resistance: Charter contains a triazole fungicide. Some loss of disease control may occur over time if Charter or other fungicides in the triazole group are used repeatedly or consecutively in successive years on same fields, because of the development of resistant strains of pathogens. Alternate use of Charter with fungicides with a different mode of action in your disease control program.

Precautions:

When handling Charter or treated seed wear chemical resistant gloves and coveralls. Avoid inhaling or breathing dust. Work in a well-ventilated area or wear a dust filtering respirator in non-ventilated areas. Wash hands and exposed skin thoroughly after handling the concentrate and after application.

Hazard Rating:

☠ Caution Poison.

Crown

Fungicide Group – 1, 7
(Refer to page 225)

Company:

Bayer CropScience PCP# 23430

Formulation:

92 g/L carbathiin and 58 g/L thiabendazole, formulated as a liquid seed treatment. Container size: 10 L.

Crops and Staging:

Lentil and chickpea. Seed treatment.

Diseases Controlled:

Seed-borne ascochyta in lentil caused by *Ascochyta lentis*; seed-borne ascochyta in chickpea caused by *Ascochyta rabiei*. Seedling blights and seed rots caused by *Botrytis*, *Fusarium* and *Rhizoctonia* fungi.

Cost (2004 suggested retail price):

Lentil: $7.88 per 25 kg of seed
Chickpea: $3.94 to $7.88 per 25 kg of seed

Rates:

Lentil: 0.15 L per 25 kg of seed.

Chickpea: 0.075 L to 0.15 L per 25 kg of seed. The lower rate will provide sufficient control of ascochyta in most cases. Use the higher rate for smaller seed size varieties.

Compatibility with Rhizobia-based Inoculants:

Crown is compatible with Rhizobia. Contains a sticker for peat Rhizobia. DO NOT TANK MIX CROWN WITH RHIZOBIA. Crown and Rhizobia can be applied simultaneously to seed through separate systems or applied sequentially. Read inoculant label before use.

Application Information:

Commercial Treaters and On-Farm Auger Treating:
Crown is a ready to use formulation designed for commercial treaters and on-farm auger treating. Crown is added directly to the seed as it enters a mixing chamber or auger. It is important that the seed and chemical be mixed quickly and uniformly. See instructions supplied with the applicable treater system for information on proper application techniques. When a grain auger is used for treating, running the auger less than full is the key to adequate mixing. Augers used for handling treated seed should not be used to move seed for food, feed or oil processing.

Applications to Seed in a Hopper Box or Seed Drill:
Partially fill the hopper box or seed drill with a pre-measured amount of seed. Apply the proper amount of Crown evenly over the seed surface. DO NOT pour in one area. Mix with a paddle until all seed is of a uniform red colour, indicating adequate coverage. DO NOT MIX WITH HANDS. Repeat this procedure until the hopper box or seed drill is filled. Seed can be planted immediately after treatment without drying.

How it Works:

The active ingredient carbathiin is a systemic fungicide and the active ingredient thiabendazole is a benzimidazole fungicide with both contact and systemic activity.

Restrictions:

Use: Treated seed may be toxic to birds and other wildlife. Clean up any spilled seed. Ensure treated seed is properly incorporated at planting. Treated seed must not be used for food or feed.

Storage: Do not store product above 35°C or below 0°C or in direct sunlight. Store in original container only, away from other pesticides, fertilizers, food or feed. Treated seed must be labelled as follows: **"This seed has been treated with Crown seed protectant, which contains carbathiin and thiabendazole. Do not use for food, feed or oil processing."**

Grazing: Do not graze or feed livestock on treated areas for 4 weeks after planting.

Precautions:

When handling Crown or treated seed, work in a well-ventilated area; wear rubber gloves and protective eyewear.

Hazard Rating:

Caution – Eye Irritant.

Cruiser 5FS

Contains insecticide only. Available to commercial seed treaters only.

Company:

Syngenta PCP# 27045

Formulation:

47.6% thiamethoxam. Container size: 1000L, 1050L, Bulk

Crops and Staging:

Wheat and barley.

Insects Controlled:

Wireworms.

Cost:

Not available.

Rate:

33 to 50 mL per 100 kg of seed. Use the higher rate for fields that have a history of moderate to high infestations of wireworms.

Application Information:

For use only in commercial seed treatment facilities with closed transfer systems that provide uniform seed coverage. All seed treated with this product must be conspicuously coloured at the time of treatment. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Allow the seed to dry before bagging or storing in bulk containers.

How it Works:

Thiamethoxam is a seed treatment insecticide in the neonicotinoid class of chemistry and controls wireworms through contact and systemic activity.

Registered Mixes:

For added control of seed and soil-borne pathogens in wheat and barley, Cruiser 5FS can be tank-mixed with Dividend XL RTA as part of a closed system application.

Restrictions:

Use: Treated seed must not be used for food, feed or oil processing. All seed must be labelled **"Seed treated with thiamethoxam insecticide."**

Storage: Store away from food and feed. Ideal storage temperature is above freezing and below 30°C. If product should freeze, bring to room temperature, then ensure the contents are.mixed well prior to application.

Grazing: Do not graze or feed livestock on treated areas for 45 days after planting.

Environmental: This product is toxic to fish and aquatic vertebrates. Do no apply directly to water or to areas where surface water is present. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or accessible to birds, promptly clean up or bury to prevent ingestion.

Precautions:

Keep out of reach of children, unauthorized personnel and domestic animals. Harmful if swallowed, inhaled, or absorbed through skin. Do not get into eyes. Avoid prolonged contact with skin. Avoid inhalation of vapours, spray mist or treated seed dust. Workers involved in treating, cleanup or maintenance of seed equipment must wear chemical-resistant overalls over long sleeved shirt and long pants, chemical-resistant gloves, head gear and respiratory protection.

Hazard Rating:

Caution Poison.

DB-Red L

Fungicide Group – M
(Refer to page 225)

Company:

Agsco Inc., distributed by United Agri Products – PCP# 27144

Formulation:

323 g/L maneb. Container size: 2 x 10 L; 57 L returnable container

Crops and Staging:

Wheat, barley, oats, rye. Seed treatment.

Diseases Controlled and Rates:

CROP	DISEASES CONTROLLED	RATE PER 25 KG SEED
Wheat	Common bunt Root rot Seedling blight (including *Fusarium*)	78 mL
Barley	Covered smut False loose smut Root rot Seedling blight	99 mL
Oats	Covered smut Root rot Seedling blight	138 mL
Rye	Common bunt Root rot Seedling blight	65 mL

Cost (2004 suggested retail prices):

Wheat: $1.41 per 25 kg seed

Barley: $1.79 per 25 kg seed

Oats: $2.49 per 25 kg seed

Rye: $1.17 per 25 kg seed

How it Works:

The ingredient maneb is a dithiocarbamate fungicide with contact activity.

Restrictions:

Use: Use only recommended rates; lower amounts may not give the desired control and higher amounts may cause seed injury. Uneven treating of seed may cause over-treatment of some seed and under-treatment on other seed. Seed should be well-cured, dry and cleaned before treatment.

Storage: Store away from heat source and in a cool, dry area. If product becomes frozen, thaw and shake or agitate. If treated seed is stored, label container, **"This seed has been treated with DB-Red L seed treatment. Do not use for food, feed or processing."** Excess treated seed should not be stored past planting time. Do not contaminate feed or food stuffs.

Grazing: Do not graze or feed livestock on treated areas.

Environment: Do not contaminate ponds, lakes or streams.

Hazard Rating:

☠ Danger Poison

DCT

Fungicide Group – 1, M
(Refer to page 225)

Company:

Norac Concepts Inc. PCP# 14986

Formulation:

6% diazinon, 18% captan, and 14% thiophanate-methyl formulated as a seed treatment powder. Container size: 10 kg.

Crops and Staging:

Field beans. Seed treatment.

Diseases Controlled:

Seedling blight, root rot, seed-borne anthracnose. This product will not control anthracnose if seed is severely infected.

Insects Controlled:

Root maggots.

Cost (2004 suggested retail price):

$5.74 per 25 kg of seed

Rates:

For slurry machines: 0.52 kg per 1 L water. Use 1 L of slurry per 100 kg of seed.

For mixing in bucket: 0.13 kg per 0.35 L water. Use 0.35 L of slurry per 25 kg of seed.

How it Works:

The active ingredient captan is a phthalimide fungicide with protective activity. The active ingredient thiophanate-methyl is a benzimidazole fungicide with systemic activity. The diazinon component is an organophosphate insecticide with contact and stomach activity.

Restrictions:

Storage: Do not store treated seed for more than one month. Treated seed should be coloured and labelled **"Poisonous to man and animals. This seed has been treated with diazinon, captan and thiophanate-methyl for control of insects and seed-borne diseases. Do not use for food or feed purposes."**

Environmental: Do not contaminate any body of water.

Precautions:

May be fatal if swallowed. Never handle material with bare hands. Do not breathe dust. When treating, auguring, or handling, work in a well-ventilated area and wear a respirator, goggles, rubber gloves and clothing that completely covers arms and legs.

Hazard Rating:

☠ Danger Poison

Dithane F-45

Fungicide Group – M
(Refer to page 225)

Company:
Dow AgroSciences PCP# 20552

Formulation:
37% mancozeb formulated as a flowable.
Container size: 10 L

Crop and Staging:
Seed potatoes in storage.

Diseases Controlled:
Fusarium dry rot.

Rate and Application Information:
1.58 L per 1000 kg seed potatoes. Apply in 4 to 8 L of water as a spray on conveyor belt prior to storage.

Cost (2004 suggested retail price):
$7.00 per L

How it Works:
Mancozeb is a dithiocarbamate fungicide with contact activity.

Restrictions:
Storage: Do not allow product to freeze. Keep away from fire and sparks. Store in a cool dry well-ventilated place away from feed or food.

Environment: This product is toxic to fish. Do not apply to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Precautions:
Keep out of reach of children. May cause irritation of nose, throat, eyes and skin.

Dividend XL RTA

Fungicide Group – 3, 4
(Refer to page 225)

Company:
Syngenta PCP# 25777

Formulation:
3.37% difenoconazole, 0.27% metalaxyl-M formulated as a flowable seed treatment.

Container size: 2 x 10 L jugs; 115 L returnable tote.

Crops and Staging:
Spring Wheat, Durum, Winter Wheat and Barley.

Seed treatment.

Rates and Diseases Controlled:

CROP	RATE/ 100 KG SEED	DISEASES CONTROLLED	DISEASES SUPPRESSED[2]	EARLY SEASON FOLIAR DISEASES CONTROLLED[1]
Winter Wheat	650 mL	Common Bunt[3], Dwarf Bunt[3], Loose Smut, Seed-borne *Septoria*, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Common Root Rot (*Cochliobolus* spp.), Take-all, *Fusarium* crown and root rot	*Septoria* Leaf Blotch
	325 mL	Common Bunt[3], Dwarf Bunt[3], Loose Smut, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Common Root Rot (*Cochliobolus* spp.), Take-all, *Fusarium* crown and root rot	
Spring Wheat	650 mL	Common Bunt[3], Loose Smut, Seed-borne *Septoria*, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Common Root Rot (*Cochliobolus* spp.), Take-all, *Fusarium* crown and root rot	
	325 mL	Common Bunt[3], Loose Smut, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Common Root Rot (*Cochliobolus* spp.), Take-all, *Fusarium* crown and root rot	
Barley	650 mL	Seed-borne *Septoria*, False Loose Smut, Covered Smut, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Take-all, *Fusarium* crown and root rot	
	325 mL	False Loose Smut, Covered Smut, Seed-borne and soil-borne *Fusarium*, General seed rots[4], *Pythium* Damping-off	Take-all, *Fusarium* crown and root rot	

[1] Dividend XL RTA provides control of early season *Septoria* leaf blotch in winter wheat for the first six (6) weeks after planting. For full season control, apply a foliar fungicide according to label directions.

[2] Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

[3] Dividend XL RTA controls both seed and soil-borne common and dwarf bunt.

[4] General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* and *Aspergillus*.

Cost (2004 suggested retail price):

$22.34 per L; $1.82 to $3.63 per 25 kg of seed

Application Information:

Dividend XL RTA is a ready-to-apply formulation for use in commercial seed treatment plants, and for on-farm treatment using standard gravity flow or mist-type seed treatment equipment which accurately meters and mixes a flowable seed treatment. The equipment must provide uniform coverage of Dividend XL RTA on the seed. Uneven seed coverage may not give the desired level of disease control.

Consult the manufacturer of the application equipment for suitability for this application and for instructions on operation and calibration of the equipment.

Water Volume: Dividend XL RTA (ready to apply) does not require addition of water for application. However, when using the Flexi-Coil Seed Treatment Unit, Dividend XL RTA must be diluted with water to reach 99 mL/10 kg of seed. The recommended dilution rate is 1 part Dividend XL RTA to 2 parts water. This is equal to 33 ml of Dividend plus 66 ml of water in order to achieve the total liquid volume requirement of 99 ml/10 kg seed.

How it Works:

The active ingredient difenoconazole is a systemic fungicide from the triazole fungicide class that provides broad-spectrum protection against seed and soil-borne diseases. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including *Pythium* damping-off.

Registered Mixes:

For the control of true loose smut in barley (*Ustilago nuda*), tank mix Dividend XL RTA with Charter, Vitavax Single Solution, Raxil 250FL or Baytan 30. Read the label directions for each product and follow the more restrictive label precautions and limitations.

Restrictions:

Use: Treated seed should be labelled **"This seed has been treated with Dividend XL RTA; do NOT use for food, feed, or oil purposes."**

Storage: Keep in heated storage. Product will freeze at -18°C. Store away from feeds and food stuffs.

Grazing: Do not graze, feed green forage or cut for hay within 55 days of planted wheat or 39 days of planting barley.

Re-cropping: Do not plant any crop other than wheat and barley within 30 days to fields in which treated seeds were planted.

Environment: Do not apply directly to water or to areas where surface water is present. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Resistance: Dividend XL RTA contains a triazole fungicide. Some loss of disease control may occur over time if Dividend XL RTA or other fungicides in the triazole group are used repeatedly or consecutively in successive years on same fields, due to the development of resistant strains of pathogens. Alternate use of Dividend XL RTA with fungicides with a different mode of action in your disease control program.

Precautions:

Keep out of reach of children. When handling Dividend XL RTA, contaminated equipment or seed treated with Dividend XL RTA, wear long pants, a long-sleeved shirt and chemical resistant gloves. In addition, wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.

Hazard Rating:

None.

Foundation Lite

Fungicide Group – 2, M

(Refer to page 225)

Company:

Bayer CropScience PCP# 25592

Formulation:

132 g/L iprodione and 88 g/L thiram.

Container size: 100 L, 200 L, 1000 L.

Crop and Staging:

Canola, mustard. Seed treatment.

Diseases Controlled:

Damping off and root rot caused by *Rhizoctonia* solani, seed-borne blackleg and seed-borne *Alternaria* black spot.

Cost (2004 suggested retail price):

$19.42 per 25 kg of seed

Rate:

563 mL per 25 kg seed; 100 L container treats 4,444 kg of seed

Application Information:

Roll drum or stir well before using. Foundation Lite treated seed should not require drying after treatment and can be stored or bagged immediately. Add product to seed slowly and mix well for 3-5 minutes in treating equipment. Thorough seed coverage is required.

How it Works:

The active ingredient iprodione is a dicarboximide fungicide with contact and eradicant activity. The active ingredient thiram is a dithiocarbamate fungicide with contact activity.

Restrictions:

Use: Treated seed must be labelled **"This seed has been treated with Foundation Lite containing iprodione and thiram. Do not use for feed, food or oil processing."** Date of treatment must also be included.

Storage: Protect product from frost. Store treated seed in cool dry conditions. Do not store treated seed for more than 6 months. Treated seed stored for periods in excess of 6 months may decrease in germination at a faster rate than untreated seed. Treated seed stored for more than 6 months should be tested for germination before planting.

Environment: Do not contaminate domestic or irrigation waters. Do not contaminate feed or food stuffs.

Precautions:

Keep out of reach of children; avoid contact with skin and eyes. When handling product or treated seed work in a well-ventilated area, wear chemical resistant gloves, coveralls and wear a suitable respirator. Do not consume alcoholic beverages 24 hours before or after working with Foundation Lite.

Hazard Rating:

Caution Poison

Gaucho CS FL / Gaucho 480 FL

Fungicide Group – 7, M
(Refer to page 225)

Gaucho CS FL and Gaucho 480 FL are available to commercial seed treaters only.

Company:

Bayer CropScience (Gaucho CS FL – PCP# 27174; Gaucho 480 FL – PCP# 26124)

Formulation:

Gaucho CS FL: 285.7 g/L imidacloprid; 47.6 g/L carbathiin; 95.3 g/L thiram. Container sizes: 10, 100, 1000 L.

Gaucho 480 FL: 480 g/L imidacloprid. Container size: 10 L.

Crops and Staging:

Canola, rapeseed and mustard. Seed treatment.

Insects Controlled:

Gaucho CS FL: Flea beetles. For areas where flea beetle populations are often high, tank mix Gaucho CS FL with Gaucho 480 FL. Under high insect pressure, a foliar insecticide may be required, therefore monitor crop regularly for insect infestation levels.

Gaucho 480 FL: Flea beetles. Suppression of Lygus bugs, cabbage seed pod weevil larva, and aphids.

Diseases Controlled:

Gaucho CS FL only: Seed-borne blackleg (*Leptosphaeria maculans*), seed rot and seedling blight caused by *Rhizoctonia*, *Pythium* and *Alternaria* spp.

Cost:

Not available.

Rates:

Gaucho CS FL: 350 mL to 525 mL per 25 kg of seed (1.4 L to 2.1 L per 100 kg seed).

Gaucho 480 FL: 208 mL to 417 mL per 25 kg of seed (0.8 L to 1.7 L per 100 kg seed).

In areas where flea beetle populations are often high, use the higher rates; OR, tank mix the lower rates of Gaucho CS FL and Gaucho 480 FL according to label directions.

Application Information:

For use in commercial seed treaters only. Prior to and during application, seed treatment must be thoroughly agitated to ensure uniform mixing of product. Keep above 10°C prior to and during application. Do not apply direct heat to container. A blue colourant must be added to treated seed.

How it Works:

Imidacloprid is a chloronicotinyl insecticide with systemic activity. The active ingredient carbathiin is carboximide fungicide with systemic activity and the active ingredient thiram is a dithiocarbamate fungicide with contact activity.

Restrictions:

Use: Treated seed must be labelled as follows: **"This seed has been treated with imidacloprid, carbathiin and thiram. Do not use for food, feed or oil processing. Store away from feeds and other foodstuffs."**

Storage: Storage of Gaucho CS FL at low temperatures is not recommended. Treated seed stored for periods in excess of 9 months should be tested for germination before planting. Do not store treated seed above 25°C or in direct sunlight.

Grazing: Do not graze or feed livestock on treated areas for 4 weeks after planting.

Environment: These products are highly toxic to birds and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. Do not contaminate water when disposing of equipment wash water. Cover or incorporate spilled treated seeds. Leftover treated seed should be double sown around the headlands, or buried away from water sources.

Precautions:

Harmful if swallowed, absorbed through skin or inhaled. When handling Gaucho or Gaucho treated seed, work in a well-ventilated area and wear chemical resistant gloves, protective eyewear and a suitable mask.

Hazard Rating:

▽ Caution Poison

Genesis 240/Admire SPT

See Imidacloprid-based Insecticide Potato Seed Treatments on page 272.

Helix / Helix XTra

Fungicide Group – 3, 4, 12
(Refer to page 225)

Helix and Helix XTra are available to commercial seed treaters only.

Company:

Syngenta (Helix – PCP# 26637; Helix XTra – PCP# 26638)

Formulation:

Helix: 10.3% thiamethoxam, 1.24% difenoconazole, 0.39% metalaxyl-M, 0.13% fludioxonil.

Helix XTra: 20.7% thiamethoxam, 1.25% difenoconazole, 0.39% metalaxyl-M, 0.13% fludioxonil.

Formulated as a liquid seed treatment.
Container size: 1000 L returnable tote

Crops and Staging:

Canola, mustard. Seed treatment.

Diseases Controlled:

Seed-borne blackleg, seed-borne *Alternaria*, and the seedling disease complex (damping off, seedling blight, seed rot, root rot) caused by *Pythium*, *Fusarium* and *Rhizoctonia* spp.

Insects Controlled:

Flea beetles:

Helix: 14 to 21 days

Helix XTra: 28 to 35 days

Cost:

Not available.

Rate:

375 ml per 25 kg seed (1.5 L per 100 kg seed).

Application Information:

For use only in commercial seed treatment facilities with closed transfer systems. Helix and Helix XTra are ready to use liquid products and contain an appropriate colourant. Apply using standard commercial seed treatment equipment that provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect or disease control. Consult the manufacturer of the seed treating equipment for advice on the operation and calibration of the equipment. Maintain constant agitation during the seed treatment process. Allow the seed to dry before bagging.

How it Works:

The active ingredient thiamethoxam is a systemic insecticide from the neonicotinoid chemical class. The active ingredient difenoconazole is a systemic fungicide from the triazole chemical class that provides broad-spectrum protection against seed and soil-borne diseases. The active ingredient metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including *Pythium* damping off. The active ingredient fludioxonil is a phenylpyrrole chemistry and has contact and local penetrant properties.

Restrictions:

Use: All treated seed must be labelled **"This seed has been treated with Helix which contains the insecticide (thiamethoxam) and the fungicides (difenoconazole, metalaxyl-M and fludioxonil). Wear long-sleeve shirt, long pants, and chemical resistant-gloves when handling treated seed. Do not use for food, feed or oil processing. Store away from food and feed."**

Storage: Store in a well-ventilated, secure area. Keep out of reach of children.

Environment: Do not apply directly to water, or to areas where surface water is present. In cleaning of equipment or disposing of wastes, do not contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes.

Precautions:

Treat seed in a well-ventilated area. All seed treaters and handlers must wear chemical-resistant coveralls over long sleeve shirt and long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e. half-mask respirator with suitable dust filter or fresh air hood).

Hazard Rating:

▽ Caution Poison

Imidacloprid-based Insecticide Potato Seed Treatments

Includes Admire SPT and Genesis 240. Contain insecticide only.

Company:

Bayer CropScience
(Admire SPT – PCP# 27702; Genesis 240 – PCP# 27349)

Formulation:

240 g/L imidacloprid. Container size: 1 L, 3.785 L

Crops and Staging:

Potato. Seed treatment.

Insects Controlled:

Control of Colorado potato beetle, potato flea beetle, potato leafhopper and aphids (including green peach, buckthorn, foxglove and potato aphid).

Cost (2004 suggested retail price):

Genesis 240: $6.72 to $10.09 per 100 kg of potato seed tubers
Admire SPT: not available

Rate:

26 to 39 ml per 100 kg of potato seed tubers. The higher rate is recommended when extended length of control is needed.

Application Information:

May be applied when potato pieces are being cut. Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system that is well contained and will prevent the loss of any liquid. Do not dilute with any more than 3 parts water to 1 part Genesis 240 or Admire SPT Insecticide. Agitate or stir spray solution as needed. Complete coverage of the seed piece is required for optimal insect control. As part of the seed cutting and treating process, application of a fungicide registered for potato seed treatment or an inert absorbent ingredient is recommended.

How it Works:

Imidacloprid is a chloronicotinyl insecticide with systemic activity.

Restrictions:

Use: Do not apply more than 0.47 L per acre per year. Plant seed pieces as soon as possible after cutting and treating. Avoid prolonged exposure of treated seed to sunlight. Do not use treated seed pieces for food, feed or fodder.

Storage: Store product in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children.

Re-planting: Use a minimum plant-back interval of 30 days for cereals, 9 months for peas and beans, and 12 months for all other food and feed crops. Green manure and other cover crops not intended for human or animal consumption do not require a plant-back interval following treatment. Do not graze or harvest cover crops for food or feed. It is not recommended that this product be used in fields treated with imidacloprid during the previous season.

Environmental: Do not plant treated seed pieces within 15 metres of well-head or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc. This product is toxic to wildlife. Keep out of lakes, streams, ponds, or other aquatic systems. Do not contaminate water when disposing of equipment wash waters. Leftover treated seed should be double sown around the headland, or buried away from water sources such as lakes, streams, ponds or other aquatic systems.

Precautions:

Keep out of reach of children. Harmful if swallowed or absorbed through skin. Applicators and other handlers must wear long-sleeved shirt and long pants, water-proof gloves and shoes plus socks. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist.

Hazard Rating:

▽ Caution Poison

Mancozeb-based Fungicides

Fungicide Group – M
(Refer to page 225)

Includes Potato ST 16, MancoPlus, and Tuberseal

Company:

Norac Concepts Inc. (MancoPlus – PCP# 26157; Tuberseal – PCP# 17042)

United Agri Products (Potato ST 16 – PCP# 24734)

Formulations:

16% mancozeb formulated as a dust. Container size: 10 kg (MancoPlus; Tuberseal); 20 kg (Potato ST 16).

Crops and Staging:

Potatoes. Seed treatment

Diseases Controlled:

Fusarium seed piece decay in potatoes.

Cost (2004 suggested retail prices):

Per 100 kg of seed: $2.55 (MancoPlus); $2.53 (Potato ST 16); $2.40 (Tuberseal)

Rates:

0.5 kg per 100 kg seed.

Application Information:

Apply product before planting. If treated whole seed is cut, make a second application to protect cut surfaces. Plant as soon as possible after treating. If cut seed is not planted immediately, store in a ventilated location to allow cut surfaces to dry.

How it Works:

The active ingredient mancozeb is a dithiocarbamate fungicide with contact activity.

Restrictions:

Use: Treated seed pieces should be labelled **"Poisonous to man and animals. This seed has been treated with mancozeb for control of *Fusarium* decay. Do not use for food or feed."**

Storage: Store product in a cool, dry, well-ventilated place. Keep away from fire and sparks.

Environment: Do not contaminate any body of water.

Precautions:

Keep out of reach of children. May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Wash thoroughly after handling and before eating, drinking or smoking. When treating or handling treated seed, work in a well-ventilated area and wear a suitable dust mask, goggles and gloves.

Hazard Rating:

☠ Warning Poison

Maxim 480 FS

Fungicide Group – 12
(Refer to page 225)

This product is available to commercial seed treaters only.

Company:

Syngenta PCP # 27001

Formulation:

40.3% fludioxonil, as a water-based slurry

Crop and Staging:

Grains: Barley, corn, buckwheat, millet, oats, rye, triticale, wild rice, wheat.

Legumes: Bean, pea, soybean, chickpea, lentil.

Oilseeds: Flax, canola, mustard, rapeseed, safflower, sunflower.

Grasses: All grasses including turfgrass, bermudagrass, bluegrass, fescue, bromegrass. All pasture and grasses grown for hay or silage.

Herbs and Spices: Anise, borage, caraway, coriander, dill, others.

Diseases Controlled:

Seed-borne and soil-borne diseases caused by *Fusarium* spp. (including seedling diseases due to *F. graminearum*) and *Rhizoctonia* spp. on listed crops, as well as *Aspergillus* and *Penicillium* spp. on corn, soybean, canola and mustard.

Seed treatment.

Rates:

1.3 to 2.6 mL to treat 25 kg of seed. Use the higher rate for crops or cultivars that are expected to be more susceptible to seed-borne or soil-borne pathogens.

Cost:

Not available.

Application Information:

For use by commercial seed treaters only. This product contains no colorant. A blue colourant must be added when this product is applied to canola/rapeseed. A red colourant must be added when this product is applied to grain. Shake or mix well before using. Use standard slurry seed treatment equipment to provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of product into the required amount of water for the slurry treater and dilution rate to be used. Maintain constant agitation of slurry during treatment. Allow the seed to dry before bagging.

How it Works:

The active ingredient fludioxonil is a phenylpyrrole fungicide with contact activity.

Registered Mixes:

Maxim 480FS will not control seed rots or blights caused by *Pythium* or *Phytophthora* species and can be mixed with a labelled seed treatment product with activity against these diseases. Consult each label for registered use rates and instructions.

Restrictions:

Use: All treated seed must be labelled **"This seed has been treated with fludioxonil fungicide. Do not use for food, feed, or oil purposes. Store away from feed and foodstuffs. Do not graze treated crops, or cut for forage within 30 days of planting."**

Storage: Store above freezing and below 30°C.

Environment: This product is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present. In cleaning of equipment or disposing of wastes, do not contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes.

Resistance: Maxim 480FS contains a Group 12 fungicide. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Alternate the use of Maxim 480FS (and other Group 12 fungicides) with products with a different mode of action.

Precautions:

Keep out of reach of children. May be harmful if swallowed. Avoid inhalation of vapour, spray mist or treated seed dust. Treat seed in a well-ventilated area. When handling product or seed, wear long-sleeved overalls and chemical-resistant gloves. When bagging or sewing bags of treated seed, wear a dust mask.

Maxim PSP

Fungicide Group – 12
(Refer to page 225)

Company:

Syngenta PCP# 26647

Formulation:

0.5% fludioxonil formulated as a dry powder.
Container size: 10kg

Crops and Staging:

Potatoes. Seed piece treatment.

Diseases Controlled:

Black scurf (*Rhizoctonia solani*), silver scurf (*Helminthosporium solani*), and fusarium dry rot (*Fusarium* spp.)

Cost (2004 suggested retail price):

$11.13 per kg. One 10 kg bag treats 2000 kg of seed.

Rate:

Use 0.5 kg per 100 kg of cut seed.

Application Information:

Apply using appropriate treater designed for treating potatoes or by dust attachment over belt. Cut pieces should be treated immediately after cutting. If treated seed pieces are bagged, they should be stored for 2-3 days in open crates before bagging. For optimum protection against silver scurf, ensure that seed tubers are completely free of soil. Total skin coverage is essential.

How it Works:

The active ingredient fludioxonil is a phenylpyrrole fungicide with contact activity.

Restrictions:

Storage: Store in a dry place. Avoid contamination of feed and foodstuffs.

Environment: Do not contaminate domestic or irrigation water supplies, lakes, streams and ponds.

Resistance: Any fungal population may contain individuals naturally resistant to fludioxonil or strains of some fungi may develop tolerance to fungicides after prolonged use. It is important to follow label rate and directions and to observe all practices that minimize occurrence of resistance. Whenever possible, alternate with products from different chemical families. In order to minimize the potential for the development of resistance among labeled disease organisms do not use Maxim PSP in any two consecutive seed generations.

Precautions:

Avoid inhalation of dust. Wear two layers of clothing, work boots, mask, and chemical-resistant gloves when handling product and treated seed. Wash hands and face after handling. May be harmful if swallowed. Keep out of reach of children.

Hazard Rating:

Caution Poison

Mertect SC

Fungicide Group – 1
(Refer to page 225)

Company:

Syngenta PCP# 13975

Formulation:

500 g/L thiabendazole formulated as a water dispersible suspension. Container size: 4 x 5 L.

Crops and Staging:

Potato. Post-harvest treatment.

Diseases Controlled:

Fusarium dry rot (*Fusarium* spp.), skin spot, silver scurf (*Helminthosporium* spp.), black scurf (*Rhizoctonia* spp.), and pocket rot (*Phoma* spp.)

Cost (2004 suggested retail price):

$75.24 per L

Rate and Water Volume:

7.5 L per 37 gallons (170 L) of water. Spray 2 L of this suspension per 1 metric tonne of potatoes.

Application Information:

Shake well before using. Do not allow suspension to stand without continuous agitation. Potatoes must rotate along conveyor line to ensure complete coverage. Prior to treating potatoes destined for export, confirm with authorities that treated potatoes will be allowed to enter importing country.

How it Works:

The active ingredient thiabendazole is a benzimidazole fungicide with systemic activity.

Restrictions:

Storage: Minimum storage temperature 0°C.

Resistance: Thiabendazole-resistant strains of silver scurf and fusarium dry rot can develop.

Strains of some fungi may develop tolerance to Mertect SC or other products in the same class of chemistry after prolonged usage. When using Mertect SC, use full rates and ensure thorough coverage. If potatoes are going to be treated with Mertect SC prior to storage in the fall, do not use any product containing thiophanate-methyl at planting. Do not use any product containing thiophanate- methyl if Mertect SC was used as the storage treatment on seed. Use seed treatment with a different mode of action on Mertect SC treated seed potatoes.

Precautions:

Wear chemical-resistant gloves when handling product. May cause eye irritation. Avoid contact with eyes, skin and clothing. Keep out of reach of children.

Polyram 16D

Fungicide Group – M
(Refer to page 225)

Company:

BASF Canada PCP# 25867

Formulation:

16% metiram formulated as dust. Container size: 10 kg.

Crops and Staging:

Potatoes. Seed treatment.

Diseases Controlled:

Fusarium seed piece decay; seed-borne common scab (*Streptomyces* spp.)

Cost (2004 suggested retail price):

$51.95 per 10 kg bag; $2.34 to $3.38 per 100 kg cut seed

Rates:

0.45 to 0.65 kg per 100 kg seed. One container treats 2,222 to 1,538 kg of seed.

How it Works:

The active ingredient metiram is a dithiocarbamate fungicide with contact activity.

Application Information:

Apply to entire surface of seed pieces after cutting. Treated potato seed should not be allowed to stand in hot sun or drying wind. Seed should not be planted into hot, dry soil or in cold, wet soil. Plant as soon as possible after treating. If not planted immediately, provide sufficient ventilation to allow cut surface to dry. If not cutting seed, treat at same rate for control of seed-borne common scab, apply a second application to control fusarium seed piece decay.

Restrictions:

Storage: Store product in cool, dry, locked, well-ventilated area without floor drain. Keep away from heat, fire or sparks. Do not allow product to become wet or overheated as this will reduce its effectiveness and may create flammable vapours. Protect from freezing. Do not ship or store near food, feed, seed or fertilizers.

Environment: Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

Precautions:

Keep out of reach of children. May cause irritation to eyes, nose, throat and skin. Avoid inhalation and contact with eyes, skin or clothes. Wear protective equipment and clothing including goggles or face shield, gloves, hat, long-sleeved shirt, trousers and rubber boots.

Potato ST 16/MancoPlus/Tuberseal

See Mancozeb-based Fungicides on page 273.

Prosper FL / Poncho 600 FS/Poncho 600

Fungicide Group – 4, 7, M
(Refer to page 225)

Poncho 600 FS contains insecticide only. Both products available to commercial seed treaters only.

Company:

Bayer CropScience (Prosper FL – PCP# 27564;
Poncho 600 FS – PCP# 27453;
Poncho 600 – PCP# 27449)

Formulation:

Prosper FL: 120 g/L clothianidin; 56 g/L carbathiin; 120 g/L thiram; 4 g/L metalaxyl.
Container sizes: 3.8L, 10L, 100L, 1000L, Bulk

Poncho 600 FS / Poncho 600: 48% clothianidin.
Container sizes: 3.8L, 10L, 100L, 200L

Crops and Staging:

Canola.

Poncho 600 FS/Poncho 600 only: corn.

Insects Controlled:

Flea beetles.

Poncho 600 FS/Poncho 600 only: wireworm, seed corn maggot, black cutworms, corn flea beetle, white grubs.

Diseases Controlled:

Prosper FL only: Seed rot, damping off, seedling blight and early season root rot caused by *Pythium*, *Rhizoctonia*, *Fusarium*; seed-borne *Alternaria* spp.; and seed-borne blackleg (*Leptosphaeria maculans*).

Cost:

Not available.

Rate:

Canola:

Prosper FL: 1250 mL per 100 kg seed for use under low to moderate flea beetle pressure.

Poncho 600 FS / Poncho 600: 250 mL/100 kg seed (for low to moderate flea beetle pressure); 333 mL per 100 kg seed (for moderate flea beetle pressure); 666 mL per 100 kg seed (for high to extreme flea beetle pressure)

Corn: 33.3 to 66.6 mL of product per 80,000 units of seed (0.25 to 0.5 mg a.i. per kernel)

Registered Mixes:

For canola only: Poncho 600 FS/Poncho 600 can be mixed with Prosper FL to provide longer season control of flea beetles and for areas where flea beetle populations are high. Refer to labels for rates and mixing instruction.

Application Information:

Prosper FL and Poncho 600 FS/Poncho 600 are for use in commercial seed treatment facilities with closed transfer systems only. Poncho 600 FS/Poncho 600 contains no colourant; a baby blue colourant must be added when this product is applied to canola. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Allow the seed to dry before bagging or storing in bulk containers.

How it Works:

Clothianidin is a chloronicotinyl insecticide with systemic activity. Carbathiin is a carboximide fungicide with systemic activity; thiram is a dithiocarbamate fungicide with contact activity; and metalaxyl is an acylalanine fungicide with systemic activity.

Restrictions:

Use: Treated seed must be labelled as follows: **"This seed has been treated with clothianidin and/or carbathiin, thiram and metalaxyl. Do not use for food, feed or oil processing. Store away from feeds and other foodstuffs."** Seed grown and harvested from Poncho 600 FS/Poncho 600 and/or Prosper FL treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.

Re-planting: Corn and canola may be replanted at any time. A 1-year plant back interval is required for leafy, root and tuber vegetables. A 30-day plant back is required for cereals, grasses, nongrass animal feeds, soybeans and dry beans.

Environmental: These products are toxic to aquatic invertebrates. Do no apply directly to water or to areas where surface water is present. Do not contaminate water when disposing of equipment wash waters. These products are toxic to birds and mammals. Any spilled or exposed seeds should be incorporated into the soil or otherwise cleaned up from the soil surface.

Precautions:

Keep out of reach of children. Work in a well-ventilated area when handling products or treated seed. Workers must wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, head gear and respiratory protection.

Hazard Rating:

Warning Poison

Raxil FL/Raxil 250 FL

Fungicide Group – 3
(Refer to page 225)

Company:

Bayer CropScience (Raxil 250 FL – PCP# 26138)

Formulation:

Raxil FL: 8.33 g/L tebuconazole

Raxil 250 FL: 6 g/L tebuconazole

Formulated as a flowable seed treatment.
Container size: 2 x10 L, 200 L

Crops and Staging:

Wheat, barley and oats. Seed treatment.

Diseases Controlled:

Wheat: Loose smut, sinking smut or common bunt; seed rot and seedling blights caused by *Fusarium* spp.; suppression of common root rot caused by *Cochliobolus sativus*; suppression of crown rot and root rot caused by *Fusarium* spp.

Barley: False loose, covered and true loose smuts; barley leaf stripe; seed rot and seedling blights caused by *Fusarium* spp. Suppression of crown rot and root rot caused by *Fusarium* spp.

Oats: Loose smut

Cost (2004 suggested retail price):

Wheat, barley: $1.80 to $2.38 per 25 kg of seed

Oat: $2.38 to $3.56 per 25 kg seed

Rates:

	RAXIL FL			RAXIL 250 FL		
CROP	UNDILUTED CONCENTRATE ML/25 KG SEED	DILUTED CONCENTRATE ML/25 KG SEED*	KG SEED TREATED/10L CONCENTRATE OR 10.6 L OF DILUTED MIXTURE*	UNDILUTED CONCENTRATE ML/25 KG SEED	DILUTED CONCENTRATE ML/25 KG SEED*	KG SEED TREATED/10L CONCENTRATE OR 10.6 L OF DILUTED MIXTURE*
Wheat	45 – 60	48 - 64	5,556 – 4,167	63 – 83	66 – 88	4,000 – 3,000
Barley	45 – 60	48 - 64	5,556 – 4,167	63 – 83	66 – 88	4,000 – 3,000
Oats	60 – 90	64 – 95	4,167 – 2,778	83 – 125	88 – 133	3,000 – 2,000

* For dilution information refer to the "triple rinse instructions" on the product label.

Note: For the suppression of common root rot of wheat, use the higher rates per 25 kg seed. Use the higher rates where disease levels are likely to be high or on highly susceptible varieties.

Application Information:

Raxil FL and Raxil 250 FL are ready to use formulations designed for commercial or on-farm treatment with conventional seed treating equipment that can accurately control application rates and provide a good distribution of chemical onto seed. It is important that the seed and chemical are mixed quickly and uniformly. See manufacturer's instructions supplied with the treater system.

Uniform coverage at the correct rate is important for satisfactory results. Under-treatment may lead to loss of efficacy and over-treatment could reduce germination. Seed may be planted immediately after treating.

Treater calibration can be affected by temperature and should be rechecked on a regular basis. Undiluted product can be used at temperatures down to -10°C.

How it Works:

The active ingredient tebuconazole is a systemic triazole fungicide that provides broad-spectrum protection against seed and soil-borne diseases.

Restrictions:

Use: Treated seed should be labelled **"This seed has been treated with Raxil FL / Raxil 250 FL (tebuconazole), do not use for food, feed or oil processing. Wear a long sleeved shirt and long pants (or coveralls) and chemical resistant gloves when handling treated seed."** Do not reuse container for any purpose. Augers used for handling treated seed should not be used to move seed for feed, food or oil processing. Do not reuse bags from treated seed to handle food or feed products.

Storage: If containers of Raxil FL and Raxil 250 FL have been stored for several months, some settling may occur that requires agitation. If 10 L containers are stored they should be shaken.

Grazing: Do not feed or graze livestock on treated areas for 4 weeks after planting.

Environment: Do not contaminate pods, lakes or streams.

Resistance: Raxil FL and Raxil 250 FL contain a triazole fungicide. Some loss of disease control may occur over time if Raxil FL and Raxil 250 FL or other fungicides in this group are used repeatedly or consecutively in successive years on same fields, due to the development of resistant strains of pathogens. Alternate use of Raxil FL and Raxil 250 FL with fungicides with a different mode of action in your disease control program.

Precautions:

Germination will not be affected by treatment as long as over-application does not occur and seed is properly stored. All seed carried from one year to the next should be checked for germination before planting.

Keep out of reach of children. Work in a well ventilated area when treating seed or while augering or handling treated seed. Wear chemical resistant gloves when handling treated seed.

Hazard Rating:

Danger skin irritant.

Senator Potato Seed Piece Treatment

Fungicide Group – 1
(Refer to page 225)

Company:

Engage Agro Corporation PCP# 26236

Formulation:

10% thiophanate-methyl formulated as dust.
Container size: 10 kg.

Crops and Staging:

Potatoes. Seed piece treatment.

Diseases Controlled:

Fusarium rot, *Verticillium* wilt, silver scurf (*Helminthosporium solani*). Also aids in control of seed piece decay and blackleg infections.

Cost (2004 suggested retail price):

$9.90 per kg; $4.95 to treat 100 kg cut seed.

Rate:

Use 500 g per 100 kg cut seed (one bag treats 2,000 kg seed).

Application Information:

Apply in a convenient container or by dust attachment over belt. Cut pieces should be treated within 6 hours of cutting. For optimum control of silver scurf, ensure that seed tubers are completely free of soil. Total skin coverage is essential. If planting is to be delayed more than 1 to 2 days, the treated pieces should be stored for 2 to 3 days in open crates before bagging.

How it Works:

The active ingredient thiophanate-methyl is a benzimidazole fungicide with systemic activity.

Restrictions:

Storage: Store product in a dry place. Avoid contamination of feed or food stuffs.

Environment: Do not contaminate domestic or irrigation water supplies, lakes, streams and ponds.

Resistance: Strains of fungi may develop tolerance to Senator PSPT or other products in the same class of chemistry after prolonged use. Use full label rates of Senator PSPT and ensure thorough coverage. When possible alternate with products from different chemical families. If potatoes are going to be treated with Mertect prior to storage in the fall, do not use Senator PSPT or any other product containing thiophanate-methyl at planting.

Do not use Senator PSPT or any other product containing thiophanate-methyl if Mertect was used as the storage treatment on the seed. Use a seed treatment with a different mode of action on Mertect treated seed potatoes. Implement an integrated disease management program that helps to maintain lower inoculum levels. This includes planting disease free tubers, crop rotation, and good hygiene in storage bins.

Precautions:

Keep out of reach of children. May be harmful if swallowed. Avoid inhalation of dust. Wash hands and face after handling.

Thiram 75WP

Fungicide Group – M
(Refer to page 225)

Company:

Bayer Crop Science PCP# 27556

Formulation:

75% thiram formulated as wettable powder. Container sizes: 5 kg, 25 kg.

Crops and Staging:

Mustard, grasses, dry beans, peas, soybeans, safflower, corn, and alfalfa. Seed treatment.

Diseases Controlled:

Seed decay, seedling blight, damping off.

Cost (2004 suggested retail prices):

Mustard, grasses, alfalfa: $3.59 per 25 kg seed
Dry bean, pea, soybean: $1.00 to $1.40 per 25 kg seed
Corn: $1.20 per 25 kg of seed
Safflower: $2.00 per 25 kg of seed

Rates:

CROP	KG THIRAM/ 25 KG SEED	AMOUNT OF SEED (KG) ONE 5 KG PACKAGE TREATS
Mustard, grasses, alfalfa	0.09	1,389
Dry beans, peas, soybeans	0.025 - 0.035	3,571 - 5,000
Field corn	0.03	4,166
Safflower	0.05	2,500

Compatibility with Rhizobia-based Inoculants:

Thiram is compatible with Rhizobia, however, some restrictions may apply in storage length with Thiram-treated seed inoculated with Rhizobia. Contact Rhizobia manufacturer on use patterns with their Rhizobia strain.

How it Works:

The active ingredient thiram is a dithiocarbamate fungicide with contact activity.

Restrictions:

Use: Treated seed should be labelled **"Do not use for food or feed. This seed has been treated with thiram"**

Storage: Store in a cool, dry, ventilated place, away from feeds and foods. Keep away from flame, sparks and heat.

Grazing: Do not graze treated areas or feed clippings from treated areas to livestock.

Precautions:

Do not consume alcohol 24 hours before or after working with thiram. Avoid breathing dust or spray mist. When treating, augering or handling treated seed, work in a well-ventilated area and wear a suitable dust mask, goggles and gloves.

Hazard Rating:

◈ Warning Poison

Vitaflo 280

Fungicide Group – 7, M
(Refer to page 225)

Company:

Bayer CropScience PCP# 11423

Formulation:

Vitaflo 280: 15.59% carbathiin and 13.25% thiram formulated as a liquid suspension.
Container sizes: 10 L, 200 L, 1000 L.

Crops, Rates and Diseases Controlled:

CROP	ML PER 25 KG SEED	DISEASES CONTROLLED	DISEASES SUPPRESSED
Barley	57 - 82[1]	False loose smut; True loose smut; Covered smut; Leaf stripe; Seed rot and seedling blight caused by *Cochliobolus sativus, Fusarium, Pythium, Aspergillus, Penicillium* and *Alternaria* spp.	Suppression of root rot caused by *Cochliobolus sativus* and *Fusarium* spp.; Suppression of net blotch
Wheat	57 - 82[1]	Loose smut; Stinking smut (bunt); Seed-borne dwarf bunt; Seed-borne *Septoria*; Seed rot and seedling blight caused by *Cochliobolus sativus, Fusarium, Pythium, Aspergillus, Penicillium* and *Alternaria* spp.	Suppression of root rot caused by *Cochliobolus sativus* and *Fusarium* spp.
Oats	82	Loose smut; Covered smut; Seed rot and seedling blight caused by *Cochliobolus sativus, Pythium, Aspergillus, Penicillium* and *Alternaria* spp.	
Rye	57 - 82[1]	Stem smut; Seed rot and seedling blight caused by *Cochliobolus sativus, Fusarium, Pythium, Aspergillus, Penicillium* and *Alternaria* spp.	Suppression of root rot caused by *Cochliobolus sativus* and *Fusarium* spp.
Triticale	50	Seed rot and seedling blight	
Dry beans	65[2]	Seed rot, seedling blight, and root rot caused by *Rhizoctonia solani*; Seed-borne anthracnose *(Colletotrichum lindemuthianum)*	
Corn	70 140 - 187[3]	Seed rot and seedling blight; Seed-borne head smut	
Flax, Linola	131	Seed rot, root rot and seedling blight caused by *Rhizoctonia solani* and *Fusarium* spp.	
Lentils	82	Seed rot, seedling blight, and early season root rot caused by *Botrytis cinerea, Rhizoctonia solani, Fusarium* and *Pythium* spp.	
Peas	65 - 82[4]	Seed rot and seedling blight caused by ascochyta (*Mycosphaerella*), *Rhizoctonia solani, Fusarium* and *Pythium* spp.	
Soybeans	65	Seed rot and seedling blight caused by *Rhizoctonia solani, Phomopsis* and *Fusarium* spp.	

(see footnotes on following page)

[1] Use the high rate to control true loose smut of barley, loose smut of wheat, and stem smut of rye. The high rate will provide a higher level of protection against seed rot and seedling blight.

[2] Vitaflo 280 will not protect against wind-borne anthracnose spores. This product will not control anthracnose if seed is severely infected.

[3] Use 140-187 mL per 25 kg seed to control head smut of corn.

[4] Use high rate on peas to control ascochyta (*Mycosphaerella pinodes*)

Cost (2004 suggested retail price):

Barley, wheat, rye: $1.35 to $1.94 per 25 kg seed
Triticale: $1.19 per 25 kg seed
Peas, lentils, oats: $1.94 per 25 kg seed
Flax: $3.10 per 25 kg seed
Beans, soybeans: $1.54 per 25 kg seed
Corn: $1.66 to $4.43 per 25 kg seed

Application:

Vitaflo 280 can be diluted with water for use in slurry treaters. Dilute Vitaflo 280 as required to apply the specified rate of undiluted Vitaflo 280. Specific directions regarding dilution are contained in the Vitaflo 280 Operating Manual. Agitate while diluting and continue agitating for approximately two minutes after diluting. Keep in constant agitation and use within 24 hours of diluting. DO NOT USE SLURRY TREATMENT FOR FLAX.

Vitaflo 280 should be used undiluted in all commercial seed treaters other than slurry treaters. Treater calibration should be rechecked if the Vitaflo 280 temperature falls below -18°C. Undiluted Vitaflo 280 can be used at temperatures down to -30°C.

How it Works:

The active ingredient carbathiin is carboximide fungicide with systemic activity and the active ingredient thiram is a dithiocarbamate fungicide with contact activity.

Tank Mixes:

Mixing of Vitaflo 280 with solvent based products may result in gelling or sedimentation. Seed treater lines and reservoirs should be thoroughly cleaned before use of Vitaflo 280. To clean, flush with water containing detergent.

Compatibility with Rhizobia-based Inoculants:

Vitaflo 280 is compatible with Rhizobia. Do not tank mix Vitaflo 280 and Rhizobia. Always check with Rhizobia manufacturers on any restrictions that may exist with seed treatments.

Restrictions:

Use: Seed treated with Vitaflo 280 must be labelled as follows **"This seed has been treated with Vitaflo 280 liquid seed protectant containing carbathiin and thiram. Do not use for feed, food, or oil processing. When augering or handling treated seed work in well-ventilated area and wear suitable mask, goggles and gloves."** Do not use treated seed for feed, food or oil processing.

Storage: Do not store Vitaflo 280 in direct sunlight or above 35°C. Vitaflo 280 will not freeze even at extreme temperatures. Do not store dry beans or soybeans treated with Vitaflo 280. Wheat, barley, rye, oats, triticale and flax seed treated with Vitaflo 280 can be stored up to 18 months without reduction in germination. Corn seed treated with Vitaflo 280 can be stored up to one year without reduction in germination.

Grazing: Do not graze or feed livestock on treated area for four weeks after planting.

Environment: Do not contaminate ponds, lakes or streams.

Precautions:

Work in a well ventilated area when treating seed or while augering or handling treated seed. Wear suitable dust mask, goggles and chemical resistant gloves. Keep out of reach of children. Do not consume alcoholic beverages 24 hours before or after treating seed with Vitaflo 280 or Vitavax Powder.

Hazard Rating:

◈ Warning Poison

Vitavax rs Fungicide

Fungicide Group – 7, M
(Refer to page 225)

This seed treatment is available to commercial seed treaters only.

Company:

Bayer CropScience PCP# 25862

Formulation:

5.97% carbathiin and 12.94% thiram as a liquid seed treatment. Container size: 10 L, 100 L

Crops and Staging:

Canola, rapeseed and mustard. Seed treatment.

Diseases Controlled:

Canola, rapeseed and mustard: Seed rot, seedling blight and damping off caused by species of *Rhizoctonia, Pythium* and *Alternaria*.

Canola and rapeseed: Seed-borne blackleg.

Cost:

Not available.

Rate:

208 to 312 mL per 25 kg of seed (0.833 to 1.25 L per 100 kg of seed).

100 L will treat 8000 to 12,000 kg of seed

Application Information:

Prior to and during application, Vitavax rs Fungicide must be thoroughly agitated to ensure uniform mixing of product. Vitavax rs Fungicide treated seed does not require drying after treatment. Refer to label for rinsate instructions.

How it Works:

The active ingredient carbathiin is a carboximide fungicide with systemic activity and the active ingredient thiram is a dithiocarbamate fungicide with contact activity.

Registered Mixes:

Can be co-applied on the seed with Apron FL (Allegiance FL) and Gaucho 480 FL.

Restrictions:

Use: Treated seed must be labelled **"This seed has been treated with Vitavax rs Fungicide, containing carbathiin and thiram, do not use for food, feed or oil processing."** Do not reuse container for any purpose. Augers used for handling treated seed should not be used to move seed for feed, food or oil processing. Do not reuse bags from treated seed to handle food or feed products. Do not contaminate feed or food-stuffs with treated seed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Treated canola or rapeseed stored for periods more than 9 months should be tested for germination before planting. Storage of Vitavax rs Fungicide at low temperatures is not recommended. Keep product above 10°C prior to and during application. Do not store treated seed above 25°C or in direct sunlight.

Grazing: Do not feed or graze livestock on treated areas for four weeks after planting.

Environment: Do not contaminate ponds, lakes or streams.

Precautions:

Work in a well ventilated area when treating seed or while augering or handling treated seed. Wear chemical resistant gloves when handling treated seed and goggles to protect eyes from possible splashing. Keep out of reach of children. Because of possible unpleasant side effects, alcoholic beverages should be avoided for 24 hours before or after treating with Vitavax rs Fungicide.

Hazard Rating:

☠ Caution Poison.

Caution – Eye irritant.

Vitavax Single Solution

Fungicide Group – 7
(Refer to page 225)

Company:

Bayer CropScience PCP# 14069

Formulation:

230 g/L carbathiin formulated as a solution.
Container sizes: 10 L, 200 L.

Crops and Staging:

Barley, wheat, oats, rye, flax. Seed treatment.

Cost (2004 suggested retail prices):

Barley, wheat: $1.53 to $1.90 per 25 kg seed
Oats, rye: $1.53 per 25 kg seed
Flax: $2.55 per 25 kg seed

Diseases Controlled:

BARLEY	WHEAT	OATS	RYE	FLAX
Covered smut True loose smut False loose smut Seed rot Seedling blight *Suppression of net blotch, leaf stripe, common root rot	Loose smut Stinking smut (bunt) Seed rot Seedling blight *Suppression of common root rot	Covered smut Loose smut Seed rot Seedling blight *Suppression of common root rot	Stem smut Seed rot Seedling blight *Suppression of common root rot	Seed rot Seedling blight

* will not protect plants from infection beyond the first leaf stage

Rates:

CROP	ML/25 KG SEED	KG SEED TREATED/ 10 L CONTAINER
Wheat**	60 - 75	4,167 - 3,333
Barley**	60 - 75	4,167 - 3,333
Oats	60	4,167
Rye	60	4,167
Flax	100	2,500

** For wheat and barley varieties highly susceptible to true loose smut and for high levels of smut or bunt on seed, the 75 mL rate will give increased disease control.

Application Information:

Vitavax Single Solution is a ready to use formulation designed for commercial treaters and on-farm auger treating. Do not dilute with water. It is important that the seed and chemical are mixed quickly and uniformly. See instructions supplied with the applicable treater system.

How it Works:

The active ingredient carbathiin is a carboximide fungicide with systemic activity.

Restrictions:

Use: Treated seed must be labelled as followed: **"This seed has been treated with Vitavax Single Solution Seed Protectant. Contains carbathiin; do not use for food, feed or oil processing."** Do not contaminate feed or foodstuffs.

Storage: Do not store below 0°C. Product is flammable. Keep away from flame and sparks.

Grazing: Do not graze or feed livestock on treated areas for four weeks after planting.

Environment: Do not contaminate ponds, lakes or streams.

Precautions:

Keep out of reach of children. Work in a well-ventilated area when treating or handling treated seed. Wear a suitable mask, gloves and goggles. Product may damage painted surfaces by removing paint or leaving stain. Do not use rubber or PVC hose.

Insect Control

Additional Resources

For information on insect control on livestock or livestock housing facilities see:

http://www.agr.gov.sk.ca/apps/insectPest/default.asp

For information on insect management in commodities other than livestock or those covered in this guide see the WCCP Guide to Integrated Control of Plant Pests at: http://www.members.shaw.ca/dan-johnson/wccpabout.html

Insect Control Decisions

Control of insects should be considered when numbers or damage exceed economic thresholds. To select an insecticide, verify the registered products for the insect and field crop in the following tables. Consideration should then be given to the rates, cost, preharvest intervals, the product application, restrictions, precautions and the hazard rating. The **preharvest interval** is the number of days that must pass between the last application of a pesticide and harvest. Harvest is the cutting of the crop or removal of the produce from the plant.

It includes direct-combining, cutting (swathing) or grazing; it does not include swath-combining or baling for hay.

Field Scouting

Field scouting is the regular examination of fields to accurately assess the kind and the number of insects present and the amount of damage being done. Scouting should be done weekly during the growing season and daily when infestations approach economic levels or when weather conditions favour the rapid development of specific pests.

To properly scout for insect pests, you must know when they occur, where they live, what they look like, and how to find and count them. Generally, in fields of less than 100 acres, check a minimum of five locations. In fields greater than 100 acres, check a minimum of 10 locations. There are several possible scouting patterns that can be used when checking fields. These options are based on pest distribution and field configuration.

Pattern 1: Used when pests are uniformly distributed.

This scouting pattern typically looks like an X, Z or W, excluding field edges. Pests that fit this pattern include aphids, bertha armyworm, diamondback moth, lygus bugs.



Pattern 2: Used when pests are at the edges of fields.

Scout by walking along field edges, fence lines or ditches.

Some examples of pests appearing at the edges of fields include flea beetles, colorado potato beetles and grasshoppers.



Monitoring methods, typical symptoms and **economic thresholds** for the more common crop pests are described in the field scouting section for each commodity. The **economic threshold** is the minimum number of insects or damage to the crop that should result in management action. Note that factors such as moisture, temperature conditions and stage of crop growth, can increase or decrease the impact of insects on crop production.

Estimating Percent Defoliation

Many action-thresholds for insects are based on percent defoliation of the plants they are feeding on. The following figure may assist in determining the percent defoliation. Although the photo below is of sunflower leaves, this figure can be used to estimate % defoliation for many crops.



Photo courtesy of North Dakota State University Extension Service

Reducing Bee Losses from Insecticides

Careless use of insecticides can kill bees and other beneficial insects such as pollinators, predatory and parasitic biological control insects. Help to reduce insecticide poisoning of bees by:

1. **Avoid applying insecticides which are toxic to bees on crops in bloom.** Any field with even a small amount of bloom, whether it is the main crop, cover crop, or weeds will probably have foraging bees visiting the flowers. If at all possible, apply insecticides before or after the crop has gone into bloom. Control all flowering weeds prior to insecticide application.

2. **Apply insecticides when bees are least active.** The highest level of bee activity occurs during the day. Apply insecticides in late evening or early morning when the bees are not foraging. As a general rule, evening applications are less hazardous to bees than morning applications.
3. Avoid insecticide drift. To avoid insecticides drifting into non-target locations, do not apply insecticides during windy conditions. Choose nozzles with a low drift rating. As a general rule, ground applications of insecticide are less prone to drift than aerial applications.
4. Contact the beekeeper before spraying. Communication and cooperation between the insecticide applicator and the beekeeper can usually prevent bee losses. Notifying the beekeeper in advance (i.e. 48 hours) of applying insecticides will allow the beekeeper to move or protect the colonies from insecticide damage.
5. If possible, use insecticides and/or insecticide formulations which are the least hazardous to bees. The following table "Field Hazard of Insecticides to Bees" will help in selecting the least hazardous insecticide. In general, dusts are more hazardous to bees than sprays. Wettable powders are more hazardous than emulsifiable concentrates (EC) or water-soluble formulations. Granular insecticides and spreadable bran bait insecticides are generally the least hazardous to bees.

Field Hazard of Insecticides to Bees

The following table indicates the maximum time required for the insecticides listed in the Guide to be degraded by weather to a low hazard level for bees. **These times are to be used as general guidelines only.** Most of these insecticides have not been tested for bee toxicity under Western Canadian conditions and environmental conditions influence the rate at which pesticides degrade.

INSECTICIDE	TOXICITY RATING		RESIDUE HAZARD (DAYS)[a]
	HONEY BEE	LEAFCUTTER BEE	
Admire	1	1	–[b]
Cygon/Lagon	1	1	10
Decis	2	2	1
Diazinon	–[b]	–[b]	–[b]
Dibrom	1	1	–[b]
Eco bran	3	3	N/A
Furadan	1	1	5
Guthion/Sniper	1	1	5
Lannate	1	1	1.5
Lorsban/Nufos/Pyrinex/ Chlorpyrifos	1	1	3.5
Malathion	2	1	2 (Honey Bee), 6 (Leafcutter Bee)
Matador	2	2	1
Monitor	1	1	1
Pounce	1	1	5
Ripcord	–[b]	–[b]	–[b]
Sevin	1	1	7
Thiodan/Thionex	2	1	2

TOXICITY RATING 1 = High 2 = Moderate 3 = Low

[a] Residue hazard represents the average time in days that residues poisonous to bees will remain on foliage. Most of this information has not been generated under Western Canadian growing conditions and should be used only as a guideline.

[b] Information currently not available, although these products are known to be toxic to bees.

Field Scouting in Alfalfa

SAP FEEDERS

Lygus bug/Alfalfa plant bug

Typical Damage: Field blooms poorly or not at all. Flower buds blasted, whitish, and dry; flowers dropping off before fully open. Collapsed seed.

When and How to Monitor: Monitor June through mid-August. Make five 180° sweeps with a 15-inch (40 cm) insect net through alfalfa canopy at each sampling site. Record total number of plant and lygus bugs (both nymphs and adults) captured. Calculate average number per sweep.

Economic Threshold: Hay: Control not recommended. Seed alfalfa at bud and early bloom: 8 lygus bugs/sweep; 4 alfalfa plant bugs/sweep; or 5 bugs if the plant bug population is a combination of lygus bugs and alfalfa plant bug. If insecticides are used, attempt to spray before the onset of bloom. Protecting insect pollinators in seed production fields is very important.

Pea Aphid

Typical Damage: Suck juices from plants; stunt growth; cause premature drying.

When and How to Monitor: Monitor July through August. Take 5 sweeps at each location. Monitor fields closely during periods of slow plant growth.

Economic Threshold: 100-200 aphids/180°sweep when crop is moisture stressed, or until mid-August.

DEFOLIATORS

Alfalfa Weevil

Typical Damage: Feed on developing buds and leaves. Stunt growth.

Economic Threshold: Alfalfa Hay: One of the best control strategies is to cut fields for hay early. If early cutting of the hay crop is not possible, treatment should be considered when 25-50% of leaves on the upper one-third of the stem show damage or 50-70% of terminals show injury. Alfalfa Seed: 20-25 larvae/sweep (90°=straight sweep) or 35-50% of foliage tips showing damage.

LEAFMINERS

Alfalfa Blotch Leafminer

Typical Damage: Females feed by puncturing leaves with their ovipositors, creating characteristic "pinholes," and consuming plant juices. A single female creates an average of 3,769 pinholes during her lifetime. Larvae create distinctive mines as they feed. Infested fields take on a whitish cast due to the larval mines in the leaves.

Economic Threshold: Treatment is suggested if 30-40% of the plants exhibit pinhole feeding injury. If insecticides are used, they must be applied during the "pinhole" stage.

Alfalfa Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Sap Feeders						
Lygus bugs	Matador (P)	0.034 L	$4.97	Do not apply within 14 days of livestock foraging.	A or G	64-110
	Decis 5EC (seed production only) (P)	0.08-0.1 L	$7.36-$9.20	20	G	395
	Malathion 500 (OP)	0.80-1.21 L	$10.72-$16.21	7	A or G	4302
	Malathion 500E (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Dibrom (OP)	0.44-0.88	$16.15-$32.30	4	A or G	345
	Cygon (alfalfa) (OP)	0.17 L	$5.19	2	A or G	450
	Cygon 480 (alfalfa seed production) (OP)	0.44 L	$10.63	28	A or G	450
	Lagon (seed production only) (OP)	0.44 L	$12.45	28	A or G	60

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Guthion SC Guthion 50/ Sniper (OP)	0.91-1.41 L 0.45-0.70 L	\$14.74-\$22.84 \$12.29-\$19.11	21	A or G	21-37 12.3-24.7
Alfalfa plant bug	Cygon 4-E (alfalfa) (OP)	0.17 L	\$5.19	2	A or G	450
	Cygon 480 (alfalfa seed production) (OP)	0.44 L	\$10.63	28	A or G	450
	Lagon (seed production only) (OP)	0.44 L	\$12.45	28	A or G	60
	Guthion SC Guthion 50/ Sniper (OP)	0.91-.141 L 0.45-0.70 L	\$14.74-\$22.84 \$12.29-\$19.11	21	A or G	21-37 12.3-24.7
Potato leafhopper	Matador (P)	0.034 L	\$4.97	Do not apply within 14 days of livestock foraging.	A or G	64-110
	Sevin (C)	1.01-1.6 L	\$15.00-\$23.76	2	A or G	649
	Malathion 500 (OP)	0.80-1.21 L	\$10.72-\$16.21	7	A or G	4302
	Malathion 500E (OP)	0.91-1.11 L	\$11.24-\$13.71	7	A or G	1375-2800
	Dibrom (OP)	0.44-0.88 L	\$16.15-\$32.30	4	A or G	345
	Lagon (OP)	0.17 L	\$4.81	2	A or G	480
	Cygon (OP)	0.17 L	\$5.19	2	A or G	450
	Guthion SC Guthion 50/ Sniper (OP)	0.91-1.41 L 0.45-0.70 L	\$14.74-\$22.84 \$12.29-\$19.11	21	A or G	21-37 12.3-24.7
	Methoxychlor (OC)	2.02-2.53 L	\$21.05-\$26.36	7	G	5,000-7,000
Pea Aphid	Matador (P)	0.034 L	\$4.97	Do not apply within 14 days of livestock foraging.	A or G	64-110
	Malathion 500 (OP)	0.80-1.21 L	\$10.72-\$16.21	7	A or G	4302
	Malathion 500E (OP)	0.91-1.11 L	\$11.24-\$13.71	7	A or G	1375-2800
	Dibrom (OP)	0.44-0.88 L	\$16.15-\$32.30	4	A or G	345
	Cygon (OP)	0.17 L	\$5.19	2	A or G	450
	Lagon (OP)	0.17 L	\$4.81	2	A or G	60
Spider mites	Malathion 500E (OP)	0.91-1.11 L	\$11.24-\$13.71	7	A or G	1375-2800
Defoliators						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	\$3.00-\$6.00	2	G	NA

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Sprays					
	Matador (young grasshoppers only) (P)	0.025-0.034 L (Ground) 0.034 L (Aerial)	$3.71-$4.85 $4.97	Do not apply within 14 days of livestock foraging.	A or G	64-110
	Sevin (C)	0.50-1.01 L	$7.43-$15.00	2	A or G	649
	Malathion 500 (young grass-hoppers) (OP)	0.80-1.21 L	$10.72-$16.21	7	A or G	4302
	Malathion 500E (young grass-hoppers) (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Cygon (OP)	0.17-0.34 L	$5.19-$10.37	2-7	A or G	450
	Lagon (OP)	0.22-0.36 L	$6.23-$10.19	2-7	A or G	60
	Guthion SC Guthion 50 (border application only) (OP)	0.44-0.70 L 0.21-0.34 L	$7.13-$11.34 $5.74-$9.29	21	A or G	21-37 12.3-24.7/7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5000-7000
Alfalfa weevil	If alfalfa has reached the bud or early bloom stage, immediate **cutting** will kill many alfalfa weevil larvae					
	Matador (P)	0.034 L	$4.97	Do not apply within 14 days of livestock foraging.	A or G	64-110
	Decis 5EC (seed crops only) (P)	0.08-0.1 L	$7.36-$9.20	20	G	395
	Malathion 500 (OP)	0.80-1.21 L	$10.72-$16.21	7	A or G	4302
	Malathion 500E (larvae only) (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Cygon (OP)	0.17 L	$5.19	2	A or G	450
	Guthion SC Guthion 50/ Sniper (OP)	0.91-1.41 L 0.45-0.70 L	$14.74-$22.84 $12.29-$19.11	21	A or G	21-37 12.3-24.7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5,000-7,000
Sweet clover weevil	Cygon 4E (OP)	0.34-0.44 L	$10.37-$13.42	28	A or G	450
	Guthion SC Guthion 50/ Sniper (OP)	0.91 L 0.45 L	$14.74 $12.29	21	A or G	21-37 12.3-24.7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5,000-7,000
Blister beetle	Sevin (C)	1.01-1.6 L	$15.00-$23.76	2	A or G	649
True Armyworm	Sevin (C)	1.01-2.12 L	$15.00-$31.48	2	A or G	649
	Methoxychlor (OC)	2.83-4.55 L	$29.49-$47.41	7	G	5,000-7,000
Beet webworm	Sevin (C)	1.01-2.12 L	$15.00- $31.48	2	A or G	649
Alfalfa looper	Dibrom (OP)	0.44-0.88 L	$16.15-$32.30	4	A or G	345

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Alfalfa Caterpillar	Sevin (C)	1.01-2.12L	$15.00-$31.48	2	A or G	649
Leafminers						
Alfalfa blotch leafminer	Sevin (C)	1.01-2.12 L	$15.00-$31.48	2	A or G	649
	Malathion 500E (OP)	1.11 L	$13.71	7	A or G	1375-2800
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Cygon 480 (OP)	0.22 L	$6.71	2	A or G	450
	Lagon (OP)	0.22 L	$6.23	2	A or G	60

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Barley – See Small Grain Cereals

Scouting for insects in Beans
(Dry Edible Beans)

BELOWGROUND FEEDERS

Seedcorn Maggot

Typical Damage: Seedcorn maggot attacks bean seed, preventing sprouting or weakening seedlings. The yellowish white maggot is found burrowing in the seeds or emerging stem. Seedcorn maggots are usually most severe in wet, cold seasons and on high organic matter soils.

SAP FEEDERS

Leafhoppers

Typical Damage: Foliage becomes dwarfed, crinkled, and curled. Small triangular brown areas appear at the tips of leaves, gradually spreading around the entire leaf margin.

Economic Threshold: An average of one leafhopper per trifoliate leaf. Do not let infestations and damage progress to the point that yellowing of foliage is easily detected.

DEFOLIATORS

Grasshoppers

Economic Threshold: Substantial yield loss does not occur until up to 35% defoliation occurs before bloom and 15% after bloom.

Beans (Dry Edible) Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders						
Wireworms	No insecticides registered					
Seedcorn Maggot	Sow seeds as shallow as possible in a warm, well-prepared seedbed. If manure is used, apply and plow it under the previous fall.					
	Agrox CD, Agrox B-2, DCT (OP)	See Seed Treatments section of Guide to Crop Protection. For control of Seedcorn maggots only			Seed treatment	N/A
	Diazinon 50W	20 g/300 mL water/4 L seed	–	–	Seed treatment	1960
Sap Feeders						
Lygus Bugs	Sevin XLR Plus (C)	2.12-2.59 L	$31.48-$38.46	5	A or G	649
	Cygon (OP)	0.28-0.40 L	$8.54-$12.20	7	A or G	450
Potato Leafhoppers	Sevin XLR Plus (C)	1.01 L	$15.00	5	A or G	649
	Malathion 500 (OP)	0.56-1.21 L	$7.50-$16.21	1	A or G	4302
	Malathion 500E (OP)	0.61-1.11 L	$7.53-$13.71	3	A or G	1375-2800
	Diazinon 50W (OP)	0.40 kg	$10.24	7	G	1960
	Diazinon 50 EC (OP)	0.445 L	$7.28	7	G	NA
	Cygon (OP)	0.28-0.40 L	$8.54-$12.20	7	A or G	450
	Thionex EC (OC)	0.6-1.0 L	$9.75-$16.25	2	A or G	110
	Thiodan (OC)	0.6-1.0 L	$9.75-$16.25	2	A or G	107.2
	Methoxychlor (OC)	1.21-3.64 L	$12.61-$37.93	7	G	5000-7000
Aphids	Malathion 500 (OP)	0.56-1.21 L	$7.50-$16.21	1	A or G	4302
	Malathion 500E (OP)	0.61-1.11 L	$7.53-$13.71	3	A or G	1375-2800
	Diazinon 50W (OP)	0.40 kg	$10.24	7	G	1960
	Diazinon 500 E (OP)	0.445 L	$7.52	3	G	1500
	Diazinon 50 EC (OP)	0.445 L	$7.28	7	G	NA
	Dibrom (Dry, Field, Lima) (OP)	0.445-0.890 L	$16.15-$32.30	4	A or G	345
	Cygon (OP)	0.28-0.40 L	$8.54-$12.20	7	A or G	450
	Thionex EC (OC)	0.6-1.0 L	$9.75-$16.25	2	A or G	110
	Thiodan (OC)	0.6-1.0 L	$9.75-$16.25	2	A or G	107.2
Defoliators						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	$3.00-$6.00	5	G	N/A
Variegated Cutworm	Sevin XLR Plus (C)	30-35 mL/100 m of row		5	A or G	649
Alfalfa looper	Dibrom (OP)	0.445-0.890 L	$16.15-$32.30	4	A or G	345

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Scouting for insects in Canaryseed

APHIDS

When and How to Monitor: Start checking for aphids during the early heading stage of canaryseed. The head should be bent and closely inspected for aphids hiding along the small stem inside the canaryseed head. Also check the stems, underside of leaves, and in the canaryseed boot.

Economic Threshold: 10 to 20 aphids on 50% of the stems prior to the soft dough stage.

Canaryseed Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
Sap feeders						
Aphids	Cygon 480 (OP)	0.20 L	$6.10	21	A or G	450
	Lagon 480E (OP)	0.20 L	$5.66	21	A or G	60
	Malathion 500E (canary grass- for seed) (OP)	0.56 L	$6.92	14	A or G	1375-2800

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: OP=organophosphates

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Field Scouting in Canola

Scouting Calendar

Early-season: Flea beetles, cutworms, red turnip beetle, diamondback moth

Mid-season: Diamondback moth, lygus bug, cabbage seedpod weevil, grasshoppers

Late season: Bertha armyworm, diamondback moth, Lygus bugs, grasshoppers, alfalfa looper

Cutworms

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in field.

When and How to Monitor: Monitor late May to mid-July.

Economic Threshold: none, but use 25-30% stand reduction.

SAP FEEDERS:

Lygus bugs

Typical Damage: Attacked buds appear shrunken and bleached white. Damaged seeds appear dark brown and shriveled.

When and How to Monitor: Monitor from the end of flowering until seeds within the pod have become firm. Make 10 180° sweeps with a 37 cm diameter insect net at each sampling site. Sample canola for lygus bugs on a sunny day when the temperature is above 20°C and the crop canopy is dry.

Economic Threshold: 10-18 lygus bugs/10 sweeps from the end of flowering to early pod development in the upper canopy and 15-25 lygus bugs/10 seeps in early pod ripening stage. Controls are not recommended when seeds are ripening (yellow or brown).

Aphids

Economic Threshold: Control aphids in canola if densities exceed 25 aphids/10 cm shoot tip after flowering.

DEFOLIATORS:

Flea beetles

Typical Damage: Shot-holes in leaves complete destruction of seedling plants late May through June. Holes chewed in pods in August (occasional).

When and How to Monitor: Monitor May through June when crop is in seedling stage. Examine 10 plants at random at each stop. Estimate overall percentage leaf loss.

Economic Threshold: When 25 percent of leaf surface is destroyed and flea beetles are present. If damage is only along the field margins and beetles are still congregated there, then control measures should be applied to the damaged areas only.

Diamondback moth

Typical Damage: Flowers clipped or chewed, outer layers of stem and pods chewed, holes chewed in pods.

When and How to Monitor: Monitor late-May through early September. Beat plants within 50 cm x 50 cm area and count larvae on the ground. Multiply by 4 to get the number of larvae per square metre. Do this in at least 5 areas of the field.

Economic Threshold: 100 to 150 larvae/m^2 in immature to flowering plants. 200 to 300 larvae/m^2 in plants with flowers and pods. Note that these threshold numbers are based on stands averaging 150-200 plants/m^2. In areas where stands are thinner, the economic threshold should be lowered accordingly. A nominal threshold of 25-33% defoliation with larvae still present can be applied for canola at seedling stage.

Bertha Armyworm

Typical Damage: Outer layers of stems and pods chewed resulting in whitish appearance, holes chewed in pods.

When and How to Monitor: Monitor larvae in late July through early August. Sample at least 3 locations a minimum of 50 metres apart. At each stop, shake plants in a 1/4 m^2 area and carefully check soil surface for dislodged larvae. During heat of the day larvae will often be found under leaves on soil surface. Adult populations can be monitored in June and July using traps baited with a pheromone, which attracts the male moths. This determines the risk of a larval infestation occurring latter.

Economic Threshold: 10 to 34 larvae/yard^2 (m^2), depending on value of crop and cost of control.

A loss of 0.058 Bu/acre for each larva/m^2 can be expected. Yield loss may be greater for canola under moisture stress.

Canola Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Cabbage root maggot	• **Increased seeding rates and increased row spacing** (to about 25 to 30 cm) can reduce damage to the roots by root maggots. • **Cultivating** prior to seeding reduces adult emergence from overwintered pupae. Root maggot infestations are greater under zero-till systems than under conventional tillage, but yields under zero tillage usually still exceed those with conventional tillage.					
Cutworms (Army, dark-sided, redbacked, pale western)	Pounce (P)	0.073-0.158	$7.30-$15.00	Treat prior to 6-leaf stage	G	1276
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486 L	$6.06-$8.92	21	A or G	205-418
Sap Feeders						
Lygus Bugs	Decis 5EC (P)	0.06 L	$5.52	7	A or G	395
	Matador (P)	0.034 L	$4.97	50	A or G	64-110
	Lorsban/Nufos/ Pyrinex/ Chlorpyrifos 480 EC (OP)	0.405 L	$7.27	21	A or G	205-418
	Gaucho Platinum (suppression only) (N)	1.667L/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	N/A

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
Turnip aphid	Cygon 480 (OP)	0.34-0.36 L	$10.37-$10.98	21	A or G	450
	Lagon (OP)	0.34-0.36 L	$9.62-$10.19	21	A or G	60
	Gaucho Platinum (suppression only) (N)	1.667L/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	N/A
Aster leafhopper	Cygon 480 (OP)	0.34-0.36 L	$10.37-$10.98	21	A or G	450
	Lagon (OP)	0.34-0.36 L	$9.62-$10.19	21	A or G	60
Defoliators						
Crucifer Flea beetle and/or striped flea beetle	**Seed Treatments**					
	Helix (N)	1500 ml/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	>5,000
	Helix Xtra (N)	1500 ml / 100 kg of seed	N/A	Seed Treatment	Seed Treatment	>5,000
	Gaucho Canola System,(N)	0.833 L/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	N/A
	Gaucho Platinum (N)	1.667L/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	N/A
	Prosper (N)	—	N/A	Seed Treatment	Seed Treatment	N/A
	Foliar Sprays					
	Decis Flowable (P)	0.04-0.06 L	$4.65-$6.98	7	A or G	15000
	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Ripcord (P)	0.02 L	$2.40	30	G	242-542
	Matador (P)	0.034 L	$4.97	50	A or G	64-110
	Sevin XLR Plus (C)	0.2 L	$2.97	Seedling application only	A or G	649
	Furadan (C)	0.06 L	$2.18	60	A or G	7.34
	Malathion 500 (OP)	0.44 L	$5.90	7	A or G	4302
	Malathion 500E (OP)	0.45-0.71 L	$5.56-$8.77	7	A or G	1375-2800
Red Turnip beetle	Furadan (C)	0.11 L	$3.99	60	A or G	7.34
Cabbage seedpod weevil	Gaucho Platinum (suppression of larvae only) (N)	1.667L/ 100 kg of seed	N/A	Seed Treatment	Seed Treatment	N/A
	Matador (adults) (P)	0.034 L	$4.97	50	A or G	64-110
	Decis 5EC (for control of adults only) (P)	0.08	$7.36	7	A or G	395
Diamondback moth	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Malathion 500 (OP)	0.22-0.34 L	$2.95-$4.56	7	A or G	4302
	Malathion 500E (larvae only) (OP)	0.22-0.34 L	$2.72-$4.20	7	A or G	1375-2800

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
	Lorsban/Nufos/ Pyrinex/ Chlorpyrifos 480 EC (for larval control only) (OP)	0.405 L	$7.27	21	A or G	205-418
Bertha Armyworm	**Seeding as early as possible** and choosing **early maturing varieties of canola** may help minimize damage in years when outbreaks are forecasted.					
	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Ripcord (P)	0.028 L (ground) 0.036 L (air)	$3.36 (ground) -$4.32 (air)	30	A or G	242-542
	Lannate (C)	0.0874-0.2064 kg	$6.48-$15.31	8	A or G	30-34
	Lorsban/Nufos/ Pyrinex/ Chlorpyrifos 480 EC (OP)	0.304-0.405 L	$5.21-$7.43	21	A or G	205-418
	Monitor (OP)	0.23-0.5 L	$6.61-$14.38	10	A or G	17
Alfalfa looper	Lannate (OP)	0.087-0.206 kg	$6.48-$15.31	8	A or G	30-34
	Lorsban/Nufos/ Pyrinex (OP)	0.304-0.405 L	$5.21-$7.43	21	A or G	272/205/418
Beet webworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Lannate (C)	0.0874-0.2064 kg	$6.48-$15.31	8	A or G	30-34
Clover cutworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Lannate (C)	0.0874-0.2064 kg	$6.48-$15.31	8	A or G	30-34
True armyworm	Lorsban/Nufos/ Pyrinex/ Chlorpyrifos 480 EC (OP)	0.304-0.405 L	$5.21-$7.43	21	A or G	205-418
Variegated cutworm	Lorsban/Nufos/ Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486	$6.06-$8.92	21	A or G	205-418
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	$3.00-$6.00	Treat only seedlings	G	N/A
	Sprays					
	Matador (young grasshoppers only) (P)	0.025-0.034 L (Ground) 0.034 L (Aerial)	$3.80-$4.97 $4.97	50	A or G	64-110
	Ripcord (young grasshoppers only) (P)	0.02-0.028 L	$2.40-$3.36	30	G	242-542
	Malathion 500 (OP)	0.45-0.69 L	$5.90-$9.11	7	A or G	4302
	Malathion 500E (OP)	0.45-0.71 L	$5.56-$8.77	7	A or G	1375-2800

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.235-0.354 L	$4.03-$6.50	21	A or G	205-418
	Cygon 480 (OP)	0.34-0.36 L	$10.37-$10.98	21	A or G	450
	Lagon (OP)	0.34-0.36 L	$9.62-$10.19	21	A or G	60
	Monitor (OP)	0.5 L	$14.38	10	A or G	17

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: N=Neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in Corn

Cutworms

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in filed.

When and How to Monitor: At each stop, examine 100 plants in a row. Calculate percentage of plants cut off or showing leaf feeding.

Economic Threshold: When 3% of plants are cut.

European corn borer

Typical Damage: Shot-holes in leaves, holes in stalk, tassels and ears.

When and How to Monitor: Field scouting should begin in early July. Older fields should be scouted first. At 10 locations, examine 10 plants for egg masses and signs of shot-holing. Calculate percentage of plants with shot-holes. Scout every 5 to 7 days until the end of July or larvae start to tunnel into the stalks.

Economic Threshold: When 50% of the plants have shot-holes in the laves, caused by young corn borers.

Corn Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Cutworms	Matador (P)	0.034 L	$4.97	1 (sweet corn) 14 (silage)	A or G	64-110
	Ripcord (black, white, darksided, redbacked, army, pale western) (P)	0.07 L	$8.40	21	G	242-542
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (dark sided, black, redbacked) (OP)	0.971 L (Pre-plant treatment), 0.486-0.971 L (seedling treatment)	Pre-plant treatment- $17.43; Seedling treatment- $8.72-$17.43	70	G	205-418
Wireworms	Poncho 250	0.25 mg of Poncho 600 per kernel	—	Seed Treatment	Seed Treatment	500-1,000

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Seed corn maggot	Agrox CD (OP)	0.05 kg/25 kg of seed		Seed Treatment	Seed Treatment	N/A
	Agrox B-2 (OP)	See label		Seed Treatment	Seed Treatment	N/A
	Diazinon 50W (OP)	0.020 kg/ 300 mL water/ 4L of seed	$0.40/4 L of seed	Seed Treatment	G	1960
Sap Feeders						
Corn leaf aphid	Lannate (sweet corn only) (C)	0.430-0.620 kg	$31.88-$45.99	5	G	30-34
	Thiodan/ Thionex EC (OC)	1.11 L	$18.04	50	G	110
Defoliators and Borers						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	$3.00-$6.00	1	G	N/A
	Sprays					
	Sevin XLR Plus (C)	0.50-1.01 L	$7.43-$15.00	1	A or G	649
European Corn Borer	Cultivars of **Bt corn** are resistant to feeding by European corn borer. If using cultivars of Bt corn, a refuge of at least 20% of the corn acres planted to non-Bt cultivars should be planted to reduce the odds of European corn borer developing resistance to Bt corn.					
	Success (sweet corn only) (S)	0.034	$15.64	7 (28 days for fodder harvest)	G	>5,000
	Dipel 2X DF (M)	0.23-0.45 kg	$20.24-$39.60	0	G	>4,000
	Matador (P)	0.034 L	$4.97	1 (sweet corn), 14 (silage)	A or G	64-110
	Decis 5EC (field corn) (P)	0.1-0.12 L	$9.20-$11.04	See product label	G only	395
	Ripcord (P)	0.07 L	$8.40	5	A or G	242-542
	Pounce (sweet corn) (P)	0.11-0.15 L	$11.00-$15.00	1	A or G	1030
	Sevin XLR Plus (C)	1.01-1.6 L	$15.00-$23.76	1	A or G	649
	Furadan (C)	0.445 L	$16.13	7	A or G, see guide	7.34
	Lannate (sweet corn only) (C)	0.253 kg	$18.77	3	See product label	30-34
	Malathion 500E (OP)	0.91-1.11 L	$11.24-$13.71	5	A or G	1375-2800
Corn Earworm	Matador (P)	0.034 L	$4.97	1 (sweet corn), 14 (silage)	A or G	64-110
	Ripcord (P)	0.07 L	$8.40	5	A or G, see product label	242-542
	Pounce (sweet corn) (P)	0.11-0.15 L	$11.00-$15.00	1	A or G	1030
	Sevin XLR Plus (C)	1.01-1.6 L	$15.00–$23.76	1	A or G	649
	Lannate (sweet corn only) (C)	0.174-0.253 kg	$12.91-$18.77	3	See product label	30-34
	Malathion 500E (OP)	0.91-1.11 L	$11.24-$13.71	5	A or G	1375-2800

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Thionex EC (OC)	1.11-1.62 L	\$18.04-\$26.33	50	G	110
	Thiodan (OC)	1.11-1.72 L	\$18.04-\$27.95	50	G	N/A
True Armyworm	Methoxychlor 240 (OC)	2.83-4.55 L	\$29.49-\$47.41	7	G	5,000-7,000
Fall armyworm	Sevin XLR (C)	1.01-1.6 L	\$15.00-\$23.76	1	A or G	649
	Matador (P)	0.034 L	\$4.97	1 (sweet corn) 14 (silage)	A or G	64-110
	Pounce (P)	0.073 L	\$7.30	1	G	1030

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: M=microbials, S=spinosyns, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in Flax

Cutworms

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in field.

When and How to Monitor: Monitor late May to mid-July. Sample 5 to 10 locations in the field. At each location, check several 0.25 m^2 (50cm x 50cm) areas. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m^2.

Economic Threshold: 4-5 larvae/m^2

Aphids

Typical Damage: Wilting of plants, plants are sticky.

When and How to Monitor: Monitor mid-July to mid-August. Check 10 plants at each stop. Record total number of aphids and calculate average per plant.

Economic Threshold: Three or more aphids per main stem at full flower or 8 or more aphids per main stem at green boll stage.

Beet webworm

Economic Threshold: >10 larvae/m^2

Flax Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Wireworm	No insecticides registered for the control of wireworms in flax					
Cutworms (Army, dark-sided, redbacked, pale western)	Decis 5EC (P)	0.08 L	$7.36	40	A or G	395
	Pounce (P)	0.073-0.158 L	$7.30-$15.00	Treat prior to 6 leaf stage	G	1030
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486 L	$6.06-$8.92	21	A or G	205-418
Sap Feeders						
Potato Aphid	Cygon 480 (OP)	0.18 L	$5.49	21	A or G	450
	Lagon (OP)	0.18 L	$5.09	21	A or G	60
Defoliators						
Grasshoppers	Decis Flowable (P)	0.032-0.048 L	$3.72-$5.58	40	A or G	15,000
	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	40	A or G	395
	Matador (young grasshoppers only) (P)	0.025-0.034 L (Ground) 0.034 L (Aerial)	$3.80-$4.97 $4.97	50	A or G	64-110
	Malathion 500 (OP)	0.44-0.68 L	$5.90-$9.11	7	A or G	4302
	Malathion 500E (OP)	0.45-0.71 L	$5.56-$8.77	7	A or G	1375-2800
Bertha Armyworm	Lannate (C)	0.089-0.109 kg	$6.60-$8.11	8	A or G	30-34
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.304-0.405 L	$5.21-$7.43	21	A or G	205-418
True Armyworm	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486 L	$6.06-$8.92	21	A or G	205-418
Clover Cutworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	40	A or G	395
Variegated cutworm	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486 L	$6.06-$8.92	21	A or G	205-418
Beet Webworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	40	A or G	395

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Forage Grasses (Timothy, etc.) Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Defoliators						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	$3.00-$6.00	1-2	G	N/A
	Sprays					
	Sevin XLR Plus (C)	0.49-0.93 L (for nymphs or sparse vegetation) 0.93-1.42 L (for adults or application to dense vegetation)	$7.13-$13.81 $13.81-$20.94	1	A or G	649
	Malathion 500 (OP)	0.69 L	$9.11	7	A or G	4302
	Cygon (OP)	0.17 L (young) 0.34 L (adult)	$5.19 $10.37	2 7	A or G	450
	Lagon (OP)	0.17-0.22 L	$4.81-$6.23	2	A or G	60
European skipper (on timothy)	Dipel 2X DF (M)	57-111 g	$4.99-$9.79	N/A	A or G	>4,000

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: M=microbials, C=carbamates, OP=organophosphates

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in **Lentils**

Grasshoppers

Economic Threshold: 2 grasshoppers/m^2 during the flowering and podding stages, especially if two-striped grasshopper is the dominant species.

Lentil Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground feeders and Cutworms						
Cutworms (Army, black, dark-sided, pale western, redbacked, white)	Decis 5EC (P)	0.08 L	$7.36	30	A or G	395
	Pounce (P)	0.073-0.158 L	$7.30-$15.00	Treat prior to 6-leaf stage	G	1030
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (for pale western cutworm only) (OP)	0.354-0.486 L	$6.06-$8.92	21 – 60	A or G	205-418
Defoliators						
Grasshoppers	Decis Flowable (P)	0.032-0.048 L (ground) 0.048 L (air)	$3.72-$5.58	30	A or G	15000
	Decis 5EC (P)	0.04-0.06 L (ground) 0.06 L (air)	$3.68-$5.52	30	A or G	395
	Malathion 500 (OP)	0.68 L	$9.11	30	A or G	4302
	Malathion 500E (OP)	0.69 L	$8.52	14	A or G	1375-2800
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.235-0.486 L	$4.03-$8.92	21-60	A or G	205-418

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, OP=organophosphates

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Mustard Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
Belowground Feeders						
Root Maggot	No insecticides registered					
Sap Feeders						
Lygus Bugs	Decis 5EC (P)	0.06 L	$5.52	7	A or G	395
Defoliators						
Flea beetles	**Seed Treatments**					
	Helix (N)	1500 ml/ 100 kg of seed	N/A	N/A	Seed Treatment	>5,000
	Helix Xtra (N)	1500 ml/ 100 kg of seed	N/A	N/A	Seed Treatment	>5,000
	Gaucho Canola System (N)	0.833 L/ 100 kg of seed	N/A	N/A	Seed Treatment	N/A
	Gaucho Platinum (N)	1.667 L/ 100 kg of seed	N/A	N/A	Seed Treatment	N/A
	Sprays					
	Decis Flowable (P)	0.04-0.06 L	$4.65-$$6.98	7	A or G	15,000
	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Matador (P)	0.034 L	$4.97	50	A or G	64-110
	Furadan 480F (C)	0.06 L	$2.18	21	A or G	7.34
	Malathion 500E (OP)	0.45-0.71 L	$5.56-$8.77	7	A or G	1375-2800
Red Turnip Beetle	Furadan 480F (C)	0.11 L	$3.99	21	A or G	7.34
Cabbage seedpod weevil	Note: Yellow mustard (*Sinapis alba*) is resistant to cabbage seedpod weevil; oriental and brown mustards (*Brassica juncea*) are susceptible to feeding by cabbage seedpod weevil.					
	Matador (adults) (P)	0.034 L	$4.97	50	A or G	64-110
	Decis 5 EC (for control of adults only) (P)	0.08 L	$7.36	7	A or G	395
Diamondback Moth	Decis 5 EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
	Malathion 500E (OP)	0.22-0.34 L	$2.72-$4.20	7	A or G	1375-2800
Bertha Armyworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
Clover Cutworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
Beet webworm	Decis 5EC (P)	0.04-0.06 L	$3.68-$5.52	7	A or G	395
Grasshopper	Matador (young grasshoppers only) (P)	0.025-0.034 L (Ground) 0.034 L (Aerial)	$3.80-$4.97 $4.97	50	A or G	64-110
	Malathion 500E (OP)	0.45-0.71 L	$5.56-$8.77	7	A or G	1375-2800

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: N= neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Oats – See small grain cereals

Grasshopper Management on Pastures, Rangelands, Hay, Headlands, and Roadsides

Note: Insects for biological control of weeds such as leafy spurge may be introduced and established in some areas of Manitoba and Saskatchewan. If grasshopper numbers become high, consider using control strategies and insecticides that will minimize harm to these biological control agents.

Insecticide and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Grasshoppers on rangelands may be effectively managed by applying certain insecticides in 30 metre swaths, alternating with 30 metre untreated swaths. For more information on managing grasshoppers by this method see: http://www.sdvc.uwyo.edu/grasshopper/raats.htm					
Spreadable Bran Baits					
Eco bran (pastures, rangelands, field borders, headlands, right-of-way, roadsides, wastelands) (C)	0.8-1.6 kg	$3.00-$6.00	0-2 (see label)	G	N/A
Sprays					
Decis Flowable (Rangeland, pastures, roadsides) (P))	0.032-0.048	$3.72-$5.58	1 (beef cattle) Dairy cattle must not be fed or grazed on Decis-treated crops	A or G	15,000
Decis 5EC (Rangeland, pastures, roadside, fence row) (P)	0.04-0.06 L	$3.68-$5.52	1 (beef cattle) Dairy cattle must not be fed or grazed on Decis 5EC treated crops.	A or G	395
Ripcord (Roadsides, headlands, and summerfallow) (young grasshoppers only) (P)	0.02-0.028 L	$2.40-$3.36	Treated areas must not be grazed or cut or hay.	G	242-542
Matador (Unimproved pasture, summerfallow) (young grasshoppers only) (P)	0.026-0.034 L (ground) 0.034 L (Aerial)	$3.80-$4.97 $4.97	14	A or G	64-110
Sevin XLR plus (ditchbanks, field borders, headlands, pastures, rangelands, right-of-way, wastelands) (C)	0.48-1.41 L	$7.13-$20.94	0 (ditchbanks, field borders, headlands, right-of-way, wasteland); 1 (pastures, rangeland)	A or G	649
Malathion 500 (Hay only) (OP)	0.69 L	$9.11	7	A or G	4302
Malathion 500E (pastures, rangelands) (young grasshoppers only) (OP)	0.69 L	$8.52	Do not apply to fields occupied by dairy animals, but may be grazed or harvested on the day of application.	A or G	1375-2800

Insecticide and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Diazinon (Rangeland, pasture, ditch banks, roadsides, fence rows, wasteland) (OP)	0.445 L (Diazinon 500E and 50 EC) 0.445 kg (Diazinon 50W)	$7.52 (500 E) $7.28 (50 EC) $11.39 (50W)	Do not spray livestock directly. Dairy and Beef cattle and sheep may be grazed or fed green forage immediately following application. Hay may be fed if not cut for 21 days following application.	G	1500 (500 E) 1960 (50W)
Dibrom (Rangeland, pastures, dairy and horse paddocks) (OP)	0.22-0.34 L (young grasshoppers) 0.28-0.4 L (adult grasshoppers)	$8.08-$12.48 (young) $10.28-$14.68 (adult)	4	A or G	345
Lagon 480E (pasture, wasteland) (OP)	0.22 L (nymphs) 0.34-0.40 L (Adults)	$6.23 $9.62-$11.32	2 days-0.22L rate 7 days-0.34L rate 28 days-0.40L rate	A or G	60
Hay	0.17-0.22 L	$4.81-$6.23	2		
Cygon (pasture, wasteland) (OP)	0.17-0.34 L	$5.19-$10.37	2-7	A or G	450
Lorsban/Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	Ungrazed and unoccupied areas such as roadsides, right of way, and fence lines adjacent to barley, wheat, oats, or canola, and lentils.			A or G	205- 418

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in Peas (Field Peas)

SAP FEEDERS

Aphids

When and How to Monitor: Monitor when field peas begin to flower. Take 180° sweeps or check 10 8-inch (20 cm) plant tips at each stop. Record total number of aphids and calculate average per sweep or plant tip.

Economic Threshold: If, at the beginning of flowering, there are 9 to 12 aphids per sweep or 2-3 aphids per 8-inch (20 cm) plant tip, an insecticide application when 50 percent of plants have produced some young pods will be cost-effective.

Peas (Field Peas) Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders						
Wireworm	No insecticides registered for the control of wireworms in peas					
Seedcorn Maggot	Diazinon 50W (OP)	20g/300ml water/4L seed	$0.51 to treat 4L of seed	Seed Treatment	Seed Treatment	1960
	Agrox CD (OP)			Seed Treatment	Seed Treatment	N/A
	Agrox B-2 (OP)			Seed Treatment	Seed Treatment	N/A
Cutworms (Army, black, dark-sided, pale western, redbacked, white)	Pounce (P)	0.073-0.158 L	$7.30-$15.00	Treat prior to 6 leaf stage	G	1030
Sap Feeders						
Leafhoppers	Malathion 500E (OP)	0.91 L	$11.24	3	A or G	1375-2800
Pea Aphid	Lannate (C)	0.206 kg	$15.31	1	G	30-34
	Malathion 500E (OP)	0.91 L	$11.24	3	A or G	1375-2800
	Lagon (OP)	0.11-0.17 L	$3.11-$4.81	3	A or G	60
Spider mites	Malathion 500E (OP)	0.91 L	$11.24	3	A or G	1375-2800
Defoliator						
Alfalfa Looper	Sevin (C)	1.90 L	$28.22	3	G	649
	Lannate (C)	0.206 kg	$15.31	1	G	30-34

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Potatoes* Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Wireworms	Thimet (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	$0.78-$1.19/ 100m row	At planting application. Do not harvest before 90 days after planting time.	G	27-31
Cutworms (Army, darksided, pale western, redbacked, black, white)	Pounce (P)	0.073-0.158 L	$7.30-$15.00	Treat prior to 6-leaf stage	G	1276
	Ripcord (P)	0.07 L	$8.40	21	G	242-542
	Pyrinex/ Nufos/Lorsban/ Chlorpyrifos 480 EC (Redbacked, black, and darksided cutworms only) (OP)	0.971 L (pre-plant) 0.486-0.971 L (seedling)	$17.43 (pre-plant) $8.72-$17.43 (seedling treatment)	7	G	205-418
Sap Feeders						
Aphids	Fulfill (FD)	78.1 g/acre	$17.53	14	G	>5,000
	Genesis (N)	11.79-17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	$3.05-$4.58/ 100 pounds of potato seed tubers	NA	Seed Treatment	>4,870
	Admire 240 F (N)	Soil Application: 0.344-0.526 L (based on 90 cm row spacing). Foliar Application: 0.081 L	Soil Application: $88.98-$136.05 Foliar Application: $20.69	7 (foliar application)	G	4143-4870
	Lannate (C)	0.2185 kg	$16.21	3	G	30-34
	Malathion 500 (OP)	056-0.80 L	$7.50-$10.72	3	Aor G	4302
	Malathion 500E (OP)	0.61-0.91 L	$7.53-$11.24	3	A or G	1375-2800
	Diazinon 50 W (OP)	0.445 kg	$11.39	14	G	1960
	Diazinon 500E (OP)	0.445 L	$7.52	14	G	1500
	Diazinon 50 EC (OP)	0.445 L	$7.28	14	G	NA
	Cygon (OP)	0.22-0.44 L	$6.71-$13.42	7	G	450
	Lagon (OP)	0.22-0.44 L	$6.23–12.45	7	G	60
	Thimet (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	$0.78-$1.19/ 100m row	90	G	27-31

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Monitor (OP)	0.71-0.91 L	$20.41-$26.16	14	A or G	17
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Orthene (OP)	0.30-0.44 kg	$16.63-$24.39	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	$11.16-$15.13	1	G	110
	Thionex EC (OC)	0.6-0.8 L	$9.75-$13.00	2	G	110
	Thiodan (OC)	0.6-0.8 L	$9.75-$13.00	1	G	107.2
Leafhoppers	Genesis (N)	11.79-17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	$3.05-$4.58/ 100 pounds of potato seed tubers	NA	Seed Treatment	>4,870
	Admire 240F (N)	Soil Application: 0.344-0.526 L (based on 90 cm row spacing).	$88.98-$136.05	NA	G	4143-4870
	Pounce (P)	0.075-0.10 L	$7.30-$10.00	1	G	1276
	Decis 5EC (P)	0.04 –0.06 L	$3.68-$5.52	3	A or G	395
	Matador (P)	0.034-0.050 L	$4.97-$7.31	7	A or G	64-110
	Ripcord (P)	0.025-0.05 L	$3.00-$6.00	7	A or G	242-542
	Sevin (C)	0.50 L	$7.43	7	A or G	649
	Furadan (C)	0.445 L	$16.13	7	G	7.34
	Lannate (C)	0.2185 kg	$16.21	3	G	30-34
	Malathion 500 (OP)	056-0.80 L	$7.50-$10.72	3	A or G	1375-2800
	Malathion 500E (OP)	0.61-0.91 L	$7.53-$11.24	3	A or G	1375-2800
	Diazinon 50 W (OP)	0.445 Kg	$11.39	14	G	1960
	Diazinon 500E (OP)	0.445 L	$7.52	14	G	1500
	Diazinon 50 EC (OP)	0.445 L	$7.27	14	G	NA
	Cygon (OP)	0.22-0.44 L	$6.71-$13.42	7	G	450
	Lagon (OP)	0.22-0.44 L	$6.23-$12.45	7	G	60
	Thimet (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	$0.78-$1.19/ 100m row	90	G	27-31
	Monitor (OP)	0.71-0.91 L	$20.41-$26.16	14	A or G	17
	Guthion SC (OP)	0.91-1.41 L	$14.74-$22.84	7	A or G	21-37
	Guthion 50 Solupak WP (OP)	0.45-0.70 kg	$12.29-$19.12	7	A or G	12.3-24.7
	Dibrom (OP)	0.44 L	$16.15	4	A or G	345
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Orthene (OP)	0.30-0.44 kg	$16.63-$24.39	21	G	1030-1447

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
	Methoxychlor (OC)	1.21-3.64 L	$12.61-$37.93	7	G	5000-7000
	Thionex 50W (OC)	0.45-0.61 kg	$11.16-$15.13	1	G	110
	Thionex EC (OC)	0.6-0.8 L	$9.75-$13.00	2	G	110
	Thiodan (OC)	0.6-0.8 L	$9.75-$13.00	1	G	107.2
Lygus bugs	Pounce (P)	0.075-0.10 L	$7.30-$10.00	1	G	1276
	Decis 5EC (P)	0.04 -0.06 L	$3.68-$5.52	3	A or G	395
	Matador (P)	0.034-0.050 L	$4.97-$7.31	7	A or G	64-110
	Ripcord (P)	0.05 L	$6.00	7	A or G	242-542
	Sevin (C)	2.12-2.59 L	$31.48-$38.46	7	A or G	649
	Furadan (C)	0.445 L	$16.13	7	G	7.34
	Lagon (OP)	0.22-0.44 L	$6.23-$12.45	7	G	60
	Guthion SC (OP)	0.91-1.41 L	$14.74-$22.84	7	A or G	21-37
	Guthion 50 Solupak WP[3] (OP)	0.45-0.70 kg	$12.29-$19.12	7	A or G	12.3-24.7
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (larvae only) (OP)	0.405 L	$7.27	7	G	205-418
	Orthene (OP)	0.30-0.44 kg	$16.63-$24.39	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	$11.16-$15.13	1	G	110
	Thionex EC (OC)	0.6-0.8 L	$9.75-$13.00	1	G	110
	Thiodan (OC)	0.6-0.8 L	$9.75-$13.00	1	G	107.2
Defoliators						
Colorado potato beetle	Note: Colorado potato beetles have been found to be resistant to several families of insecticides in localized areas of Manitoba. Rotation between different families of insecticides is essential.					
	Success (S)	0.05-0.066 L	$15.64-$30.82	7	G	>5,000
	Novodor FC (M)	4.0-8.0 L	$58.20-$116.40	1	G	74, 000
	Genesis (N)	11.79-17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	$3.05-$4.58/ 100 pounds of potato seed tubers	NA	Seed Treatment	>4,870
	Admire (N)	In furrow: 0.345 -0.525 L Foliar: 0.080 L	In furrow: $88.98-$136.05 Foliar: $20.69	7	G	4143-4870
	Pounce (P)	0.075 -0.10 L	$7.30-$10.00			
	Decis 5EC (P)	0.04 -0.06 L	$3.68-$5.52	3	A or G	395
	Matador (P)	0.034-0.050 L	$4.97-$7.31	7	A or G	64-110
	Ripcord (P)	0.025-0.05 L	$3.00-$6.00	7	A or G	242-542
	Sevin (C)	0.50 L	$7.43	7	A or G	649
	Furadan (C)	0.222 L	$8.05	7	G	7.34

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
	Malathion 500 (OP)	056-0.80 L	$7.50-$10.72	3	A or G	4302
	Malathion 500E (OP)	0.61-0.91 L	$7.53-$11.24	3	A or G	1375-2800
	Diazinon 50 W (OP)	0.445 kg	$11.39	14	G	1960
	Diazinon 500E (OP)	0.445 L	$5.52	14	G	1500
	Diazinon 50 EC (OP)	0.445 L	$7.28	14	G	NA
	Thimet (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	$0.78-$1.19/ 100m row	90	G	27-31
	Monitor (OP)	0.71-0.91 L	$20.41-$26.16	14	A or G	17
	Guthion SC (OP)	0.70 L	$11.34	7	A or G	21-37
	Guthion 50/ Sniper (OP)	0.34 kg	$9.29	7	A or G	12.3-24.7
	Dibrom (OP)	0.44 L	$16.15	4	A or G	345
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Lorsban/ Nufos/ Pyrinex/ Chlorpyrifos 480 EC (larvae only) (OP)	0.405 L	$7.27	7	G	205-418
	Thionex 50 W (OC)	0.45-0.61 kg	$11.16-$15.13	1	G	110
	Thionex EC (OC)	0.6-0.8 L	$9.75-$13.00	2	G	110
	Thiodan (OC)	0.6-0.8 L	$9.75-$13.00	1	G	107.2
	Methoxychlor (OC)	1.21-3.64 L	$12.61-$37.93	7	G	5000-7000
Potato Flea Beetle	Genesis (N)	11.79-17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	$3.05 - $4.58/ 100 pounds of potato seed tubers	NA	Seed Treatment	>4,870
	Admire 240 F (suppression) (N)	Soil Application: 0.344-0.526 L (based on 90 cm row spacing).	$88.98-$136.05	NA	G	4143-4870
	Pounce (P)	0.075-0.10 L	$7.30-$10.00	1	G	1276
	Decis 5EC (P)	0.04 –0.06 L	$3.68-$5.52	3	A or G	395
	Matador (P)	0.034-0.050 L	$4.97-$7.31	7	A or G	64-110
	Ripcord (P)	0.025-0.05 L	$3.00-$6.00	7	A or G	242-542
	Sevin (C)	0.50 L	$7.43	7	A or G	649
	Furadan (C)	0.222 L	$8.05	7	G	7.34
	Lannate (C)	0.2185 kg	$16.21	3	G	30-34
	Diazinon 50 W (OP)	0.445 Kg	$11.39	14	G	1960
	Diazinon 500E (OP)	0.445 L	$7.52	14	G	1500
	Diazinon 50 EC (OP)	0.445 L	$7.28	14	G	NA

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
	Thimet (reduction of Flea beetles) (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	$0.78-$1.19/ 100m row	90	G	27-31
	Monitor (OP)	0.71-0.91 L	$20.41-$26.16	14	A or G	17
	Guthion SC (OP)	0.91-1.41 L	$14.74-$22.84	7	A or G	21-37
	Guthion 50 Solupak WP (OP)	0.45-0.70 kg	$12.29-$19.12	7	A or G	12.3-24.7
	Dibrom (OP)	0.44 L	$16.15	4	A or G	345
	Imidan (OP)	0.91 kg	$22.20	7	G	285
	Lorsban/ Nufos/ Pyrinex/ Chlorpyrifos 480 EC (OP)	0.405 L	$7.27	7	G	205-418
	Orthene (OP)	0.30-0.44 kg	$16.63-$24.39	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	$11.16-$15.13	1	G	110
	Thionex EC (OC)	0.6-0.8 L	$9.75-$13.00	1	G	110
	Thiodan (OC)	0.6-0.8 L	$9.75-$13.00	1	G	107.2
	Methoxychlor (OC)	1.21-3.64 L	$12.61-$37.93	7	G	5000-7000
Blister beetle	Methoxychlor (OC)	1.21-3.64 L	$12.61-$37.93	7	G	5000-7000
Variegated cutworm	Pounce (P)	0.075 L	$7.30	1	G	1276
	Ripcord (P)	0.07 L	$8.40	7	G	242-542
	Sevin (C)	0.1-0.125 L/ 300 m of row	$1.49-$1.86/ 300 m of row	7	A or G	649
			Stem Borers			
European Corn Borer	Success (S)	0.059 L	$27.14	7	G	>5,000
	Sevin (C)	1.01-2.59 L	$15.00-$38.46	7	A or G	649
	Guthion SC (OP)	0.91-1.41 L	$14.74-$22.84	7	A or G	21-37
	Guthion 50 Solupak WP (OP)	0.45-0.70 kg	$12.29-$19.12	7	A or G	12.3-24.7

*Before using any pesticide on potatoes, consult the list of **Agricultural Pesticides Approved for Use**, available from Simplot Canada and McCain Foods (Canada).

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: M=microbials, S=spinosyns, FD=feeding disruptors, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in Rye

BELOWGROUND FEEDERS AND CUTWORMS

Cutworms

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in field.

When and How to Monitor: Monitor late May to mid-July. Sample 5 to 10 locations in the field. At each location, check several 0.25 m^2 (50cm x 50cm) areas. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m^2.

Economic Threshold: Pale western cutworm – 3-4/m^2; Redbacked and army cutworm – 5-6/m^2. Well established crops with good moisture conditions can tolerate higher numbers.

SAP FEEDERS

Aphids

Typical Damage: Visible wilting of plants, yellow patches in fields, plants are sticky.

When and How to Monitor: Monitor July through early August. Check 20 stems in each of 5 areas. Counts should be at least 50 paces apart, and observations should be made well into the center of the field. Too frequently farmers become alarmed after checking a few plants along the margins, especially near shelterbelts, where populations are high. Record the total number of aphids and calculate the average per plant.

Economic Threshold: 12-15 aphids/stem until about 2 weeks after flowering.

DEFOLIATOR

Armyworms

Typical Damage: Leaves stripped from plants, awns chewed from heads, heads clipped.

When and How to Monitor: Monitor mid-June through July. At each stop shake plants in 1 yard^2 (m^2) area and carefully check soil surface for dislodged larvae.

Economic Threshold: 10 larvae/m^2. For migrating Armyworms: Treat a couple of swaths ahead of the infestation in the direction of movement to form a barrier strip.

Rye Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Wireworms	No insecticides registered for the control of wireworms in rye.					
Cutworms (Army, black, darksided, pale western, redbacked, white)	Pounce (P)	0.075-0.158 L	$7.30-$15.00	Treat pior to 6 leaf stage	G	1030
Sap Feeders						
Aphids	Malathion 500 (OP)	0.60-0.80 L	$8.04-$10.72	7	A or G	4302
	Malathion 500E (OP)	0.45-1.11 L	$5.56-$13.71	7	A or G	1375-2800

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
			Defoliators			
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6 kg	\$3.00-\$6.00	14	G	N/A
	Sprays					
	Sevin XLR Plus (C)	0.50-1.01 L	\$7.43-\$15.00	14	A or G	649
	Malathion 500 (OP)	0.69 L	\$9.11	7	A or G	4302
	Malathion 500E (OP)	0.91-1.11 L	\$11.24-\$13.71	7	A or G	1375-2800
	Lagon 480E (OP)	0.22-0.40 L	\$6.23-\$11.32	2	A or G	60
	Guthion (Borders Only) (OP)	0.44-0.70 L (Guthion SC) 0.21-0.34 kg (Guthion 50 Solupak WP)	\$7.13-\$11.34 (Guthion SC) \$5.74-\$9.29 (Guthion 50 Solupak WP)	30	A or G	21-37/12.3-24.7
	Sniper (OP)	0.21-0.34 kg	\$6.55-\$10.60	30	A or G	12.3-24.7
True armyworm	Sevin XLR Plus (C)	1.01-2.12 L	\$15.00-\$31.48	14	A or G	649
	Malathion 500 (OP)	0.60-0.80 L	\$8.04-\$10.72	7	A or G	4302
	Malathion 500E (OP)	0.45-1.11 L	\$5.56-\$13.71	7	A or G	1375-2800
	Methoxychlor (OC)	2.83-4.55 L	\$29.49-\$47.41	7	G	5,000-7,000

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Safflower Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
			Defoliators			
Grasshoppers	Lagon (OP)	0.22-0.40 L	\$6.23-\$11.32	21	A or G	60

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: OP=organophosphates

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Field Scouting in Small Grain Cereals (wheat, barley, oats)

BELOWGROUND FEEDERS AND CUTWORMS

Cutworms

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in field.

When and How to Monitor: Monitor late May to mid-July. Sample 5 to 10 locations in the field. At each location, check several 0.25 m^2 (50cm x 50cm) areas. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m^2.

Economic Threshold: Pale western cutworm-3-4/m^2; Redbacked and army cutworm-5-6/m^2. Well established fall-seeded crops or spring seeded crops with good moisture conditions can tolerate higher numbers.

SAP FEEDERS

Aphids

Typical Damage: Visible wilting of plants, yellow patches in fields, plants are sticky.

When and How to Monitor: Monitor July through early August. Check 20 stems in each of 5 areas. Counts should be at least 50 paces apart, and observations should be made well into the center of the field. Too frequently farmers become alarmed after checking a few plants along the margins, especially near shelterbelts, where populations are high. Record the total number of aphids and calculate the average per plant.

Economic Threshold: 12-15 aphids/stem prior to the soft dough stage.

Barley Thrips

When and How to Monitor: Sampling should begin when the flag leaf is first visible and continue until the head is completely emerged from the boot. Barley thrips exhibit an edge effect; there are usually more thrips near protected field margins than other areas of the field. Most thrips can be found under the top 2 leaf sheaths. Unroll the leaf sheaths away from the stem to find the thrips.

Economic Threshold: Insecticide treatments are only effective when applied before heading is complete.

Treat when thrips are equal to or greater than the number calculated by:

Threshold (Thrips/stem) = (Cost of Control ÷ expected $ value per bushel)/0.4

DEFOLIATORS

Grasshoppers

Typical Damage: Black strips along margins of newly emerging crops, head clipping later in season.

When and How to Monitor: Monitor May through July. Check along edges of crop, particularly areas adjacent to hayland, pastures and roadsides. Estimate number of hoppers/yard^2 (m^2).

Economic Threshold: 8-13 grasshoppers/m^2. Early in the season, when grasshoppers are small, 18 grasshoppers/m^2 and visible crop damage may be a more appropriate threshold.

A rough estimate for an economic threshold for grasshoppers in crops to be used as greenfeed has been suggested at 20 grasshoppers/m^2 or higher.

Armyworms

Typical Damage: Leaves stripped from plants, awns chewed from heads, heads clipped.

When and How to Monitor: Monitor mid-June through July. At each stop shake plants in 1m^2 area and carefully check soil surface for dislodged larvae.

Economic Threshold: 10 larvae/m^2. For migrating Armyworms: Treat a couple of swaths ahead of the infestation in the direction of movement to form a barrier strip.

SEED FEEDERS ONLY

Wheat Midge (wheat only)

When and How to Monitor: Monitor in July when crop emerges from boot stage until flowering. Check crop canopy at dusk for signs of wheat midge adult activity. At each stop, examine 10 heads. Record the number of midge adults observed on or near heads. Calculate average number of midge per head.

Sticky traps may be used to capture adult midge activity in wheat fields.

Economic Threshold: *For yield only:* 1 adult midge per 4 to 5 heads. At this level of infestation, wheat yields will be reduced by approximately 15% if the midge is not controlled. *To maintain optimum grade:* 1 adult midge per 8 to 10 wheat heads during the susceptible stage.

Small Grain Cereals (wheat, barley, oats) Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Wireworms	Cruiser (barley and wheat only) (N)	33 – 50 mL/ 100 kg seed			Seed Treatment	5,523
Cutworms (Army, dark sided, pale western, redbacked)	Decis 5EC (P)	0.08 L	$7.36	31 (oats) 40 (barley, wheat)	A or G	395
	Ricord (barley and wheat only) (P)	0.07 L	$8.40	21	G	242-542
	Pounce (P)	0.075-0.158	$7.30-$15.00	Treat prior to 6-leaf stage	G	1030
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354-0.486 L	$6.06-$8.92	60	A or G	205-418
Sap Feeders						
Aphids	Malathion 500 (OP)	0.60-0.8 L	$8.04-$10.72	7	A or G	4302
	Malathion 500E (OP)	0.45-1.11 L	$5.56-$13.71	7	A or G	1375-2800
	Cygon 480 (OP)	0.17 L	$5.19	21	A or G	450
Thrips	Lannate (C)	0.1214kg	$9.01	20	A or G	30-34
	Cygon (OP)	0.40 L	$12.20	21	A or G	450
	Lagon (OP)	0.40 L	$11.32	7	A or G	60
Defoliators						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6kg	$3.00-$6.00	14 (oats, wheat) 28 (barley)	G	
	Sprays					
	Decis Flowable (P)	0.32-0.048 L	$3.72-$5.58	31 (oats) 40 (wheat, barley)	A or G	15000
	Decis 5EC (P)	0.04-0.06 L (ground) 0.06 L (air)	$3.68-$5.52 (ground), $5.52 (air)	31 (oats) 40 (wheat, barley)	A or G	395
	Ripcord (young grasshoppers only) (wheat and barley only) (P)	0.02-0.028 L	$2.40-$3.36	30 (wheat) 60 (barley)	G	242-542
	Matador (young grasshoppers only) (P)	0.025 – 0.034 L (Ground) 0.034 L (Aerial)	$3.80-$4.97 $4.97	Do not apply within 28 days of harvest or 14 days of livestock foraging	A or G	64-110
	Sevin XLR Plus (C)	0.50-1.01 L	$7.43-$15.00	14 (wheat, oats) 28 (barley)	A or G	649
	Malathion 500 (OP)	0.68 L	$9.11	7	A or G	4302

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
	Malathion 500E (young grasshoppers) (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.235-0.354 L	$4.03-$6.50	60	A or G	205-418
	Lagon (OP)	nymphs-0.22 L adults-0.34-0.40 L	$6.23 $9.62-$11.32	2	A or G	60
True Armyworm	Sevin XLR Plus (C)	1.01-2.12 L	$15.00-$31.48	14 (wheat, oats) 28 (barley)	A or G	649
	Lannate (C)	0.1093-0.2185kg	$8.11-$16.21	20	A or G	30-34
	Malathion 500 (OP)	0.60-0.80 L	$8.04-$10.72	7	A or G	4302
	Malathion 500E (OP)	0.45-1.11 L	$5.56-$13.71	7	A or G	1375-2800
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.354 –0.486 L	$6.06-$8.92	60	A or G	205-418
	Methoxychlor (OC)	2.83-4.55 L	$29.49-$47.41	7	G	5000-7000
Pests of Seed Only						
Wheat Midge (a pest of wheat only)	Rotate Crops – Continuous wheat cropping encourages higher wheat midge populations.					
	Lorsban/ Nufos/Pyrinex/ Chlorpyrifos 480 EC (OP)	0.336-0.405 L	$5.76-$7.43	60	A or G	205- 418
	Cygon 480/ Lagon (OP)	0.40 L	$11.32-$12.20	21	A or G	450
Stem-Borers						
Hessian Fly	• Never plant wheat in the same field 2 years in a row in areas where Hessian flies are a problem. • The spring wheat cultivar Superb is partially resistant to the Hessian fly. • Early seeded spring wheat is less susceptible to stem breakage caused by Hessian fly than later seeded wheat. • Winter wheat planted in September will likely be free of Hessian flies.					
Wheat stem sawfly	Solid-stem wheat varieties (such as the hard red spring wheat varieties AC Eatonia and AC Abbey) can reduce damage by wheat stem sawfly larvae compared to susceptible varieties, however the level of control can vary depending on environmental conditions.					

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Scouting for insects in Soybeans

Soybean Aphid

Note – Although soybean aphids have been found in Manitoba since 2001, populations have never been found in economic numbers. Monitoring soybean fields for soybean aphids and other insects is advised however. Note that very high numbers of soybean aphid are required to cause economic damage to soybeans.

Typical Damage: Soybean aphids suck sap from soybean plants. Infested leaves may wilt or curl when infestations are large. Other symptoms may include plant stunting, reduced pod and seed count, and yellowing of leaves.

Economic Threshold: When there are on average more than 250 aphids per plant, and the plants are in the R1 (beginning bloom) to R4 (full pod) growth stages, treatment would be economical.

Soybean Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD$_{50}$ (Mammalian Toxicity)[2]
Belowground Feeders						
Wireworms	No insecticides registered for the control of wireworms in soybeans					
Seedcorn Maggot	Agrox CD, Agrox B-2	See "Disease Control – Seed Treatments" section of this guide.				N/A
	Diazinon 50W	20 g/300 mL water/4L seed	–	–	Seed Treatment	1960
	Sap Feeders					
Soybean Aphid	Lagon (OP)	0.28-0.40 L	$7.92-$11.32	30	A or G	60
Leafhoppers	Lagon (OP)	0.28-0.40 L	$7.92-$11.32	30	A or G	60
Lygus bugs	Lagon (OP)	0.28-0.40 L	$7.92-$11.32	30	A or G	60
Spider mites	Cygon 480 (OP)	0.40 L	$12.20	30	A or G	450
	Lagon (OP)	0.40 L	$11.32	7	A or G	60
Defoliators						
Grasshopper	No insecticides registered for control of grasshoppers in soybeans.					

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: OP=organophosphates, OC=organochlorines

[2] LD$_{50}$ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD$_{50}$.

Stored Grain Insect Control –

See Insect Control in Stored Grain at the end of the Insect Control section of this guide.

Summerfallow –

See grasshopper management on Pastures, etc.

Scouting for insects in Sunflower

BELOWGROUND FEEDERS AND CUTWORMS

Cutworm

Typical Damage: Plants cut off at or near soil surface, some leaf feeding above ground, plants missing from rows, bare patches appearing in field.

When and How to Monitor: Monitor late May to mid-July. Sample 5 to 10 locations in the field. At each location, check several 0.25 m^2 (50cm x 50cm) areas. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m^2.

Economic Threshold: When cutworm densities exceed 10/m^2.

DEFOLIATOR

Sunflower Beetle

Typical Damage: **Adults:** Leaves of seedling plants chewed or completely destroyed late May through June, shot-holes or large areas of leaves chewed July through August. **Larvae:** Leaves of plants chewed or completely destroyed.

When and How to Monitor: **Adults:** Monitor May through June on seedling plants. Examine 10 plants at random at each stop. **Larvae:** Monitor July through mid-August. Examine 10 plants at random at each sampling site. Peel back leaves around growing tip and record total number of larvae found. Calculate average number per plant.

Economic Threshold: **Adults:** 1-2/seedling; **Larvae:** 10 to 15/plant or 25-30% defoliation.

Insects affecting the seeds

PESTS OF SEED ONLY

Red Sunflower Seed Weevil

Typical Damage: Seeds partly or completely destroyed, exit hole in hull. Shriveled kernels, kernels completely destroyed.

When and How to Monitor: Monitor fields when ray petal being to form and continue every 2 to 3 days until pollination is complete. When scouting, use the X pattern and begin counts at least 70 to 100 feet into the field to avoid margin effects. Examine 5 plants at each site for a total of 25 plants. For checking individual sunflower heads, brush the face of the head vigorously to bring the weevils to the surface, or use a commercial preparation of mosquito repellent containing diethyl toluamide (DEET) to spray the heads. This will cause the weevils to move out of hiding spots. Record total number of weevils and calculate average per head.

Economic Threshold:

Confection Sunflowers: 1-2 weevils/plant. Control is based on a need to keep seed damage below 3 or 4% because of industry standards.

Oilseed sunflowers: 12-14 weevils/head.

The ideal plant stage to treat is when most plants in the field are at 40% pollen shed (R5.4).

Banded Sunflower Moth

When and How to Monitor: Scouting should be conducted prior to the R5.1 plant growth stage. Count moths on 20 plants from 5 different sites for a total of 100 plants. Sample in early evening or early morning when the moths are most active to get the most accurate counts.

Economic Threshold: 1 moth per 2 plants when scouting at dusk or later. If treatment is warranted, it should be applied at the R5.1 sunflower plant growth stage.

Sunflowers Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Belowground Feeders and Cutworms						
Wireworms	Wireworms may sometimes damage sunflowers. Seeding sunflowers when the soil temperature is at least 8 to 10°C at 1 to 1.5 inches depth may minimize damage by wireworms.					
Cutworms	Pounce (P)	0.075-0.158 L	$7.30-$15.00	Treat prior to 6-leaf stage	G	1030
	Lorsban/Pyrinex/ Nufos/Chlorpyrifos 480 EC (OP)	0.486 L	$8.72	42	G	205-418
Defoliators						
Grasshoppers	No insecticides registered for control of grasshoppers in sunflowers.					
Sunflower Beetle	Decis 5EC (P)	0.04 L	$3.68	70	A or G	395
	Matador (P)	0.017 – 0.025 L (Ground) 0.034 L (Aerial)	$2.49-$3.80 $4.97	70	A or G	64-110
	Ripcord (P)	0.028 L	$3.36	70	A or G	242-542
	Furadan (C)	0.11 L	$3.99	60	G	7.34
	Thionex EC (OC)	0.6 L	$9.75	60	G	110
	Thiodan (OC)	0.6 L	$9.75	60	G	107.2
Pests of Seed Only						
Sunflower Seed Weevil	Early planting helps to reduce seed damage by sunflower seed weevils					
	Ripcord (P)	0.028 L	$3.36	70	A or G	242-542
	Lorsban/Nufos (OP)	0.486 L	$8.72	42	A or G	272/205
Banded sunflower moth	Late planting may provide some control					
Sunflower Moth	Dipel (young larvae) (M)	0.1275- 0.2529 kg	$11.22-$22.26	N/A	A or G	> 4,000

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: M=microbial, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Field Scouting in Sweet Clover

DEFOLIATOR

Sweetclover Weevil

Typical Damage: Adults chew crescent-shaped and jagged notches in leaves and can completely defoliate plants.

When and How to Monitor: Inspect clover seedlings for weevil damage in spring as the seedlings emerge. In mid-summer and throughout August, inspect first-year clover stands for damage along crop margins. Invading weevils move into these stands only as far as necessary to satisfy their food requirements, so an insecticide application to affected field margins is usually all that is required. Visually estimating the number of weevils per plant must be done carefully because weevils fall from plants easily and are difficult to see on the ground.

Economic Threshold: 1st year stands: 1 weevil adult/3 seedlings (1/5 seedlings under dry conditions). 2nd year stands: 9-12 weevil adults/plant.

Sweet Clover Insect Management Chart

Insect	Insecticide (and insecticide group[1])	Rate/Acre	Cost/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD_{50} (Mammalian Toxicity)[2]
Defoliators						
Grasshoppers	**Spreadable Bran Baits**					
	Eco bran (C)	0.8-1.6kg	$3.00-$6.00	2	G	N/A
	Sprays					
	Sevin XLR Plus (C)	0.50-1.01 L	$7.43-$15.00	2	A or G	649
	Malathion 500E (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Guthion SC Guthion 50 Solupak (Border Application Only) (OP)	0.44- 0.70 L 0.21-0.34 L	$7.13-$11.34 $5.74-$9.29	21 21	A or G	21-37 12.3-24.7/7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5000-7000
Sweetclover Weevil	Locate new seedlings as far as possible from 2nd-year clover. Cultivating second-year stands of sweet clover silage and hay as soon as possible after the crop is taken kills the new- generation weevil larvae in the soil.					
	Malathion 500 (OP) Malathion 500E (OP)	0.56-1.01 L 0.61-1.01 L	$7.50-$13.53 $7.53-$12.47	7-Cattle may be returned immediately after spraying.	A or G	1375-2800
	Cygon/Lagon (OP)	0.34-0.44 L	$9.62-$13.42	28	A or G	450/60
	Guthion SC Guthion 50/ Sniper (OP)	0.91 L 0.45 L	$14.74 $12.29	21	A or G	21-27 12.3-24.7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5000-7000
Alfalfa Weevil	Sniper (OP)	0.45-0.71 kg	$13.10-$20.37	21	A or G	12.3-24.7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5000-7000
Blister Beetle	Sevin XLR Plus (C)	1.01-1.6 L	$15.00-$23.76	2	A or G	649
True Armyworm	Methoxychlor (OC)	2.83-4.55 L	$29.49–$47.41	7	G	5000-7000
Beet Webworm	Sevin XLR Plus (C)	1.01-2.12 L	$15.00-$31.48	2	A or G	649
Sap Feeders						
Lygus Bugs	Dibrom (OP)	0.44-0.88 L	$16.15-$32.30	4	A or G	345
	Guthion SC Guthion 50/ Sniper (OP)	0.91-1.41 L 0.45-0.70 L	$14.74-$22.84 $12.29-$19.12	21	A or G	21-37
Leafhoppers	Sevin XLR Plus (C)	1.01-1.6 L	$15.00-$23.76	2	A or G	649
	Malathion 500E (OP)	0.91-1.11 L	$11.24-$13.71	7	A or G	1375-2800
	Dibrom (OP)	0.44-0.88 L	$16.15-$32.30	4	A or G	345
	Guthion SC Guthion 50/ Sniper (OP)	0.91-1.41 L 0.45-0.70 L	$14.74-$22.84 $12.29-$19.12	21	A or G	21-37 12.3-24.7/7
	Methoxychlor (OC)	2.02-2.53 L	$21.05-$26.36	7	G	5000-7000
Aphids	Dibrom (OP)	0.44-0.88 L	$16.15-$32.30	4	A or G	345

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

[1] Insecticide Group: C=carbamates, OP=organophosphates, OC=organochlorines

[2] LD_{50} values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD_{50}.

Timothy –

See forage grasses

Wheat –

See small grain cereals

Insecticide Poisoning in Humans

Organophosphate (OP) and carbamate insecticides (identified on the Insecticide Groups chart page 318) can pose a more serious risk to unprotected persons. Poisonings can occur while mixing, loading and/or during the application of these products without the appropriate protective equipment or measures. These pesticides are readily absorbed through the skin or the lungs, and can act as nervous system toxins. Overexposure can produce symptoms such as headache, nausea, pupil dilation and excessive sweating and salivation. Higher doses may cause breathing difficulties, muscle twitching, weakness and spasms. Very high doses have caused respiratory failure and death.

Both OP and carbamate pesticides inhibit an enzyme called cholinesterase. Measurements of cholinesterase in the blood before and during the application season can indicate harmful exposures to OPs and carbamates. **Persons who intend to mix, load and/or apply these types of pesticides repeatedly during a season, need a baseline and repeat measurements. Consult your doctor before the spraying season to arrange for these measurements.** For more information, see ***Monitoring Exposure to Organophosphorous and/or Carbamate Insecticides,*** a joint publication of Saskatchewan Labour and Saskatchewan Health. This publication can be viewed on the SK Labour web site: www.labour.gov.sk.ca.

Degree of Risk and Hazard Rating:

(see page 10 for full description)

Danger poison: LD_{50} is less than 500 mg/kg and indicates high toxicity.

Warning poison: LD_{50} is between 500-1,000 mg/kg and indicates moderate toxicity.

Caution poison: LD_{50} is between 1,000-2,500 mg/kg and indicates low toxicity.

(LD_{50} values are used to rate the toxicity of pesticides. They refer to the dose of pesticide (in mg per kg) that is lethal to 50 percent of a test animal population).

Resistance of Insects to Insecticides

Repeated use of the same insecticide, or insecticides with the same mode of action, against a particular insect in a given area may result in the effectiveness of the insecticide being reduced.

To delay or prevent resistance of insects to insecticides:

1. Integrate different control methods (cultural, biological, chemical) into insect control programs whenever possible.
2. Use insecticides only when the economic threshold for a pest has been surpassed and natural controls fail to limit economic damage, and
3. Rotate between insecticides with different modes of action, particularly if several applications are made in a season.
4. Keep accurate records of insecticides used for each of your fields.

Insecticides can be classified according to their similarity in chemical structure (chemical group in the table below), and by mode of action (the process by which the insecticide kills the insect). The "Group" column in the table below separates insecticides based on their mode of action. By selecting products with different modes of action for an insecticide rotation program, insecticide resistance can be prevented.

Insecticide Groups Based on Modes of Action

GROUP	CHEMICAL GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ENTRY
1A	Carbamates	Sevin, Eco Bran	carbaryl	contact/ ingestion (Sevin) ingestion (Eco Bran)
		Lannate	methomyl	contact/ ingestion
		Furadan	carbofuran	contact/ ingestion
1B	Organophosphates	Malathion	malathion	contact
		Diazinon	diazinon	contact/ ingestion
		Orthene	acephate	contact/ ingestion
		Dibrom	naled	contact/ ingestion
		Imidan	phosmet	
		Lorsban, Pyrinex, Nufos, Chlorpyrifos	chlorpyrifos	contact/ ingestion/ inhalation
		Cygon, Lagon	dimethoate	contact/ ingestion
		Thimet	phorate	ingestion
		Monitor	methamidophos	contact/ ingestion
		Guthion, Sniper	azinphos-methyl	contact
2A	Organochlorines	Thiodan, Thionex	endosulfan	contact/ ingestion
3	Diphenylethanes	Methoxychlor	methoxychlor	contact/ ingestion
	Pyrethroids	Decis	deltamethrin	contact/ ingestion
		Ripcord	cypermethrin	contact/ ingestion
		Matador	cyhalothrin-lambda	contact/ ingestion
		Pounce	permethrin	contact/ ingestion
		Tempo	cyfluthrin	
4	Neonicotinoids	Helix, Cruiser	thiamethoxam	ingestion
		Admire, Gaucho, Genesis	imidacloprid	contact/ ingestion (Admire) ingestion (Gaucho, Genesis)
		Prosper, Poncho	clothianidin	ingestion
5	Spinosyns	Success	spinosad	
8B	Fumigant	Phostoxin	aluminum phosphide	
9B	Feeding Disruptors	Fulfill	pymetrozine	ingestion mainly, some contact activity
11	Microbials	Dipel	*Bacillus thuringiensis var. Kurstaki*	ingestion
		Novodor	*Bacillus thuringiensis var. Tenebrionis*	ingestion

A more detailed table showing insecticides organized by mode (site) of action, and specific information on the mode (site) of action for the different groups can be found in Appendix III of PMRA Regulatory Directive DIR99-06 found at: http://www.hc-sc.gc.ca/pmra-arla/english/pdf/dir/dir9906-e.pdf

Insecticide Directory

For rates, costs, and preharvest intervals for insecticides, see the insect management charts on pp. 282-318.

Admire

Company:

Bayer CropScience PCP# 24094

Formulation:

240 g/L imidacloprid formulated as a flowable

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, aphids, leafhoppers, potato flea beetles (suppression only)

Application:

Do not apply by air.

Soil application: For best results, direct spray on the seed pieces or seed potatoes in the furrow. The higher rate is recommended when extended length of control is needed. Do not follow a soil application with a foliar application.

Foliar application: A maximum of two foliar applications of Admire may be made per season.

How it Works:

Admire is a systemic, chloronicotinyl insecticide.

Restrictions:

Do not apply within 15m of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc. Do not apply through any type of irrigation system.

Precautions:

Do not re-enter treated areas for 24 hours after foliar application of Admire. Admire is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Highly toxic to aquatic invertebrates and birds.

Hazard Rating:

Caution Poison

Cygon/Lagon

Company:

IPCO (Cygon 4E - PCP# 09807)
Cheminova (Cygon 480-Ag - PCP# 25651)
United Agri Products (Lagon 480E - PCP# 9382)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulations:

Cygon/Lagon-480 g/L dimethoate formulated as an emulsifiable concentrate. Container size-10 L.

Insects Controlled and Registered Crops:

CROP	Cygon 480-Ag INSECT	Lagon 480E INSECT	Cygon 4E INSECT
peas		aphids	
potatoes	aphids, leafhoppers	aphids, leafhoppers, Lygus bugs	aphids (ground application only), leafhoppers
alfalfa (*seed production only)	aphids, leafhoppers, Lygus bugs, plant bugs*, alfalfa blotch leaf miner, grasshoppers, reduction of alfalfa weevil larvae	aphids, leafhoppers, alfalfa blotch leafminers, grasshoppers, reduction of alfalfa weevil larvae, Lygus bugs*, plant bugs*	aphids, grasshoppers, leafhoppers, Lygus bugs, plant bugs, reduction of alfalfa weevil larvae, sweet clover weevil
canaryseed	aphids	aphids	
canola / rapeseed	aphids, leafhoppers, grasshoppers,	aphids, leafhoppers, grasshoppers,	
forage crops	Lygus bugs, plant bugs, grasshoppers	grasshoppers, aphids (suppression only of Russian wheat aphid)	aphids, grasshoppers, leafhoppers, Lygus bugs, plant bugs, reduction of alfalfa weevil larvae, sweet clover weevil
sweet clover, red clover, alsike clover	sweet clover weevil	sweet clover weevil	
pastures, wastelands	grasshoppers	grasshoppers	grasshoppers
wheat	wheat midge, thrips, aphids (suppression only of Russian wheat aphid)	thrips, grasshoppers, Russian wheat aphid	
barley, oats	thrips, aphids (suppression only of Russian wheat aphid)	thrips, grasshoppers,	
flax	aphids	aphids	
soybeans	aphids, leafhoppers, Lygus bugs, mites	aphids, leafhoppers, Lygus bugs, spider mites	

Application:

May be applied by air or ground equipment (unless otherwise specified above). Apply when insects exceed economic threshold numbers and use sufficient water for good coverage. Use higher rates for adult insects, heavy infestations or dense canopy.

How it Works:

Broad spectrum, systemic (within the plant) and contact, organophosphate insecticide and acaricide.

Restrictions:

Grazing: Remove cattle prior to spraying. Read label carefully to determine livestock reentry period.

Storage: Store at temperatures between 4°C and 30°C and in areas away from feed and food.

Decis

Company:

Bayer CropScience
(Decis 5EC - PCP# 17734; Decis Flowable - PCP# 20078)

Formulations:

50 g/L deltamethrin formulated as an emulsifiable concentrate and as a flowable.

Container sizes: Decis 5EC-2.5 L and 2 L jugs; Decis Flowable - 3 L jugs.

Insects Controlled and Registered Crops:

DECIS 5EC

CROPS	INSECT
Alfalfa (seed crops only)	Lygus bug, alfalfa weevil
Field corn	European corn borer
Potatoes	Colorado potato beetle, potato flea beetle, Lygus bugs, leafhopper
Canola (rapeseed), mustard	Flea beetle, clover cutworm, bertha armyworm, diamondback moth, beet webworm, cabbage seedpod weevil, Lygus bugs, grasshoppers
Sunflowers	Sunflower beetle
Wheat, barley, oats, lentils, flax	Cutworms, grasshoppers
Rangeland, pastures, roadside, fence row	Grasshoppers
Flax	Cutworms, beet webworm
Canola	Flea beetle
Wheat, barley, oats, flax, lentils	Grasshoppers
Rangeland, pastures, roadsides	Grasshoppers

Application:

Decis may be applied by air or ground equipment to all crops with the exception of alfalfa and corn, which require ground application only. Apply when insects exceed economic threshold numbers with sufficient water for good coverage. Use higher rates for severe infestations, on dense foliage or when a number of insect growth stages are present. Works as contact and stomach poison usually with very quick results.

Tank Mixes:

May be tank mixed with the following herbicides:

Hoe-Grass 284 (5 EC only) Hoe-Grass II (5 EC only)
Pardner, Buctril M
Banvel, MCPA
2,4 -D,
Achieve (Decis Flowable only)
Puma^{120} Super

How it Works:

Decis is a non-systemic, synthetic pyrethroid which works by contact and ingestion.

Effects of Weather:

Do not spray under a strong temperature inversion, or when temperature exceeds 25°C. Do not apply within 1 hour of rain.

Restrictions:

Grazing: Do not feed treated alfalfa to beef cattle within 90 days of harvest. Do not allow beef cattle to graze treated cereals, flax, lentils, pasture or rangeland within 1 day of application. Dairy cattle must not be fed or grazed on Decis treated crops.

Storage: Do not store below freezing. Do not store near feed, food, seeds or fertilizer. Keep away from heat, sparks and open flames. If stored for 1 year or longer, shake well before using.

Others: Observe a 16 yard (15 m) buffer zone from environmentally sensitive areas (for example, wetlands, sloughs, rivers, houses, farm buildings) when applying by ground equipment and a 110 yard (100m) buffer zone when applied by air. The number of aerial applications permitted varies with crop. Consult container label. In soils with high organic content (muck soils), Decis 5 EC should be applied only once per season, prior to August 1, and at rates of no more than 80 mL/acre.

Decis Flowable

CROPS	INSECT
Canola (rapeseed), mustard	Flea beetle
Wheat, barley, oats, lentils, flax	Grasshoppers
Rangeland, pastures, roadside	Grasshoppers

Precautions:

Decis is of high mammalian toxicity and is a severe eye and skin irritant. Avoid contacting or breathing spray mist. Very toxic to aquatic organisms and fish. Toxic to bees. Avoid spraying when bees are foraging. Wear protective clothing, including goggles and respirator, when handling or spraying. Do not contaminate or store near feed or foodstuffs. Wash thoroughly after using Decis.

Hazard Rating:

☠ Danger Poison

Diazinon

Company:

United Agri Products (Diazinon 500E - PCP# 11889; Diazinon 50W - PCP# 19576)

IPCO (Diazinon 50 EC PCP# 27538)

Formulations:

Diazinon 500E – 50% diazinon formulated as a liquid. Container size: 4 L, 10L

Diazinon 50 EC – 50% diazinon formulated as an emulsifiable concentrate. Container size: 10 L

Diazinon 50W – 50% diazinon formulated as a wettable powder. Container size: 2 kg

Different trade names refer to different companies. Note that products may have different label registrations and recommendations. Check the label for more information.

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	Aphids, flea beetles, Colorado potato beetles, leafhoppers
Corn, peas, beans	Seedcorn maggot
Beans (field)	Aphids
Rangeland, pasture, ditch banks, roadsides, fence rows, wasteland.	Grasshoppers

Application:

Apply Diazinon 50W by ground only.

How it Works:

Diazinon is an organophosphate insecticide with systemic and contact activity.

Effects of Weather:

Diazinon 50 EC/500E and 50W work most effectively if the temperature is 20°C or more or when temperatures will reach or exceed this minimum.

Restrictions:

Beneficial insects: Avoid spraying open flowers. Highly toxic to bees exposed to direct treatment or to residues on crops.

Environment: Do not contaminate any body of water, waterway or water source. Toxic to birds, fish, and wildlife, including waterfowl.

Precautions:

May be fatal if swallowed. Avoid contact with skin and inhalation of spray mist. Wear an approved respirator. Wash thoroughly after handling and before eating, drinking and smoking.

Hazard Rating:

☠ Danger Poison

Dibrom

Company:

United Agri Products PCP# 7442

Formulations:

864 g/L naled formulated as an emulsifiable concentrate.

Container size – 4 x 3.78 L jugs per case and 2 x 9.46 L jugs per case

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, Clover, Vetch	Aphids, loopers, leafhoppers, Lygus bugs
Potatoes	Flea beetles, Colorado potato beetles, leafhoppers
Rangeland, field areas and pastures.	Grasshoppers

Application:

Apply Dibrom by ground or air. Use designated amounts in full volumes of water. For ground application use 100 – 300 L of water per hectare. For aerial use 11-22 L of water per hectare unless otherwise stated.

How it Works:

Dibrom is an organophosphate insecticide. It acts as a contact and stomach poison.

Effects of Weather:

Do not apply Dibrom when air temperature is greater than 32°C.

Restrictions:

Environment: Do not contaminate any body of water, waterway or water source. Dibrom is highly to moderately toxic to birds, aquatic animals and other wildlife.

Precautions:

Concentrate may cause skin damage. Do not get on skin, eyes or clothing. Use waterproof gloves and face shield or goggles when handling concentrate. Harmful if swallowed. Avoid breathing spray mist.

Avoid contamination of feed, foodstuffs and drinking water.

Highly toxic to bees, avoid application during periods of bee activity.

Hazard Rating:

☠ Danger Poison

Dipel 2X DF

Company:

Valent BioSciences PCP# 26508

Formulation:

32 BIU/kg of *Bacillus thuringiensis subsp. Kurstaki* formulated as a dry flowable

Insects Controlled and Registered Crops:

CROP	INSECT
Sunflower	Sunflower moth
Timothy	Essex (European) skipper
Corn (Sweet and Field)	European corn borer larvae

Application:

Treat when larvae are young (early instars) before the crop is damaged. A spreader sticker such as Triton B1956 should be used to give thorough foliage coverage.

How it Works:

A biological stomach insecticide resulting in the larvae ceasing to eat in a few hours, with death usually occurring within 1-3 days.

Restrictions:

Storage: Store at temperatures between 0° and 25°C (cooler temperatures preferable).

Others: Do not allow dilute spray to stand in tank for more than 12 hours. Use product within 24 months of date of manufacture if stored at cool temperatures. Final spray solution for Bt should have a pH of 5-7.

Precautions:

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. In case of contact with eyes or skin, immediately flush eyes or skin with plenty of water.

Hazard Rating:

☠ Caution Poison

Eco Bran

Company:

Peacock Industries PCP# 25815

Formulation:

Wheat bran infused with carbaryl insecticide

Container size – 20 kg bag; 1kg bottle

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, clover, corn (field and sweet), beans, oats, rye, wheat, barley, canola, pastures, rangelands, forage grasses, field borders, headlands, rights-of-way, roadsides, wastelands	Grasshoppers

Preharvest Intervals and Livestock Re-entry Periods:

Crop	Preharvest Interval/ Livestock re-entry period
Corn (field and sweet)	1
Alfalfa, clover	2
Beans	5
Oats, rye, wheat	14
Barley	28
Canola	Treat only seedlings
Field borders, headlands, rights-of-way, roadsides, wastelands	0
Entry of beef cattle or other livestock to pastures, rangelands or forage grasses	1
Entry of dairy cattle to pastures or rangelands, harvest of forage crops	2

Application:

For ground application only. Do not apply by air.

Broadcast application may be made with spreaders, hand applicators, or by hand. Use gloves and wash thoroughly following application. More information on application and applicators can be found at: http://www.grasshoppercontrol.com

Restrictions:

Do not apply within 50m of sloughs, ponds, streams, dugouts or open water. Apply when winds are between 3-8 kph and do not favour drift.

Presence of product on flowering crops such as alfalfa and clover will not harm foraging honey or leafcutter bees.

May be used in pastures while beef cattle are grazing.

Precautions:

Harmful if inhaled or swallowed. Avoid breathing dust or vapour from bait. Use only in well-ventilated areas. May cause eye irritation. Avoid contact with eyes and skin. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat, sparks and open flame.

Fulfill 50WG

Company:

Syngenta Crop Protection Canada, Inc. PCP # 27274

Formulation:

50% pymetrozine formulated as a wettable granule

Insects Controlled and Registered Crop:

CROP	INSECT
Potatoes	Aphids including: Green peach, Potato, Foxglove, Buckthorn

Application:

Apply by ground only. Apply Fulfill 50WG to plant foliage. Thorough spray coverage is essential for best performance. Apply Fulfill 50 WG with sufficient water (minimum of 100 L/ha) to ensure good coverage of all plant surfaces. Higher water volumes will generally result in better coverage, especially under adverse conditions (hot, dry), where a dense plant canopy exists and/or aphid infestations are high. One additional application may be needed to control persistent aphid populations. Allow a minimum of 7 days between applications.

The use of a non-ionic adjuvant is recommended to improve the performahce of Fulfill 50WG insecticide under drought stress conditions.

How it Works:

Fulfill 50WG is a systemic insecticide and works primarily by ingestion and but has some contact activity. Affected aphids stop feeding shortly after exposure, but may remain on the plant foliage until they die, which is usually within 2-4 days. Fulfill 50WG Insecticide has residual activity in the plant and will control aphids that move onto the plant after spraying.

Fulfill 50WG has shown no phytotoxicity on the varieties of potato tested when applied at the label rates.

Effects of Weather:

Fulfill 50WG Insecticide exhibits movement through the leaf surface into plant tissue and is rainfast as soon as the spray solution has dried.

Restrictions:

Storage: Store in a cool, dry, place away from food, drinks, and animal feeding stuffs. Keep in the original container tightly closed.

Others: Do not apply by air. Do not exceed 2 applications (386 g product/ha) per crop per season. Do not apply directly to aquatic systems, permanent water bodies or areas where surface water is present. Do not contaminate water when cleaning equipment or disposing of equipment wash water.

Do not apply Fulfill 50WG insecticide through chemigation.

Do not use in nurseries or in plant propagation houses, or on any plants grown for use as transplants. A re-cropping restriction of 30 days is required for all crops.

Precautions:

May cause skin sensitization reactions. Applicators and other handlers must wear personal protective equipment including, long-sleeved shirt, long pants, waterproof gloves and shoes plus socks. Do not enter or allow entry into treated areas for 12 hours. Do not use, pour, spill, or store near heat or open flame.

Hazard Rating:

Caution Poison

Furadan

Company:

Bayer CropScience PCP# 10828

Formulations:

480 g/L carbofuran formulated as a flowable.

Container sizes: Furadan 480F-4 L jug

How it Works:

Furadan is a broad spectrum, carbamate insecticide, acaricide and nematicide.

Effects of Weather:

Avoid spraying when winds are more than 15 km/hr or less than 5 km/hr. When conditions are hot and dry, set up equipment to produce larger droplets to reduce evaporation effects.

Tank Mixes:

Furadan 480F is compatible with fungicides commonly used on the same crops. Do not mix with Bordeaux and hydrated lime.

Restrictions:

Storage: Do not store below 2°C. Store separately from food and feed.

Aerial restrictions: Do not apply by air on potatoes or sunflowers. Do not spray fields of corn smaller than 12.5 acres (5 ha) by air.

Do not apply Furadan to areas where bees are actively foraging or near apiaries. Do not spray sunflowers after plants are more than 24 inches (60 cm) in height or after heads have started to form. The use of Furadan 480F may be hazardous to burrowing owls. Do not apply within 270 yards (250 m) of known burrowing owl nests. Do not make more than 2 applications per season to potatoes, corn, and sunflower.

Insects Controlled and Registered Crops:

FURADAN 480F

CROP	INSECT
Potatoes	Colorado potato beetle, potato flea beetle, potato leafhopper, tarnished plant bug
Corn (sweet, field, silage)	European corn borer
Sunflower	Sunflower beetle
Canola, mustard	Flea beetle, red turnip beetle

Application: Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage.

Precautions:

Of high acute mammalian toxicity. Highly toxic to bees, waterfowl, birds, fish and other wildlife. Poisonous by swallowing or inhalation. Do not breathe spray mist. Do not get in eyes. Wear goggles at all times. Use an approved respirator. Do not re-enter fields less than 48 hours following application, unless appropriate protective clothing is worn. (i.e. long-sleeved shirt and long pants).

Hazard Rating:

Danger Poison

Guthion/Sniper

Last date for use of this product is December 31, 2005

Company:

Bayer CropScience (Guthion - PCP# 12287)

United Agri Products (Sniper - PCP# 23337)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulations:

240 g/L azinphos-methyl formulated as a spray concentrate and 50% azinphos-methyl formulated as a wettable powder.

Container sizes-
Guthion SC / Sniper – 240E -2 x 10 L
Guthion Solupak-4 x 2 kg

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, clover	Sweet clover weevil, grasshopper*, lygus bug, alfalfa plant bug, leafhoppers, alfalfa weevil, spittlebugs
Rye	Grasshopper
Potatoes	Colorado potato beetle, tarnished plant bug*, flea beetles*, leaf hoppers*

*Guthion only

Application:

May be applied by air or ground equipment. Apply when insects exceed economic threshold levels and use sufficient water for good coverage. Use higher rates for severe infestations, dense canopy or for adult insects (grasshoppers). Refer to label for specific mixing instructions.

How it Works:

Guthion and Sniper are contact, non-systemic, organophosphate insecticides and acaricides.

Restrictions:

Storage: Store at temperatures above freezing. Keep away from heat and open flame. Keep separate from food or feed.

For grasshopper control in rye do not apply more than once per season. Do not apply to alfalfa or clover more than once per cutting at rates higher than 0.91 L/acre. Do not apply to crops in bloom or allow spray to drift toward beehives.

Precautions:

Re-entry Period: Do not re-enter treated fields without protective equipment within 2 days of application.

Guthion and Sniper are of high acute mammalian toxicity and are poisonous if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe fumes or spray mist. Wear an approved agricultural respirator, protective clothing, natural rubber gloves and goggles when mixing or applying. Wash hands, arms and face thoroughly with soap and warm water after use. Wash all contaminated clothing with soap and hot water before re-use.

Highly toxic to bees. Do not spray crops in bloom. To protect fish and wildlife do not contaminate streams, lakes or ponds.

Hazard Rating:

☠ Danger Poison

Imidan

Company:

United Agri Products PCP# 23006

Formulation:

50% phosmet formulated as a wettable powder in water soluble sachets.

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa	Alfalfa weevil, Alfalfa blotch leafminer
Potatoes	Colorado potato beetle, potato flea beetle, potato leafhopper, potato aphid

Application:

Apply by ground only.

Imidan 50-WP instapak is packaged in water soluble sachets that are to be dropped into the spray tank unopened. Do not use in low-volume, gear-type spray equipment.

How it Works:

Imidan is an organophosphate insecticide.

Restrictions:

Storage: Keep sachets dry and do not allow sachets to contact any moist surface prior to adding to spray tank. Keep water soluble sachets in the protective container and store in a cool, dry place. Do not store at temperatures above 40°C.

Precautions:

Harmful if swallowed, inhaled or absorbed through the skin. Wear protective clothing, including rubber gloves and goggles, during mixing, loading and spraying.

Imidan is toxic to fish and bees. Keep out of any body of water.

Hazard Rating:

☠ Danger Poison

Lannate

Company:

E.I. duPont Canada PCP# 10868

Formulations:

90% methomyl formulated as a water soluble powder. Container size: 24 x 225 gram water soluble bags.

Insects Controlled and Registered Crops:

CROP	INSECT
Canola	Alfalfa looper, bertha armyworm, clover cutworm, beet webworm
Flax	Bertha armyworm
Peas	Alfalfa looper, pea aphid
Wheat, oats, barley	True armyworm, thrips
Potatoes	Aphids, leafhoppers, flea beetles, variegated cutworm

Application:

May be applied to canola, flax and cereals by air or ground equipment. Ground applications only to peas and potatoes. Apply when insects exceed threshold levels using sufficient water for good coverage.

Use higher rates for mature insects, dense canopy or when infestations are heavy. Early morning or late evening sprays are recommended.

When applied by air, pilot should not assist in mixing and loading operations. Apply a minimum of 22.5 L of water per ha (9.1 L/acre) for aerial application.

How it Works:

Lannate is a carbamate insecticide that works by contact and ingestion and has some systemic action. Rapidly degraded in green, growing plants; short term residual. Rapid knockdown.

Restrictions:

Storage: Do not store below 0°C. Store in original container away from other pesticides, fertilizer, food or feed.

Others: Sprays or drift must not contact workers, other persons or animals. The area being treated must be vacated by unprotected persons.

Precautions:

Lannate is of high acute mammalian toxicity. May be fatal or cause blindness if swallowed. Poisonous if inhaled. Do not breathe vapours or spray mist. Do not get in eyes, on skin or on clothing.

Toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where run-off is likely to occur. Do not allow to drift from treated areas. Highly toxic to bees exposed to direct application. Do not apply to areas being visited by bees. Time applications to coincide with minimum bee activity.

When mixing, loading or applying Lannate, wear protective clothing, goggles and an approved respirator. Wear clean clothes daily. Wash thoroughly after handling or applying.

First Aid:

If swallowed, drink 1 to 2 glasses of water and induce vomiting. Get medical attention immediately. If inhaled, remove from exposure. If patient is not breathing, begin artificial respiration immediately. In case of contact, wash skin with plenty of soap and water. If in eyes, flush with water for 15 minutes. Remove and wash contaminated clothing before re-use. Atropine sulphate is an antidote.

Hazard Rating:

☠ Danger Poison

Lorsban/Nufos/Pyrinex/Chlorpyrifos

Company:

Dow AgroSciences (Lorsban 4E - PCP# 14879)
United Agri Products (Pyrinex 4E - PCP# 23705)
Cheminova Inc. (Nufos 4E – PCP# 25831)
IPCO (Chlorpyrifos 480EC- PCP# 27479)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulations:

480 g/L chlorpyrifos formulated as an emulsifiable concentrate. Container sizes-10 L jug, 115 L returnable container, 208 L drum.

Insects Controlled and Registered Crops:

CROP	INSECT
Barley, oats, wheat	Army, dark sided, pale western and redbacked cutworms; armyworms, grasshoppers
Wheat	Russian wheat aphid, brown wheat mite, wheat midge
Canola	Dark-sided, redbacked, variegated, pale western, and army cutworms; bertha army worm, alfalfa looper, armyworm, diamondback moth larvae, grasshoppers, lygus bug
Flax	Dark-sided, redbacked, variegated, pale western, and army cutworms, armyworm, bertha armyworm
Lentils	Pale western cutworm, grasshoppers
Sunflowers	Redbacked, pale western and army cutworms, sunflower seed weevil
Field and sweet corn	Dark sided, black and redbacked cutworms
Potatoes	Colorado potato beetle (larvae), potato flea beetle, tarnished plant bug, red-backed cutworm, black cutworm, darksided cutworm

Application:

May be applied by air or ground equipment. Ground application only for redbacked cutworm control in corn and sunflower. Ground application only for potatoes. Uniform coverage of the crop is essential in aerial applications. Apply when insects exceed economic threshold levels and use sufficient water for good coverage. Use higher rates for heavy infestations, mature insects, heavy crop canopy, or under dry soil conditions.

How it Works:

Lorsban, Pyrinex, Nufos and Chlorpyrifos are broad spectrum, non-systemic insecticides. They work by contact, ingestion and vapour action (inhalation).

Effects of Weather:

Avoid application under hot temperatures. Best results will be obtained when application is made in evening (after 7 p.m.) or morning (before 8 a.m.) – wheat midge and cutworms. Do not apply to plants under extreme drought stress or crop injury may occur.

Tank mixes:

Lorsban 4E may be tank mixed with the following herbicides:

Avenge 200-C
Banvel plus 2,4-D amine
Buctril M
MCPA ester and amine
2,4-D amine and ester

When tank mixing always add the herbicide to the spray tank, then add Lorsban 4E. If Lorsban 4E is added first, settling out may occur causing plugging of lines or nozzles.

Restrictions:

Grazing: Treated cereals grown for cover crop should not be used for human or animal consumption within 60 days of harvest.

Storage: Combustible. Do not store near heat or flame. Do not store with food, feed, drugs or clothing.

Others: Do not use more than 9 applications on potatoes per season.

Do not use more than 5 applications or more than a total of 2.162 L/acre on wheat, barley and oats.

Precautions:

Lorsban 4E, Pyrinex 480 EC, Nufos 4E and Chlorpyrifos 480EC have a high acute mammalian toxicity. Very toxic to bees, fish, birds, aquatic organisms and other wildlife. May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. May cause skin or eye irritation. Wear protective clothing, impervious gloves and goggles. Wash thoroughly with soap and water after handling and applying. Immediately remove contaminated clothing and wash before re-use. Do not apply or allow to drift on to workers or other persons.

Do not apply directly to water or where runoff could occur to adjacent aquatic sites. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Avoid use when bees are actively foraging

Hazard Rating:

☠ Danger Poison

Malathion

Company:

United Agri Products (Malathion 500E - PCP# 4709)

IPCO (Malathion 500 - PCP# 5821)

Different companies produce malathion. Note differences in label registrations and recommendations. Check your label for more information.

Formulations:

500 g/L malathion formulated as an emulsifiable concentrate.

Application:

May be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels using sufficient water for good coverage. Use higher rates for heavy infestations, dense canopy or mature stages of insects.

Insects Controlled and Registered Crops:

CROP OR STRUCTURE	INSECT
Alfalfa, clover	Grasshopper (young), aphid, lygus bug, alfalfa weevil larvae, leafhoppers, alfalfa blotch leafminer, spider mites
Canola	Flea beetle, diamondback moth, grasshopper (young)
Wheat, barley, oats, rye	Grasshopper (young), aphid, true armyworm
Potatoes	Colorado potato beetle, leafhopper, aphid, spider mites
Canaryseed	Aphids
Sweet clover	Sweet clover weevil
Flax, lentils, hay, pasture	Grasshopper (young)
Corn (grain, forage)	Earworms, European corn borers
Beans, peas	Aphids, leafhoppers, spider mites
Empty grain bins, grain elevators, grain box cars, flour mills	Confused flour beetles, flat grain beetles, granary weevils, grain mites, Indian meal moths, lesser grain borers, red flour beetle, rice weevils, rusty grain beetles, sawtoothed grain beetles

How it Works:

Malathion is a non-systemic, contact, organophosphate insecticide and acaricide of brief to moderate persistence. Generally non-phytotoxic.

Effects of Weather:

For best results apply when daytime temperatures are above 20°C.

Restrictions:

Grazing: When spraying forages and pastures, cattle should be removed and returned after spraying.

Storage: Do not store near food or feed. Store in a cool dry place but not below -10°C. Protect from heat.

Others: Do not apply to any plant in bloom. Apply to crops when bees are absent from field. Avoid contact with automobile paint and wash immediately if exposure occurs.

Precautions:

Malathion has a low acute mammalian toxicity. Highly toxic to bees and fish. Wear protective clothing to reduce skin and eye exposure.

Hazard Rating:

Caution Poison

Matador

Company:

Syngenta Crop Protection Canada, Inc. PCP# 24984

Formulations:

120g/L cyhalothrin-lambda formulated as an emulsifiable concentrate

Container size - 4 x 3.78 L.

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	Colorado potato beetle, potato flea beetle, potato leafhopper, tuber flea beetle, Lygus bugs
Canola, mustard	Crucifer flea beetle, grasshoppers, Lygus bug, cabbage seedpod weevil
Sunflower	Sunflower beetle
Wheat, barley, oats	Grasshoppers
Unimproved pasture, summerfallow, alfalfa	Grasshoppers
Flax	Grasshoppers
Alfalfa	Alfalfa weevil, Lygus bugs, pea aphid, potato leafhopper
Corn	European corn borer, corn earworm, fall armyworm, cutworms

Application:

Aerial: Canola, mustard, sunflower, flax, alfalfa, unimproved pasture, summerfallow - Do not make more than 1 application at the 33.2 ml/ac (83 ml/ha) rate per year.

Ground: Corn, potatoes, wheat, barley, oats – Do not make more than 2 applications at the 33.2 ml/ac (83 ml/ha) rate per year.

Timing:

For potato insects, timing of application should be based on the presence of vulnerable pest developmental stages and significant populations as determined by local monitoring.

For sunflower beetles, use the high rate to control adults.

For flea beetles, to prevent migration of overwintering adults throughout the field, spray a 15 m strip around the field at the first sign of flea beetle feeding.

For grasshoppers, apply the low rate when grasshoppers are up to the 3rd nymphal stage (up to 1 cm in length) or when insect numbers are low. Apply the high rate when insects are larger, up to but not including, winged adults or when insect numbers are high.

How it Works:

Matador is a synthetic pyrethroid insecticide. It is a fast acting stomach and contact insecticide effective against a broad spectrum of foliar pests. It has no fumigant or systemic activity.

Effects of Weather:

For best results, apply Matador 120 EC during the early morning before temperatures rise, and during the evening, past the heat of the day.

Tank Mixes:

Matador may be tank mixed with Horizon herbicide.

Restrictions:

Grazing: Do not graze or feed lactating dairy animals on treated green cereal forage or treated pasture. Do not apply within 14 days of livestock foraging.

Storage: Store above 0°C. Storage below 0°C will not impair the effectiveness of Matador 120 EC, however, following such storage, agitate well before use.

Others: Allow a 7-day interval between applications. Do not apply within 15m of productive fisheries, water or waterfowl habitat.

Precautions:

Matador has potential for skin and eye irritation. Avoid splashing in eyes or on skin, particularly the face. If hands are contaminated, wash with soap and water before touching other areas of skin. This product is very toxic to bees. Avoid spraying when bees are foraging. Spray deposits should be dry before bees commence foraging in treated crops.

Hazard Rating:

☠ Danger Poison

Methoxychlor

Use of this product will not be allowed after 2005

Company:

United Agri Products PCP# 13727

Formulation:

240 g/L methoxychlor formulated as an emulsifiable concentrate.

Insects Controlled and Registered Crops:

CROP	INSECT
Beans	Leafhoppers
Potatoes	Blister beetles, Colorado potato beetles, flea beetles, leafhoppers
Alfalfa, Clover	Alfalfa weevil, armyworms, sweet clover weevil, leafhoppers, grasshoppers
Wheat, oats, barley, rye, corn	Armyworms

Application:

Ground application only

Apply in sufficient water to obtain thorough uniform coverage (usually 1000 L of water is required to cover 1 ha). Use lower rate when crop is young or infestation is light. Use higher rate when crop is more advanced or infestation is severe.

How it Works:

Methoxychlor is an organochlorine insecticide effective against a wide range of insect pests.

Restrictions:

Storage: Store away from food or feed.

Others: Do not treat alfalfa or clover during bloom. Protect from freezing.

Precautions:

Avoid inhaling spray mist. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and food. Wash after using.

Methoxychlor is very highly toxic to fish and aquatic invertebrates. Avoid contamination of ponds, lakes, streams, and other bodies of water that contain fish life or which may be used for irrigation or domestic purposes.

Monitor

Company:

Bayer CropScience, United Agri Products PCP# 12287

Formulations:

480 g/L methamidophos formulated as a liquid. Container size-10 L pail.

Insects Controlled and Registered Crops:

CROP	INSECT
Canola	Bertha armyworm, grasshoppers
Potatoes	Colorado potato beetle, potato flea beetle, potato leafhopper, aphids

Application:

May be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels using sufficient water for good coverage. Use higher rates for mature stages of insects, dense canopy or heavy infestations.

How it Works:

A broad spectrum organophosphorous insecticide and acaricide which works by contact and systemic action. Contact effectiveness may persist for 7 to 21 days.

Restrictions:

Storage: Store in a cool, dry place but not below -10°C. Do not store near food or feed. Protect from heat.

Others: Only 2 applications per season may be made on canola for bertha armyworm and grasshoppers. Highly toxic to bees exposed to direct treatment or residues on crops. Avoid use during flowering and pollination periods. Do not contaminate any bodies of water.

Precautions:

Monitor is of high acute mammalian toxicity and is extremely toxic to fish, wildlife and bees. Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Wear an approved respirator, natural rubber gloves, protective clothing and goggles. Wash hands, arms and face with soap and water after handling and applying. Wash all contaminated clothing with soap and hot water before re-use. Keep unprotected persons out of operating or spray areas or where drift could occur.

Do not re-enter treated areas until drifting insecticide and volatile residues have dissipated. Do not store or transport with feed or food.

Hazard Rating:

☠ Danger Poison

Orthene

Company:

Distributed by United Agri Products PCP#14225

Formulation:

75% acephate as a water soluble powder. Container size-case of 12 x 1.5 kg

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	green peach aphid, potato aphid, potato flea beetle, potato leafhopper, tarnished plant bug

Preharvest Interval

Do not apply within 21 days of harvest.

Application:

Apply with conventional ground equipment only. Do not apply by air.

Apply only when insects exceed economic thresholds.

Use higher rate only for heavy infestations.

How it Works:

Acephate is an organophosphate systemic insecticide that works through contact and as a stomach poison.

Effects of Weather:

Do not apply if rainfall is expected within 48 hours after application. Treatment areas should not be irrigated for at least 48 hours after application.

Restrictions:

Storage: Store in cool, dry place, in the original container away from food or feed. Protect from excessive heat.

Environment: Orthene has the potential to leach through soils to ground water. It is recommended that Orthene is not used on coarse textured soils or in areas where the water table may be high.

Do not feed foliage to livestock or allow animals to graze on treated areas.

Do not make more than 4 applications per season.

Others: Orthene is not registered in the United States. Therefore Orthene should not be used on any produce destined for markets in the United States.

Precautions:

Orthene is toxic to birds and mammals.

Orthene is toxic to aquatic invertebrates. Do not apply where runoff or drift is likely to occur.

Orthene is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging in the treatment area. Beekeepers should be warned to protect bees from treated areas for one week after treatment.

First Aid:

If swallowed, induce vomiting and obtain medical attention or call a poison control centre immediately. In case of contact with skin, wash with soap and water. If in eyes, flush with water. See a physician if eye irritation persists. Atropine is an antidote.

Hazard Rating:

☠ Caution Poison

Pounce

Company:

Distributed by United Agri Products PCP# 16565

Formulations:

384 g/L permethrin formulated as an emulsifiable concentrate. Container size-1 L (12 x 1L).

Insects Controlled and Registered Crops:

CROP	INSECT
Cereals, canola, corn, flax, lentils, peas, potatoes, sunflowers,	Redbacked cutworm, Pale Western cutworm European corn borer, corn earworm, Fall armyworm
Potatoes	Colorado potato beetle, potato flea beetle, potato leafhopper, tarnished plant bug, variegated cutworm

Application:

Apply Pounce by ground only. Apply when insects exceed economic threshold numbers and use sufficient water for good coverage. Use higher rates for heavy infestations, adult insects and dense foliage. For cutworm control application should be made under warm, moist conditions in the evening or at night. Use high rates if larvae are near maturity or soil conditions are dry. Do not disturb soil surface for five days after treatment.

How it Works:

Pounce is a synthetic pyrethroid insecticide. It is a stomach and contact insecticide with no systemic or fumigant effects.

Restrictions:

Grazing: Cover crops or crops treated with Pounce should not be used as a green feed for animals.

Storage: Store above 0°C.

Others: Observe a 16 yard (15 m) setback distance for ground and 110 yard (100 m) setback distance by air near water bodies or other sensitive areas.

Precautions:

Pounce is of low acute mammalian toxicity. Both are very toxic to bees and fish. Do not contaminate ponds, lakes or streams while filling sprayer or spraying. Avoid spraying when bees are foraging.

Hazard Rating:

▽ Caution Poison

Ripcord

Company:

BASF Canada PCP#15738

Formulation:

407 g/L cypermethrin formulated as an emulsifiable concentrate.

Container size-1 L

Insects Controlled and Registered Crops:

CROP	INSECT
Wheat, barley	Grasshoppers, cutworms
Canola	Grasshopper, flea beetle, bertha armyworm
Roadsides, headlands, summerfallow	Grasshopper
Sunflower	Sunflower beetle, sunflower seed weevil
Corn	European corn borer, cutworms, corn earworm
Potatoes	Colorado potato beetle, flea beetle, leafhopper, tarnished plant bug, cutworms

Application:

Ripcord may be applied by ground application only for control of grasshoppers on wheat, barley, roadsides, headlands and canola; for flea beetle control on canola; and for control of cutworms.

Ripcord may be applied by ground or air for bertha armyworm, sunflower beetle, sunflower seed weevil, corn earworm, European corn borer and Colorado potato beetle, flea beetle, leafhoppers and tarnished plant bug on potatoes. Only 1 aerial application is permitted per season for bertha armyworm on canola and sunflower beetle and seed weevil on sunflower, and 2 applications per season for corn and potatoes.

Observe a 16 yard (15 m) setback from water bodies and other sensitive areas when applying Ripcord with ground application. Observe a 110 yard (100 m) setback from water bodies and sensitive areas when applying by air.

Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage. Use higher rates for mature insect stages (grasshoppers) or severe infestations.

How it Works:

A pyrethroid insecticide that works as a contact and stomach poison.

Effects of Weather:

Activity of Ripcord on grasshoppers is reduced as soil temperature increases. Application for grasshopper control should be made at temperatures below 25°C. Spraying for grasshoppers should be delayed until evening if daytime temperatures are above 25°C.

Restrictions:

Grazing: Treated crops must not be grazed or cut for hay.

Storage: Keep in original container during storage. Do not contaminate or store near foodstuffs.

Precautions:

Ripcord is of low to moderate acute mammalian toxicity and very toxic to bees, fish, and aquatic organisms. Harmful or fatal if swallowed. May be harmful if absorbed through skin.

Severely irritating to eyes. Causes skin irritation and sensitization. Wear long-sleeved protective clothing and gloves when handling or applying. Wear face shield or goggles when mixing. Do not apply where streams, lakes and ponds may be contaminated. Avoid spraying when bees are foraging.

Hazard Rating:

☠ Caution Poison

Sevin XLR / Sevin XLR Plus

Company:

Bayer CropScience
PCP#: 27876 (Sevin XLR); 19531 (Sevin XLR Plus)

Formulation:

466 g/L carbaryl formulated as a liquid suspension (Sevin XLR)

480 g/L carbaryl formulated as a liquid suspension (Sevin XLR Plus).

Container size -10 L.

Insects Controlled and Registered Crops:

CROP	INSECT
Canola	Flea beetle
Alfalfa, clover	Grasshopper, blister beetle, Leafhoppers, alfalfa caterpillar, armyworm
Barley, oats, rye, wheat	True armyworm, grasshopper
Ditchbanks, field borders, forage grasses, headlands, pastures, rangelands, rights-of-way, wastelands	Grasshopper
Corn (field and sweet)	European corn borer, corn earworm, fall armyworm, grasshopper
Potatoes	Colorado potato beetle, flea beetle, leafhopper, European corn borer, climbing cutworms
Beans	Leafhoppers, lygus bugs, climbing cutworms

Application:

May be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage. Use lower rates on young plants and early stages of insects and higher rates on mature plants and advanced stages of insects, or mature insects.

Sevin XLR Plus applications are more resistant to wash-off when applied as a concentrated suspension. To ensure wash-off resistance is retained, apply dilutions of 1 part Sevin XLR Plus to no greater than 39 parts water. For example, if applying 1 L/ac of Sevin XLR Plus, do not exceed 40 L/ac total application volume. Applications should be made to dry foliage to maximize wash-off resistance.

How it Works:

A carbamate insecticide that works by contact (approx. 20%) and ingestion (approx. 80%). Moderate to rapid in speed of action with short to moderate residual activity (2 to 4 weeks).

Restrictions:

Storage: Do not store in areas where temperatures frequently exceed 38°C. Store in original container in a cool dry area out of reach of children and animals and away from food and feed.

Grazing: Remove cattle from area when spraying. Crops may be grazed or harvested for feed: Dairy animals-48 hours after treatment; Meat animals-24 hours after treatment.

Precautions:

Sevin is of moderate acute mammalian toxicity. This product is highly toxic to honeybees exposed to direct treatment on blooming crops or weeds. Harmful if inhaled or swallowed. Avoid contact with skin and eyes. Wear long-sleeve work clothing and change to clean clothing daily. Wash hands and face after handling. Avoid contamination of food, feed, water supplies, streams and ponds.

Hazard Rating:

◈ Warning Poison

Success 480 SC

Company:

Dow AgroSciences PCP# 26835 (Supplemental)

Formulation:

480 g/L spinosad formulated as a suspension concentrate
Container size-1L jug

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	Colorado potato beetle larvae and European corn borer larvae
Sweet corn	European corn borer

Application:

Apply Success 480 SC as a foliar spray by ground only. Do not apply by air. Apply when scouting indicates the target pest species have reached economic threshold levels. For Colorado potato beetle, target eggs at hatch or small larvae. For control of European corn borer, time the application to coincide with peak egg hatch Use higher application rate for higher pest pressure or when extended egg hatch is anticipated. If pest populations persist, a repeat application 7 to 10 days after the initial application may be necessary.

How it Works:

Success 480 SC is in the spinosyn class of insecticides. It is a contact and stomach insecticide. It is derived from the fermentation of *Saccharpolyspora spinosa*.

Effects of Weather:

This product has the potential for run-off. Do not spray immediately after a rainfall or if rain is forecast within 48 hours after application.

Restrictions:

Storage: Avoid freezing. Do not store or ship with food, feeds, drugs or clothing.

Others:
Potatoes - Do not apply more than a maximum seasonal rate of 100 ml/acre.
Sweet corn - Do not exceed 33 ml/acre per application - maximum of 2 applications per season.
Do not apply within 7 days of harvest. Do not apply within 28 days of fodder harvest or within 7 days of forage harvest.

Precautions:

Success 480 SC is highly toxic to honey bees exposed to direct spray. Avoid spraying on crops in bloom or other vegetation when bees are actively foraging. This product is toxic to other beneficial parasites and predators. Do not contaminate aquatic habitats such as ponds, lakes, rivers, streams and wetlands during application or when rinsing equipment and containers. A buffer zone of 2 metres (early season) or 1 metre (late season) is required between downwind edge of spray boom and sensitive aquatic habitats. Avoid contact with eyes, skin, and clothing. Do not enter or allow worker entry into treated areas for a period of 4 hours after application.

Hazard Rating:

▽ Caution Poison

Thimet 15G

This product will no longer be available for potatoes after December 31, 2006

Company:

BASF Canada PCP# 10532

Formulations:

15% phorate formulated as a granular.
Container size-20 kg bag.

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	Aphids, leafhoppers, reduction of potato flea beetle and wireworm damage, and early season control of Colorado potato beetle
Beans	Leafhoppers, aphids, lygus bugs, thrips, mites

Application:

Ground application at seeding time: Distribute granules evenly in furrow or granules may be banded on each side of the row at planting time. Use low rate for sandy or light soils and high rate for silt or heavy soils. Works as a systemic poison with effective initial residual activity on soil and foliar insects.

How it Works:

An organophosphate insecticide that works as a systemic poison, with effective initial residual activity on soil and foliar insects.

Restrictions:

Storage: Do not store in or around the home. Store away from food or feed. Store open bags in labelled sealed drums or heavy plastic bags.

Others: Do not use in muck soils. Do not apply later than at planting time. Will provide reduction of potato flea beetle and wireworm damage and early season potato beetle control.

Precautions:

Thimet is of high acute mammalian toxicity and is highly toxic to fish, birds and other animals. Do not allow product to contact eyes and skin. Poisonous by skin contact, inhalation or swallowing. Repeated inhalation or skin contact with Thimet 15G, other organophosphorus or carbamate insecticides may, without symptoms, progressively increase susceptibility to poisoning. Wear freshly-laundered, long-sleeved work clothing daily. Use rubber gloves when transferring from package to equipment. Sleeve cuffs should be worn over gloves to prevent granules from falling into the gloves. Rubber gloves should be washed with soap and water after each use. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. While emptying bags into equipment, pour downwind and allow as little free fall as possible. The use of a dust mask is recommended.

Hazard Rating:

☠ Danger Poison

Thiodan/Thionex

Company:

Bayer CropScience (Thiodan 4EC- PCP# 15747)

Makhteshim-Agan of North America Inc. (Thionex 4EC-PCP# 23453) distributed by United Agri Products

United Agri Products (Thionex 50W- PCP#14617)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulation:

Thiodan/Thionex-400 g/L endosulfan formulated as an emulsifiable concentrate. Container size-10 L

Thionex 50W-50% endosulfan formulated as a wettable powder. Container size-1 kg.

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Aphid, Colorado potato beetle, flea beetle, leafhopper
Sunflower	Sunflower beetle
Beans	Aphids, potato leafhoppers

Application:

May be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage.

How it Works:

Organochlorine insecticides that work as contact and stomach poisons.

Tank Mixes:

Compatible with most commonly used insecticides and fungicides, except Bordeaux mixture and hydrated lime.

Restrictions:

Grazing: Do not feed treated crop residue to livestock.

Storage: Do not store below freezing. If stored one year or longer, shake well before using.

Others: On sunflowers, do not make more than 1 application per season.

Precautions:

Thiodan/Endosulfan/Thionex is of high, acute mammalian toxicity and is toxic to bees. Highly toxic to fish. Hazardous if swallowed, inhaled or absorbed through the skin. Wear synthetic rubber gloves and approved respirator in prolonged spray-mixing and loading operations. Do not apply or allow spray drift to areas occupied by unprotected persons and animals or onto streams, lakes or ponds.

Hazard Rating:

☠ Danger Poison

Insect Control in Stored Grain

Prevention

Clean Storage Facilities. Storage and transport facilities should be cleaned thoroughly and sprayed or dusted, if needed, with a recommended insecticide before storing grain in the bin (e.g. malathion, diatomaceous earth or cyfluthrin - refer to product labels for details). Clean storage will eliminate one source of infestation. However, grain stored for long periods of time still has the potential for renewed infestations.

Dry and Cool Grain. Ideally, the grain should be dry before being put into storage, and cooled as quickly as possible. For long-term storage, producers are urged to lower the grain temperature below 15°C as soon as possible after the grain is placed in storage. At 15°C the stored product insects stop laying eggs and development stops. Aeration systems used during the night immediately after harvest should have the grain below 15°C in about 2 weeks. Grain that is not moved or aerated after harvest can remain warm enough to allow insects to survive the winter. Convection currents arising from this warm air can also promote condensation, sprouting (heating) and mould growth in unmanaged grain. These conditions are very attractive to stored product pests and support their development.

Monitoring for Insects

Regular monitoring of the stored grain is the next step in determining the presence and potential for serious infestations. Either the presence of insects or damaged kernels will give an indication of a problem. Probably the most sensitive means of detecting insects in stored grain is through the use of probe traps, available from numerous crop protection agents. Often the first indication of an infestation will be found near the top centre of a storage bin, and therefore, this is where traps should be placed. Monitoring should take place once every 7-10 days during the onset of storage (1st 60 days) and then the frequency of monitoring may be adjusted.

Identifying insects in stored grain

Correct identification of insects found in stored grain is important in determining the most appropriate control methods. Some of the insects found in stored grain feed on the grains, while others feed on fungus that may be developing in the stored grain.

Grain feeders:

Insects that feed on the grain include rusty grain beetles, red flour beetles, and saw-toothed grain beetles.

The **rusty grain beetle** is the most common stored product insect. Heavy infestations of this insect cause grain to heat and spoil.

The **red flour beetle** is the another common insect pest of stored grain in the prairies. Red flour beetles cannot feed on undamaged, dry seed with less than 12% moisture content. They prefer grain dust, broken grain and milled stocks. Red flour beetles can be controlled by moving grain, whether or not it is cold.

Saw-toothed grain beetles are more common in stored oats than in stored wheat and barley.

Fungus feeders:

Insects that feed on fungus in the grain bin include the foreign grain beetle, hairy fungus beetle, psocids, and grain mites.

Foreign grain beetles resemble the rusty grain beetle, but can be distinguished from it by club-shaped antennae. Also, when placed in a glass jar, foreign grain beetles will climb up the sides, while rusty grain beetles cannot. While foreign grain beetle is considered a fungus feeder, they will feed on grain if the moisture content is in the high end of the acceptable range (eg. 14.5% mc wheat).

Grain mites are whitish, about 0.2 to 0.5 mm long, and can be hard to see with the naked eye. About eight kinds of mites are common in farm granaries and elevators.

Psocids are soft-bodied insects, about 1 mm long, with long antennae relative to the body size.

Fungus feeding insects and mites cannot survive in dry grain. Chemical control is not necessary for fungus feeding pests in stored grain. Practices that result in the grain drying may be all that is needed to control such pests.

Information to help identify insect pests of stored grain can be found at:
http://res2.agr.ca/winnipeg/storage/pages/princ_e.htm

Control Techniques:

The Canada Grain Act states there is zero tolerance for any primary insects (those that feed on whole, sound grain) in grain delivered to elevators. Outlined below are some control techniques and when and how these techniques can be best used.

Cold Temperatures. Rusty grain beetles are cold hardy and can survive subzero temperatures. Rusty grain beetles and other stored grain insects can be killed by reducing core grain temperatures as follows:

Time Required to Kill Insects
at Various Grain Temperatures

Grain Temperature	Time required to kill insects
-5°C	12 weeks
-10°C	8 weeks
-15°C	4 weeks
-20°C	1 week

Cooling the grain, through aeration or moving the grain several times during mid-winter, should provide effective control of rusty grain beetles.

Moving Grain with Pneumatic Conveyors. Moving the grain using pneumatic conveyors (grain vacs) has also been shown to be an effective means of controlling free-living insects in stored grain.

Phostoxin

Company: Degesch America Inc. (Phostoxin)

Formulation: 55% aluminum phosphide.

Formulation	Primary Use	Container Size
Phostoxin tablets (3g each)	On the farm or country elevator	500 tablets
Phostoxin pellets (0.6g each)	In terminals	2500 pellets
Phostoxin tablets prepac	Containers	4 strips of 33 tablets to a pouch

Insects controlled: Rusty grain beetle, red flour beetle, saw-toothed grain beetle, granary weevil, Indian meal moth and yellow mealworm.

Approved for use on the following stored grains: Barley, corn, dried peas, lentils, oats, rice, rye, soybeans, sunflower seeds, triticale and wheat.

Restricted Product: The use and sale of Aluminum Phosphide (Phostoxin) is restricted to licensed pesticide applicators possessing a valid fumigation licence (Saskatchewan) or stored agricultural products license (Manitoba).

Rate and Minimum Exposure Period: Refer to labels to determine rate. For grain bins a dosage of 250-500 tablets (or 880-2560 pellets) per 100m^3 of bin space being treated (not volume of grain) is recommended. It is important to ensure that bins are relatively secure. It is not advisable to use phosphine products in bins that are leaky or not well sealed.

The following table may be used as a guide to determine the minimum length of exposure period to phostoxin at the indicated temperatures:

Temperature	Exposure Period
Below 5°C (40°F)	Do not fumigate
5°C-12°C (40°-53°F)	10 days
13°-15°C (54°-59°F)	5 days
16°-20°C (60°-68°F)*	4 days
above 20°C (68°F)	3 days

If the grain is less than 5°C then the tablets will not release the gas until the grain temperature warms up. This may result in poor control and accidental exposure to phostoxin at a later date in grain handling facilities. Very dry grain will also slow the release of the gas from the pellets. A shortened exposure period cannot be compensated for by increased dosage. Also ensure that storage is well ventilated for at least 24 hours after the required time for fumigation.

Cost: $75/500 tablets; $75/2500 pellets; $45/pouch (132 tablets/pouch)

Protect-It, Insecto

Company: Hedley Technologies Inc. (Protect-It) PCP#24259; Natural Insecto Products Inc. (Insecto) PCP#22489

Formulation: 90 percent Diatomaceous Earth (DE)

Insects controlled: Rusty grain beetle (Protect-It only), rice weevil, granary weevil, Angoumois grain moth, Mediterranean flour moth, Indian meal moth, red flour beetle and Tribolium.

Approved for use on the following stored products: Feed grains, seed, stored grains, wheat, barley, buckwheat, corn, oats, rye, flax, peas, soybeans and sorghum. Also registered for structural treatment of empty grain storage and transportation containers.

How it works: Diatomaceous earth damages the cuticle of the insect, reducing the insect's ability to retain moisture. The insect eventually dies from dehydration.

Rate, for empty storage structures: Use a dust blower or aeration fan to get diatomaceous earth into the cracks, crevices and void spaces of the structure being treated. Dust areas at a rate of 1 kg per 200 square metres (5 g/m^2).

Rate, while grain is being placed into storage: Protect-It; The application rate for Protect-It varies by crop and insect species, ranging from 100 g/tonne for control of rusty grain beetle in wheat to 1000g/tonne for red flour beetle in corn. Refer to the label for details. Insecto; Apply to grain at the time of storage at a rate of 0.5 to 1 Kg per metric ton of grain (500-1000 ppm).

Cost: Protect-It - $10.40/kg. Insecto - $8.69/kg.

Precautions: The application of DE will lower the test weight measurement of the grain, but usually not to the point of downgrading. If test weight loss is excessive, the grain can be diluted with untreated grain. DE is non-toxic to humans and animals.

Malathion Grain Protectant Dust

Company: Interprovincial Co-operative PCP# 17222

Formulation: 2% malathion

Insects controlled: confused flour beetles, flat grain beetles, granary weevil, Indian meal moth, lesser grain borer, rusty grain beetle and sawtoothed grain beetle.

Approved for use on the following stored grains: Wheat, rye, barley and oats as stored grains.

Malathion Grain Protectant can be applied to grain as it is being loaded into a bin or being turned by adding gradually at the grain auger. It can also be used to control surface infestations by applying to the grain surface and raking in to 15 cm depth of the grain. Malathion controls insects by ingestion and contact and insects must be active for it to be effective.

Rate and Cost:

Crop	Rate-g/1000 kg (tonne) grain	Cost/1000 kg grain
Wheat	415	$1.19
Rye	450	$1.29
Barley	520	$1.49
Oats	735	$2.10

Do not apply to grain within 7 days of sale.

Be aware that the Canadian Grain Commission allows only 8 ppm of malathion residues in stored grains.

Malathion 500, Malathion 500 E

Refer to labels for these products for insect and mite control in *empty* grain bins, grain elevators, grain box cars and flour mills.

Malathion 500 (IPCO)

Insect	Rate	Cost	Note
Rusty grain beetle, sawtoothed grain beetle, confused flour beetle, grain mite, granary weevil, Indian meal moth, lesser grain borer (empty grain bins)	250-300 ml/5 L of water on 100 m^2	$3.16-$3.80/ 100 m^2	May be used within 1 day of grain storage

Malathion 500E (United Agri Products)

Insect	Rate	Cost	Note
Rusty grain beetle, red flour beetle, sawtoothed grain beetle, confused flour beetle, grain mite, granary weevil, Indian meal moth, lesser grain borer, flat grain beetles, rice weevils (empty grain storage facilities)	333 ml/5L of water on 100 m^2	$4.10/ 100 m^2	Wait until spray has thoroughly dried before storing grain in treated areas.

Tempo 20 WP

Company: Bayer CropScience PCP# 25673

Formulation: 20% cyfluthrin. Tempo is a group 3 (pyrethroid) insecticide.

Application: Tempo can be used to control insects in grain storage facilities, truck beds and other areas where grain is stored *before filling these areas with grain*. Cleaning of all areas prior to use of Tempo 20 WP insecticide will increase levels of control. See the insecticide label for specific mixing instructions.

Additional information on controlling insects in stored grain can be found at: http://res2.agr.ca/winnipeg/storage/pages/smmry_e.htm

Notes

2005 Forage Crops Chart – Recommended Herbi

CROPS AND WEEDS / HERBICIDES	Weed Resistance Grouping	CROPS (S = seedling only; E = established o																			
		GRASSES																			
		Annual Ryegrass	Altai Wild Rye Grass	Bromegrass	Creeping Red Fescue	Crested Wheatgrass	Intermediate Wheatgrass	Kentucky Bluegrass	Meadow Fescue	Meadow Foxtail	Millets	Northern Wheatgrass	Orchardgrass	Pubescent Wheatgrass	Reed Canarygrass	Russian Wild Ryegrass	Slender Wheatgrass	Streambank Wheatgrass	Tall Fescue	Tall Wheatgrass	Timothy
◇ 2,4-D[8]	4	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	S	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X[8]	X
⯃ 2,4-DB[8]	4			S[4]	S[4]	S[4]	S[4]		S[4]				S[4]					S[4]	S[4]	S[4]	S
▽ ACHIEVE LIQUID	1																				
▽ ALLY	2				X[2]	X	X						X								
▽ ASSERT	2	X[2]																			
— ASSURE II	1				X[2]																
⯃ ATTAIN	4																		X[2]		
◇ AVADEX BW	8																				
◇ AVENGE	8			S[1]	S[1]	S[1]		S[1]	S[1]				S[1]		S[1]	S[1]					S
▽ BASAGRAN	6			S[2]	S[2]	S[2]				S[2]	S		S[2]								S
◇ BROMOXYNIL + MCPA ester[3]	4,6			S[2]	S[2]	S[2]	S[2]		S[2]	S[2]			S[2]		S[2]	S[2]	S[2]	S[2]	S[2]	S[2]	X
— CLETHODIM	1																				
— CURTAIL M	4																				X
▽ DICAMBA + 2,4-D	4			S	S	S	S		S	S			S	S			S	S	S	S	S
— EDGE	3																				
▽ EPTAM 8-E	8																				
— HOE-GRASS 284	1			S[4]	S[4]	S[4]	S[4]									S[4]					
▽ HOE-GRASS II	1,6			S[2]	S[2]	S[2]	S[2]									S[2]					
— KERB	15																				
▽ LONTREL	4		X	X	X	X	X	X	X	X			X		X	X	X	X	X	X[9]	X
◇ MCPA[6]	4	X[8]		X[8]	X[8]	X[8]	X[8]	X[8]	X[8]				X[8]		X[8]	X[8]	X[8]			X[8]	X
⯃ MCPB + MCPA	4		S[4]	S[4]	S[4]	S[4]	S[4]		S[4]			S[4]		S[4]	S[4]	S[4]	S[4]	S[4]	S[4]	S[4]	S
◇ ODYSSEY	2																				
⯃ PRESTIGE	2,4																		E[2]		
◇ PARDNER	6			S[2]	S[2]	S[2]	S[2]		S[2]		S		S[2]		S[2]	S[2]	S[2]			S[2]	S
▽ POAST ULTRA	1				X[2]																
▽ PRINCEP NINE-T/SIMAZINE	5																				
— PURSUIT	2																				
— REFINE EXTRA TOSS-N-GO	2			X	X	X	X	E				X	X	X			X	X	X	X	
▽ TARGET	4			X[9]	X[9]	X[9]	X[9]	E[9]	E[9]	E[9]			X[9]	E[9]			E[9]	E[9]	E[9]	E[9]	X
— TRIFLURALIN[8]	3																				
◇ TRIUMPH PLUS	1,2,4			S[4]	S[4]		S[4]					S[4]					X[4]		S[4]	S[4]	
— VELPAR L	5																				
▽ VENTURE L	1				X[2]																

Poison Hazard
See Safe Use of Herbicides, Fungicides and Insecticides Section in *2005 Guide to Crop P*

⯃ Danger ◇ Warning ▽ Caution — No warning symbol required

erbicides

ablished only; X = seedling or established)																	WEEDS																																																					
				LEGUMES								COVER CROPS					GRASSES						BROADLEAF WEEDS																																															
Tall Fescue	Tall Wheatgrass	Timothy	Western Wheatgrass	Alfalfa	Alsike Clover	Cicer Milkvetch	Red Clover	Sainfoin	Sweet Clover	Trefoil, Bird's foot	White Dutch Clover	Barley	Flax[11]	Oats	Canola	Wheat	Barnyard Grass	**Foxtail Barley**	Foxtail, Green	Foxtail, Yellow	Quackgrass	Wild Oats	**Bindweed, Field**	Bluebur	Buckwheat, Wild	Burdock	Catchfly, Night flowering	**Chickweed**	Cleavers	Cockle, Cow	Cocklebur	Dandelion	**Dock, Curled**	Flixweed	Goat's-beard	Goosefoot, Oak-leaved	Hawk's beard, Narrow-leaved	**Hemp-nettle**	Knotweed, Prostrate	Kochia	Lamb's-quarters	Lettuce, Prickly	**Mustard, Ball**	Mustard, Dog	Mustard, Hare's-ear	Mustard, Indian	Mustard, Tumble	**Mustard, Wild**	Mustard, Wormseed	Pigweed, Prostrate	Pigweed, Redroot	Pigweed, Russian	**Plantain, Common**	Purslane	Radish, Wild	Ragweed, Common	Ragweed, False	**Ragweed, Giant**	Russian thistle	Scentless Chamomile	Shepherd's Purse	Smartweed, Lady's Thumb	**Sow Thistle (Annual)**	Sow Thistle (Perennial)	Stinkweed	Sunflower, Prairie/Annual	Sweet Clover, White	**Thistle, Canada**
X[6]	X[6]	X[6]	X[6]									X				X								●							●	☐	●	●[g]			●[g]			●	●	●	●	●	●	●	●	●	●	●	●	●				●	●	●	●		●[g]			☐	●[g]			☐
S[4]	S[4]	S[4]		S	S		S			S	S	X		X		X							●		●							☐	S			●	●				●		●					●	●		●		●			●	●	●			●	S		S	●			☐
				S[1]	S[1]		S[1]	S[1]	S[1]	S[1]	S[1]	X				X	●		●	●		●																																														
												X				X								●	S			●		●				●				●		●	S		●					●		●	●								S	●	●	●	S	S	●			S
																						●			S																							●																	●			
				X[1]	S[2]	S	S[2]		S[2]	S[2]	S[2]		X		X		●		●			●																																														
X[2]																							☐[h]	●	●	●		S	●		●	●[g,h]	●[h]	●	●	●[h]				●	●	●	●	●[h]	●	●	●	●	●		●[h]		●		●	●	●	●	●[h]		●	●[h]	S	☐	●	●	●	S
				S[1]	S[1]		S[1]		S[1]	S[1]		X	X		X	X						●																																														
		S[1]		S[1]			S[1]		S[1]	S[1]		X				X[10]						●																																														
		S[2]		X[2]	S[2]		S[2]	S[2]	E[2]				X										☐					●	●		●										●							●			●			●	●	●		●	●		●	●			●			☐
S[2]	S[2]	X										X	X	X		X								●	●		●			●	●			●						●	●		●					●	●		●[f]					●			●	●[g]	●	●		☐	●	●		☐
				S									X		X		●		●	●	S	●																																														
		X										X	X	X		X									●							●[g]		●[g]						S	●							●			●	●								●[g]	●[g]	●	●	☐	●[g]	●		☐
S	S	S										X				X									●	●			●	●	●			●						●											●	●			●				●		●	●		☐		●		☐
				S[2]											X		●		●	●		●			●			●	S	●								S		●	●									●	●			●					S			S						
				S[2]		S[2]			S[2]	S[2]							●		●	●								●													●									●	●			●														
				S[4]	S[2]		S[4]	S[2]	S[4]			X	X		X	X	●		●	●		●																																														
																	●		●	●		●			●		●			●										●	●							●			●								●	●		●			●			
				E						E								●			●	●						●																																								
X	X[9]	X										X	X	X	X	X									●																																			●				☐				●
	X[6]	X[6]		S[5,6]	S[5]		S[5]					X	X	X		X								●		●					●		●	●		●		S[d]		●	●	●	●	●	●	●	●	●	●		●	●				●	●	●	☐		●[g]		☐	☐	●[g]			☐
S[4]	S[4]	S[4]	S[4]		S		S				S	X		X		X							☐										●					S			●		●					●	●		●		●		S	●	●	●			●		S	☐	●			●
				X[2]													●		●			●			S			●	●					●				S		S	S							●			●								●		●	●			●			
E[2]												X				X									●			S	●			●[g]		●[g]				S		●	●							●			●	●								●	●	●	●	S	●	●		S
	S[2]	S[2]		X[2]								X	X	X		X								●	●					●	●									●	●							●		●	●	●				●			●			●			●			
				X	X	X		X	X				X		X		●	S	●	●	S	●																																													
				E						E							●			●		●			●																●													●		●	●	●				●					●	
				X[15]											X[11]				S																						S							●			●										S	●			●			
X	X		X									X		X		X									●			●	S	●				●				●		●	●		●					●			●								●	S	●	●		S	●	●		S
E[9]	E[9]	X[9]	E[9]									X		X		X							☐		●		●		●	●				●				●	●	●	●		●					●	●	●	●					●			●		●	●	●	☐	●			☐
				S[13]	S[14]	S[14]	S[14]	S[12]	S[12]	S[14]			X[7]		X		●		●	●		●			●			●		●									●		●										●			●					●									
S[4]	S[4]		S[4]													X			●	●		●			●	●		●		●	●			●[g]				●		●	●	●	●		●	●	●	●	●		●	●	☐		●	●	●	●	●		●[g]	●			●[g]	●		☐
				E[2]																												☐					●																							●				☐				
				X			X[4]			X[4]			X		X		●		●	●		●																																													

uide to Crop Protection

ymbol required

REFER TO THE 2005 GUIDE TO CROP PROTECTION AND PRODUCT LABELS FOR COMPLETE DETAILS

						VOLUNTEER CROPS				
Sow Thistle (Annual)	Sow Thistle (Perennial)	Stinkweed	Sunflower, Prairie/Annual	Sweet Clover, White	Thistle, Canada	Cereal Grains (Wheat, Barley)	Cereal Grains (Oats)	Flax	Canola	Alsike Clover
	☐	●g			☐					
	S	●			☐					
							●			
S	S	●			S			S	●a	
		●								
						●	●			
S	☐	●	●	●	S			●	●	
		●			☐					
	☐	●	●		☐				●	
						●	●			
●	☐	●g	●		☐				●	
	☐		●		☐				●	
						●e				
						●	●			
		●								
						●	●			
	☐				●					●
☐	☐	●g			☐					
S	☐	●			●				●	
		●				●	●		●a	
●	S	●	●		S			●	●	
		●								
						●	●			
				●						●
		●							●a	
	S	●	●		S				●a	
●	☐	●			☐				●	
		●g	●		☐				●a	
	☐									
						●	●			

NOTES: CROPS

1. Underseeded only.
2. For seed production only.
3. Seedlings for seed production only. Established for seed or forage.
4. Do not graze or harvest for livestock in the year of treatment.
5. Use MCPA sodium salt on seedling forages only when underseeded to flax, oats, wheat or barley. Do not use on Flemish varieties of alfalfa.
6. For use as a spot treatment only to control red bartsia.
7. Apply the fall prior to seeding.
8. Not all products registered for crops and weeds indicated. Check product labels.
9. For forage production only.
10. Check recommendations for varietal restrictions.
11. CLEARFIELD varieties only.
12. Liquid formulations in spring only.
13. Advance 10G, Bonanza 10G, Treflan E.C. (spring only), Treflan QR5 only.
14. Treflan E.C. in spring only.
15. Seed production only. Apply to seedling stands that will be in production for three years after application and established stands that will be in production for 2 years after application.
16. May not include solin. Check product label for restrictions.

LEGEND: WEEDS

●	Controlled by rates recommended for crops. Where more than one brand is recommended, the weed may not appear on all labels.
☐	Top growth suppression only.
S	Suppression.
a	Will not control CLEARFIELD varieties.
b	Do not use the quackgrass rate in seedling sweet clover or seedling alsike clover.
d	Controlled by MCPAK only.
e	Controls volunteer wheat, suppresses volunteer barley.
f	Controls redroot pigweed only when a cereal cover crop is used.
g	Spring seedlings only.
h	Controlled only when growing rapidly.

STAGES OF BIRD'S-FOOT TREFOIL LEAF DEVELOPMENT



1 TRIFOLIATE 2 TRIFOLIATE 3 TRIFOLIATE

STAGES OF ALFALFA, RED CLOVER AND ALSIKE CLOVER LEAF DEVELOPMENT



UNIFOLIATE 1 TRIFOLIATE 2 TRIFOLIATE 3 TRIFOLIATE

For information on pasture weed control, consult the "Weed Control in Pastures and Hayfields" table in the *2005 Guide to Crop Protection*.

Pesticide Tank Mix Combinations

The tank mixes given on the product pages in the guide are limited to those listed on that product's label. The following chart is designed to allow the user to find all tank mix options available for a product.

To find mixes recommended on the product label, find the product of interest on the list to the left. Hits along that row correspond to tank mixes recommended by that product.

To find additional mixes that may be recommended by other products, find the product of interest in the top listing. Hits in that column will either be common recommendations or exclusive recommendations.

	2,4-D	2,4-DB	Accent	Achieve Liquid	Ally	Assert	Assure II	Atrazine	Attain	Avadex	Avenge	Basagran/Basagram Forte*	Bromoxynil	Bromoxynil+MCPA ester	Clethodim	Curtail M	Dicamba	Dicamba+mecoprop+MCPA	Dichlorprop/2,4-D	Dual II Magnum	DyVel	DyVel DSp	Eptam	Eradicane	Everest	Express Pack/Express TNG*	Frontline	Frontline 2,4-D	Fusion	Glyphosate	HoeGrass 284	Hoegrass II	Horizon	Liberty	Linuron	Lontrel	MCPA	Mecoprop	Muster	Odyssey	Pinnacle	Poast Ultra	Prestige	Primextra II Magnum	Puma™ Super	Pursuit	Refine Extra	Reflex	Sencor	Spectrum	Thumper	Trifluralin (liquid)	Triumph Plus	Trophy	Ultim	Unity	**Fungicides**	Bravo	Quadris	Tilt	**Insecticides**	Decis	Diazinon	Lorsban/Pyrinex	Malathion	Matador	Ripcord
2,4-D	■				X	X					X						X																														X																				
2,4-DB		■																																			X																														
Accent			■										X				X																																																		
Achieve Liquid	⊖	⊖		■					X					⊖		X			X																	⊖	⊖						X								X											⊖					
Achieve Liquid Gold																																					⊖																									⊖					
Ally	X				■		X				X																						X				⊖								X																						
Amitrol																														⊖																																					
Assert	X					■			X							X			X							X	X										⊖						X		⊖		X			X						⊖											
Assure II					X		■					⊖																											X		X																										
Atrazine								■					X				⊖			X				X											X																																
Attain				X		X			■																X								X												X																						
Avadex (liquid formulations)										■																																										X															
Avenge	X	⊖			X						■		X	X		X			X																		X										⊖				X																
Basagran/Basagran Forte*	⊖											■																													X							X																			
Bromoxynil*	⊖		X					X			X		■				⊖													X	X		X				X																														
Bromoxynil+MCPA ester*					⊖			⊖			X			■	X										X				X				X				⊖					X			X		⊖																				
Clethodim														X	■	⊖																		X		X	⊖		⊖							⊖																					
Curtail M				X		X					X					■									X								X												X		X																				
Dicamba	X		X		⊖												■													X							X												X																		
Dicamba+mecoprop+MCPA																		■															X		⊖														X																		
Dichlorprop/2,4-D*				X		X					X								■														X												X																						
Dual II Magnum								X												■																																															
DyVel																					■				X								X																																		
DyVel DSp								⊖														■			X																																										
Eptam																							■																													X															
Eradicane								X												⊖				■																																											
Escort	⊖																																																																		
Everest	⊖				⊖				X				⊖	X		X		⊖	⊖		X	X			■	⊖	X	X									⊖						X				⊖			X	⊖					⊖											
Express Pack / Express TNG*						X											⊖									■				X							⊖**								X				⊖**																		
Frontline						X																			X		■						⊖												X																						
Frontline 2,4-D						⊖																			X			■																	⊖																						
Fusion														X															■							X	⊖		⊖																												
Glyphosate*	⊖						⊖	⊖					X	⊖			X									X				■						X	⊖																														
Harmony Total																	⊖																				⊖																														
HoeGrass 284													X																		■					X																										X					
Hoegrass II																																■					⊖																									X					
Horizon	⊖				X				X				X	X		X		X	X		X												■			⊖	⊖	⊖					X				X				X			X		⊖				X		⊖				X	
K2	⊖																																																																		
Liberty															X																			■																																	
Linuron								X												⊖															■		X							⊖																							
Lontrel	⊖														X														X	X	X					■	⊖			X		X					X						X														
MCPA											X		X				X																		X		■												X																		
Muster							X																																■			X																									
Odyssey																																				X				■																											
Pinnacle							X					X																													■																										
Poast Ultra														X																				⊖		X	⊖		X			■				⊖																					
Prestige				X		X																			X								X										■		X																						
Puma™ Super	⊖				X				X					X		X			X		⊖	⊖				X	X									⊖	⊖	⊖					X		■		X			X	X			⊖		X											
Refine Extra	X					X			⊖							X	⊖																X			X	⊖										■																				
Reflex												X																																				■																			
Sencor	⊖																X	X					⊖														X												■			X															
Shotgun			⊖																																																				⊖												
Spectrum						X																			X								⊖												X					■																	
Sundance	⊖												⊖	⊖																							⊖																														
Thumper				X							X																						X												X						■																
Trifluralin (liquid)										X													X																										X			■															
Triumph Plus																																				X																	■														
Trophy				⊖		⊖																											X												X									■													
Unity	⊖																																			⊖	⊖								X											■											
Venture L		⊖			⊖																															⊖			⊖																												
Fungicides																																																																			
Bravo																																																										■	X								
Polyram DF																																																															⊖		⊖		
Quadris																																																										X	■								
Ronilan EG																																																																⊖			
Tilt	⊖												⊖	⊖																			X				⊖																							■							⊖
Insecticides																																																																			
Decis	⊖												⊖	⊖			⊖														X	X					⊖								⊖																	■					
Lorsban	⊖										⊖			⊖			⊖																				⊖																											■			
Matador																																	X																											⊖						■	

* Not all mixes registered with all products. See product page for details.

X Mix is recommended by both products.

⊖ Mix is recommended by the product on the left of the chart only.

** Only in the mix of Express TNG + Sencor/Evict + MCPA. 2,4-D not included.

Weatheradio Canada

Weatheradio is a network of radio stations operated by the weather service of Environment Canada which transmits weather information continuously over VHF-FM radio. The service operates 24 hours each day, seven days a week, providing up-to-the minute weather forecasts and reports detailed to each broadcast area.

Weatheradio is transmitted from nine locations in Saskatchewan:

Prince Albert, Stranraer, Regina Beach, Lanigan, Lloydminster and Midale on frequency 162.400;

Regina, Swift Current, Yorkton and Saskatoon at 162.550; and

North Battleford, Lake Diefenbaker and Whitewood at 162.475.

Medicine Hat at 162.550 can also be received west of Maple Creek.

Meteorological Service of Canada

General Inquiries (Saskatoon)	(306) 975-4264
Weather Information	
Broadview	(306) 696-2229
Estevan	(306) 634-2833
Hudson Bay	(306) 865-2721
Lanigan	(306) 365-3011
Lloydminster	(780) 875-7709
Medicine Hat, AB	(403) 526-0483
North Battleford	(306) 445-7000
Prince Albert	(306) 929-2114
Regina	(306) 780-5744
Saskatoon	(306) 975-4266
Swift Current	(306) 773-5599
Whitewood	(306) 735-2928
Yorkton	(306) 782-1511
Severe Weather Reporting and Weather Awareness	
Call No Charge	1-800-239-0484

The **Crop Protection Planner**, a Windows-based computer program, is available to accompany this publication. The Planner will help you in making quick, accurate decisions about chemical pest control options. Access to an IBM-compatible computer is required.

For the current pricing or further details on this software, contact the U-Learn Centre at the University of Saskatchewan:

Phone: (306) 966-5565
Fax: (306) 966-5567
Web site: www.extension.usask.ca
Look for the link under non-credit agriculture courses.

The Pest Control Products (Saskatchewan) Act and Regulations to that Act require that any person who applies a pesticide for commercial purposes must have a valid applicator's licence.

To be eligible for a licence, appropriate training is required from the Agriculture Division of SIAST:

Phone: 1-800-667-7730

Further details regarding licensing requirements can be obtained from the Inspection and Regulatory Management Branch of Saskatchewan Agriculture, Food and Rural Revitalization:

Phone: (306) 787-4662

Crop Protection Companies

Ag West Inc.
1-403-938-4849

Arvesta
www.arvesta.ca
1-866-761-9397

BASF
www.agsolutions.ca
1-877-371-2273

Bayer CropScience
www.bayercropscience.ca
1-888-283-6847

Cheminova
1-888-316-6260

Degesch America Inc.
www.degeschamerica.com
1-800-330-2525

Dow AgroSciences
www.dowagro.ca
1-800-667-3852

E.I. Dupont Canada
www.dupont.ca/ag
1-800-667-3925

Engage Agro
www.engageagro.com
1-800-900-5487

Gowan Company
www.gowanco.com
1-800-883-1844 ext.2

Hedley Technologies
www.hedleytech.com
1-888-476-4473

IPCO
www.ipco.ca
1-204-233-3461

Monsanto
www.monsanto.ca
1-800-667-4944

Norac Concepts
www.noracconcepts.com
1-613-841-2907

Nufarm
www.nufarm.ca
1-800-868-5444

Peacock Industries
www.grasshoppercontrol.com
1-306-225-4691

Syngenta
www.syngenta.ca
1-800-665-9250

United Agri Products
www.uap.ca
1-800-561-5444

Valent BioSciences
www.valent.com
1-800-898-2536

HERBICIDE ROTATION - 2005

To delay the onset of resistance, include herbicides from as many different herbicide groups as possible for each weed. Refer to page 27 for more information.

ROTATION GRASS HERBICIDES

GROUP 1
Achieve, Assure II, Clethodim, FlaxMax, Fusion, Horizon, Hoe-Grass 284, Hoe-Grass II*, Poast Ultra, Puma^{120} Super, Triumph Plus*, Venture

GROUP 11
Amitrole

GROUP 2
Accent, Adrenalin*, Assert, Everest, Odyssey, Pursuit, Sundance

GROUP 10
Liberty

GROUP 3
Edge, Trifluralin

GROUP 8
Avadex, Avenge, Eptam, Eradicane

GROUP 9
Glyphosate, PrePass*, Rustler*

ROTATION BROADLEAF HERBICIDES

GROUP 2
Adrenalin*, Ally, Assert (Wild Mustard), Express Pack*, Frontline*, Muster, Odyssey, Pinnacle, Prism, PrePass*, Pursuit, Refine Extra, Spectrum*, Sundance, Triumph Plus*, Unity*

GROUP 3
Edge, Trifluralin

GROUP 11
Amitrole

GROUP 4
2,4-D, 2,4-DB, Adrenalin*, Attain, bromoxynil+MCPA ester*, Curtail M, dicamba, dicamba+mecoprop+MCPA, dichlorprop+2,4-D, Dyvel, Dyvel DSp, Eclipse*, Express Pack*, FlaxMax, Frontline*, Lontrel, MCPA, MCPB+MCPA, Prestige, Rustler*, Shotgun*, Spectrum*, Thumper*, Trophy, and mixes containing these active ingredients

GROUP 10
Liberty

GROUP 9
Glyphosate, PrePass*, Rustler*

GROUP 8
Eptam, Eradicane

GROUP 5
Atrazine, Laddok*, Sencor, Shotgun*, Simazine

GROUP 6
Basagran, bromoxynil, bromoxynil + MCPA ester*, Hoe-Grass II*, Laddok*, Thumper*, Unity*

GROUP 7
Linuron

* Some products contain more than one active ingredient and therefore may appear in more than one group.

SASKATCHEWAN AGRICULTURE, FOOD AND RURAL REVITALIZATION

3085 Albert Street Regina, Saskatchewan CANADA S4S 0B1

www.agr.gov.sk.ca

Printed January 2005
35M ISBN 088656-59953 0164